SHORT NOTES

INLAND NESTING OF RED-BILLED GULLS IN NORTHLAND

The township of Kohukohu lies at the inland extremity of the Hokianga Harbour, where it branches into the Waihou and Mangamuka Rivers. On 12/12/65 a small colony of Red-billed Gulls (*L. scopulinus*) was discovered at Kohukohu; three nests on the abandoned Kohukohu jetty and fourteen nests on the abandoned jetty at Mill Wharf, about $1\frac{1}{2}$ miles from Kohukohu township. The nesting season was well advanced; one nest at Kohukohu jetty held two eggs, and at Mill Wharf each of three nests contained one downy chick, and each of two a feathered chick being fed by parents. Another feathered, non-flying chick, was swimming with parents in attendance.

flying chick, was swimming with parents in attendance.

In 1966 I visited the Mill Wharf colony on five occasions during the period October/December. On October 8th 23 birds were present and two were seen carrying straw. By 30th October 21 nests were finished or partly built, and from the shore five of these were seen to contain eggs. On 5th November we swam out, climbed the piles, and inspected 21 nests, of which 8 contained one egg, 10 two eggs and three three eggs. Hatching had started by 12th November and the parents were very aggressive. By 10th December there were still chicks in twelve of the original nests, and four additional late nests contained respectively one egg, two eggs, two eggs, and one egg one chick. The total Mill Wharf nesting population of gulls was therefore 25 pairs in the 1966 season.

Three pairs laid their eggs in hollows on top of a rotted girder, without any nest lining. Other nests on piles and girders were constructed of straw, small twigs and straw, or small twigs without any straw. One of the late nests was made of straw and dry seaweed, another of small twigs and seaweed. Two nests were right on top of the piles; many of the girders have fallen off or been removed, and the small shelf left is a favoured nest site. All the nests were at least three feet above the water at high tide.

A White-fronted Tern (S. striata) which was building a straw nest on top of a pile on 30th October, had laid one egg by 5th November. On 12th November the tern's egg had gone and there was a Red-billed Gull's egg in the nest. On 10th December, when the main gull nesting was nearly over, five pairs of terns were on nests, three of which were on piles and unlined. Two birds were sitting on disused gull nests, to which some additional material appeared to have been added when they were taken over by the terns.

_ R. S. COWAN

UNUSUAL NESTINGS OF NEW ZEALAND DOTTEREL

In the 1965/66 season two pairs of New Zealand Dotterel (C. obscurus) nested about 300 yards apart on sandbanks on the seafront of Mr. C. F. Duder's farm at North Road, Clevedon. On 23/11/65, No. 1 pair had three small young. Of these one was banded on 6/12/65 and another on 30/12/65. The third one was not banded.

On 6/12/65 No. 2 pair had four eggs in nest. On visiting the nest on 4/1/66 I found that an unknown person had placed three small stakes round the nest and two eggs had gone. These were later deserted and found to be bad.