## LITERATURE CITED

ANDERSON, E. 1949. Introgressive hybridization. Pp. 1-109. New York: John Wiley. ARCHEY, G.; LINDSAY, C. 1924. Notes on the birds of the Chatham Islands. Records of the Canterbury Museum 2: 187-201.

the Canterbury Museum 2: 187-201.
CROOK, J. H. 1965. The adaptive significance of avian social organisations. Pp. 181-218 in: ELLIS, P. E. (ed.) Social organisation of animal communities. Pp. 1-314. Symposia of the Zoological Society of London, No. 14.
FALLA, R. A.; SIBSON, R.B.; TURBOTT, E. G. 1970. A field guide to the birds of New Zealand (Second edition). Pp. 1-256. London: Collins.
FLEMING, C. A. 1939. Birds of the Chatham Islands. Part I. Emu 38 (4): 380-413. Part II. Emu 38 (5): 492-509.
FLEMING, C. A. 1952. Historical factors affecting the ecology of isolated areas. New Zealand Science Review 10 (6): 86-88.
FORD, E. 1964. Ecological genetics. Pp. 1-335. London: Methuen.
FORSHAW, J. M. 1973. Parrots of the World. Pp. 1-584. Melbourne: Lansdowne Press.
HARRISON, M. 1970. The Orange-fronted Parakeet (Cyanoramphus malherbi). Notornis 17 (2): 115-125.

HARRISON, M. 1970. The Orange-tronted Parakeet (cyanorange).
HARRISON, M. 1970. The Orange-tronted Parakeet (cyanorange).
HARRISON, M. 1970. The Orange-tronted Parakeet (cyanorange).
HACK, D. 1968. Ecological adaptations for breeding in birds. Pp. 1-409. London: Methuen.
MAYR, E. 1947. Ecological factors in speciation. Evolution 1 (4): 263-288.
MAYR, E. 1963. Animal species and evolution. Pp. 1-797. Cambridge, Mass.: Harvard Univ. Press.
RITCHIE, I. M. 1970. A preliminary report on a recent botanical survey of the Chatham Islands. Proceedings of the New Zealand Ecological Society 17: 52-56.
SIBLEY, C. G. 1961. Hybridization and isolating mechanisms. Pp. 69-88 in: BLAIR, W. F. (ed.) Vertebrate speciation. Pp. 1-642. Austin: Univ. of Texas Press.

R. H. Taylor,

Ecology Division, DSIR, P.O. Box 30466. Lower Hutt.

## SHORT NOTE

------ \* ------

## UNUSUAL NEST SITES OF HOUSE SPARROW AND PARADISE DUCK

On 17 November 1974, while surveying a square for the Mapping Scheme in the upper Lindis (Geordie Hills station) area, I came across two unusual nest sites. The first was that of a House Sparrow in a small matagouri bush, about 2 m from the ground, and right adjacent to, and almost touching, a nest of a blackbird. Both nests were occupied. It was in open tussock country with no large trees or buildings anywhere near, and about 2000' (710 m) above sea-level.

About 2 km away, in an adjoining square, two local men pointed out to me a small old musterer's hut, no longer in use by the station, with about a foot of 'starling straw' on the floor, in the middle of which a Paradise Duck had built her nest and apparently successfully hatched a clutch of about ten. The walls and roof were well inhabited by Starlings. Only the door was open.

PETER CHILD

10 Royal Terrace. Alexandra