

## LITERATURE CITED

- GEDDES, D. 1983. Crested grebes on frozen lakes. *Notornis* 30: 21-22.
- MARCHANT, P.J. and HIGGINS, M. (Editors). 1991. *Handbook of Australian, New Zealand and Antarctic Birds*. Oxford University Press, Melbourne. Volume 1A. pp 115-124.
- O'DONNELL, C.F.J. 1985. Lake Ellesmere: a wildlife habitat of international importance. *Fauna Survey Unit Report No. 40*. New Zealand Wildlife Service, Department of Internal Affairs, Christchurch. pp 41-42.
- O'DONNELL, C.F.J. 1988. Southern Crested Grebes on a lowland coastal lake in winter. *Notornis* 35: 75-77.
- SAGAR, P. and O'DONNELL, C.F.J. 1982. Seasonal movements and populations of the Southern Crested Grebe in Canterbury. *Notornis* 29: 143-149.
- STEWART, I.D. and WARD, J.C. 1990. Birds of Lake Alexandrina, South Canterbury. *Tane* 32: 137-148.
- WARD, J.C. and STEWART, I.D. 1990. *An approach to lake management ecology, values and conflict at Lake Alexandrina*. Centre for Resource Management, Lincoln University, Canterbury.
- WESTERSKOV, K. 1971. Distribution and numbers of the crested grebe (*Podiceps cristatus australis*) in Canterbury. *Notornis* 18: 1-29.
- WILSON, H.D. 1992. Banks Ecological Region. *Protected Natural Areas Report No. 21*. Department of Conservation, Christchurch.

**KEYWORDS:** Southern Crested Grebe, *Podiceps cristatus australis*, distribution, Lake Forsyth, land use.

A.R. RIDLEY and W.J. SULLIVAN, *Ecology and Entomology Group, P.O.Box 84, Lincoln University, Canterbury, New Zealand*

Manuscript received: 14 March 1998, revised & accepted 13 August 1998



### Channel-billed Cuckoos in New Zealand in spring 1996

Channel-billed Cuckoos (*Scythrops novaehollandiae*) range from Sulawesi in the west through the New Guinea region to the Bismarck Archipelago and northern and eastern Australia. The birds in Australia belong to the nominate race *S. n. novaehollandiae* and migrate south from the New Guinea region to subcoastal forests and woodlands in southern New South Wales, arriving in August to October (Mason & Forrester 1996). Their arrival in northern Australia during their northward migration in autumn coincides with the onset of the wet season, giving rise to such colloquial names as "storm bird", "flood bird" and "rain bird" (Readers Digest 1976). In Australia nesting birds usually parasitise currawongs and butcherbirds (Cracticidae) and crows (Corvidae) (Readers Digest 1976). Vagrants reach Tasmania, New Caledonia and New Zealand (Mason & Forrester 1996).

Until 1996 there were only two New Zealand records. The first is a bird presented to the Otago Museum by Mrs Thompson of Invercargill in December 1924 (Marples 1942). Its label gives no details of when or where the specimen was actually found




FIGURE 1 – Channel-billed Cuckoos: whole body and wing (upper and under surface), reg. no. 25346; tail upper and under surface, reg. no. 25345.

(pers. obs.), thus there is a possibility that it was brought to New Zealand already preserved. It was probably presented by “Mrs Thomson” (rather than “Thompson”), because its original label is from “Thomson & Co.” of Invercargill (pers. obs.). The second record is of a bird seen in Northland from October 1986 to January 1987, during which time it fed on the fruit of a Moreton Bay Fig (Howell 1987, Taylor *et al.* 1988, D. Ogle, pers. comm). In spring 1996 there were three additional New Zealand records (all dead birds), from Northland, Waikato and Wellington. This note documents each of the 1996 records. These specimens are held in the Museum of New Zealand (their registration numbers are given below).

1. Ocean Beach, Raglan, 14 October 1996. Reg. number 25346. Found washed up on the beach by Paul Cuming (Woolley 1996). This was the least decomposed of the birds found (Table 1.). The iris was red; tarsus and feet grey; bill dark grey, becoming whitish at the tip. The plumage and measurements (Table 1) indicate it was an adult female, i.e. head and neck grey, upper parts dark grey with dark feather ends and red iris (see Readers Digest 1976, Strahan 1994).

The primaries and secondaries were slightly worn, upper wing coverts mostly slightly worn, but some very worn. The outer four tail retrices were slightly worn but the central

pair were very worn. The remainder of the plumage (head and body) was slightly or moderately worn. No active moult was found.

The stomach of this bird was empty. A male louse (*Cuculiphilus cuculiphilus platygaster*) collected from this specimen is the first New Zealand record of this species, which is host specific to the Channel-billed Cuckoo (R. Palma pers. comm.).

2. Te Pahi, Northland 30 October 1996. Reg. number 25345. The dried out corpse was found by Department of Conservation volunteer workers in mixed pasture/shrubland (Brackenbury 1997). As with bird no.25346, the tarsus and feet were grey, and the bill dark grey becoming white at the tip. The grey head, and dark grey upper wing feathers with blackish feather ends, together with its larger bill and wing lengths (Table 1), indicate that the bird was an adult male. The head, wing and tail feathers were all slightly worn except for the eighth outer primary on the left wing which was more worn than the adjacent wing feathers. There was no active moult. All body feathers were missing.

3. Pukerua Bay, Wellington 9 November 1996. Reg. number 25344. This bird was found dead on the beach by Alan Tennyson. The remains comprised the left wing only. The wing feathers were all slightly worn (including the eight remaining primaries) and the dark grey feathers with blackish ends and wing length (Table 1) indicate it was an adult male.

All three cuckoos appear to have arrived in New Zealand at about the same time, approximately the second week of October 1996. At that time, a low pressure system of about 980 millibars located in the south Tasman Sea spread a strong westerly airflow between Australia and New Zealand. It is quite likely that the associated strong winds blew the birds off course and across the Tasman Sea during their southward migration. This agrees well with the fact that migrating Channel-billed Cuckoos reach Sydney mainly in October (Strahan 1994). The 1996 New Zealand records coincide with records of two or three vagrant Channel-billed Cuckoos on Norfolk Island, north of New Zealand, also in October 1996 (Anon 1997, Christian 1997). Since 1984 there has been a "very marked increase" in numbers of this species in the Sydney region (Larkins 1994). An increase in numbers of this species in the southern part of its range could explain why four birds have been recorded in New Zealand since 1986 compared to only one record before that time.

### ACKNOWLEDGEMENTS

We thank Paul Cuming for providing the Waikato specimen, Ricardo Palmo for details of the bird louse, Erena Barker for access to the Otago Museum specimen, Michael Hall for taking the photographs, Walter Boles and Jim Moore for helping obtain references, Barbara Lyford for typing the original draft, and Ray Pierce and Mike Rudge for helpful comments on the manuscript.

### LITERATURE CITED

- ANON. 1997. Recent sightings. Norfolk Island Nature Notes 12: 435.  
BRACKENBURY, G. 1997. "Toucan" sightings wanted. OSNZ News 82:8.  
CHRISTIAN, M. 1997. Is Norfolk a cuckoo's nest? Norfolk Island Nature Notes 12: 437-438.  
HOWELL, L. 1987. Regional Roundup - Far North. OSNZ News 34:7.  
LARKINS, D. 1994. The Channel-billed Cuckoo: behaviour at nests of Pied Currawongs. Australian Birds 28: 7-10.

TABLE 1 - Details of New Zealand Channel-billed Cuckoo specimens (all measurements are in mm).

Reg. no.	Date	<sup>1</sup> Dead for	Age	<sup>2</sup> Sex	Culmen length (feathers to bill tip)	Bill length (anterior margin of nostril to bill tip)	Wing length	Tail length	Tarsus length	Mid toe and claw length
<sup>3</sup> A2469	2 Dec 24	?	Ad	M	80.0	-	368	270	45.5	56.4
25346	14 Oct 96	5 days	Ad	F	74.2	60.5	341	267	45.5	57.7
25345	30 Oct 96	3 weeks	Ad	M	84.1	64.7	356	261	45.4	-
25344	9 Nov 96	4 weeks	Ad	M	-	-	c.350	-	-	-

<sup>1</sup>Estimated time since death.

<sup>2</sup>Based on measurements (see Mason & Forrester 1996) and an ovary present in bird 1.

<sup>3</sup>Otago Museum specimen examined in 1998 (pers. obs.).

MARPLES, B.J. 1942. Notes on cuckoos. Bull. OSNZ 2: 10-11

MASON, I.J.; FORRESTER, R.I. 1996. Geographic differentiation in the Channel-billed Cuckoo *Scythrops novaehollandiae* Latham, with description of two new subspecies from Sulawesi and the Bismarck Archipelago. Emu 96: 217-233.

READERS' DIGEST 1976. Complete Book of Australian Birds. Readers' Digest, Sydney.

STRAHAN, R. (Ed) 1994. Cuckoos, Nightbirds and Kingfishers of Australia. Angus & Robertson, Sydney.

TAYLOR, A.; HOWELL, A.V.; WILDIN, R. 1988. Classified summarised notes. Notornis 35: 306

WOOLLEY, B. 1996. Regional Roundup - Waikato. OSNZ News 81:13-14.

KEYWORDS: cuckoo, *Scythrops novaehollandiae*, distribution, migration, vagrant

Alan Tennyson, *Museum of New Zealand Te Papa Tongarewa, P.O. Box 467, Wellington, New Zealand*

Gerry Brackenbury, *Department of Conservation, P.O. Box 842, Whangarei, New Zealand*

Manuscript received 17 June 1998, Revised and accepted 15 October 1998



### Chestnut-breasted Shelducks and other wetland birds at Tupuangi, Chatham Islands

On 4 December 1997, I saw three Chestnut-breasted Shelducks (*Tadorna tadornoides*) on a lake and adjacent farmland at Tupuangi (44°15'S 176°12'W), Pitt Island, in the Chatham Islands group. These were identified from their plumage (Marchant & Higgins 1990), as one adult male and two adult females. This is the first record of this species from the Chatham Islands and the most eastern occurrence of this species.

This Australian species was first confirmed in New Zealand in 1973 (Grant 1989), although there is an earlier unconfirmed record (Fennell *et al.* 1983). The next records are of at least 56 individuals throughout New Zealand between December 1982 and May