Dr. R. A. Falla and Professor B. J. Marples—also respectively, the society's first president and secretary—who were joint editors during the extremely important period up to 1945; and to Mr. R. H. D. Stidolph, who has so admirably expanded and maintained 'New Zealand Bird Notes' since then, and who will, we hope, survive the impact of the new name to be proposed for the journal.

"In the early stages of the society in particular, the summarised notes were most important in drawing contributions from so many members. As the society and the status of New Zealand bird study moves towards maturity, it may have to be considered whether some of this accumulating material might not be better collated at longer intervals instead of being written up annually. It might be noted that the ambitious aim of the early summarised notes was to give each year an indication of the distribution and status of native and introduced birds. Subsequently, more general data on such aspects as breeding and migration accumulated, and the summarising of all this material is likely to be an increasingly difficult task with every year. Such thoughts as this indicate that the society and its journal have still perhaps not ended their growing pains.

"On the other hand, signs of maturity are the increasingly high standard of individual contributions to the bulletin; the initiation, with the ringing of the first bird on February 27, 1950, of our long-planned ringing scheme; the establishment of a Checklist Committee of certain members who have undertaken to draw up a revised check-list of New Zealand birds for the society; and, lastly, the natural growth in several centres of groups of our members who join as often as they can in field work and discussion, discussion interrupted, I may say, only by the impulse to leave again for the field. With these brief remarks I have much pleasure in welcoming you here again and in opening this meeting."

BIRDS AT WAIMUMU BUSH, SOUTHLAND.—The following birds were recorded during a walk of about 1½ miles through bush beside the Upper Waimumu Stream, Hokonui Hills, 9 miles from Gore, on 29/1/49: Three wood pigeons, 3 riflemen, 6 fantails, 4 tomtits, one silver-eye and two small flocks of brown creepers, 5 plus and 10. Grey warblers were heard on four occasions. The weather at the time was very windy, and only three bellbirds were heard. Several days later, during a calm misty morning, many bellbirds were singing gloriously and continuously.—W. A. Watters, Gore.

ADVANCE OF THE ROSELLA IN NORTH AUCKLAND.—Fleming (1944: N.Z. Bird Notes, Vol. 1, p. 60 and map p. 59) gives the range of the eastern rosella (Platycercus eximitus) in 1944 as extending north to Wellsford, and to Leigh on the east coast; there had been no records from the Kaiwaka-Maungaturoto district immediately to the north of Wellsford, or from the country bordering on the Kaipara Harbour north of The following reports which I received from more northern districts indicate the extent to which the rosella is increasing its range. On 26th November, 1949, Mrs. J. A. Goodwin wrote describing two rosellas which she had seen at Maungaturoto for some weeks previously, with a sketch of one which had been killed by a cat. Mr. R. F. Graham has given me information from the Mangapai South district, some fifteen miles in a direct line to the south of Whangarei: the parakeet was first seen here in April or May, 1949, and has since been observed regularly. On the east coast, a member of the society, Mr. D. G. McMillan, reports that the rosella has been well established at Mangawai since 1946 when he came to the district; Mr. McMillan also mentions this bird's attacks on fruit. On occasional trips along the Auckland-Whangarei main highway between 1946 and 1950 I have observed the rosella only as far north as Wellsford-Te Hana, and have received reports from others who in passing along this highway have noticed it in the latter district.-E. G. Turbott, Auckland Museum.