

SUMMARISED CLASSIFIED NOTES.

LIST OF CONTRIBUTORS.

S. I. Anderson, Miss, Herbert.	C. J. Lindsay, Wellington.
H. L. Annabell, Wanganui.	M. J. McCallum, Clevedon.
P. H. Basley, Ohope.	N. Macdonald, Miss, Howick.
M. A. E. Bayne, Miss, Dunedin.	C. W. McLatchie, Hilderthorpe.
F. R. Beatson, Dunedin.	H. R. McKenzie, Clevedon.
M. J. S. Black, Rotorua.	R. V. McLintock, Stirling.
C. D. Blomfield, Whakatane.	B. J. Marples, Dunedin.
A. A. Boulit, Wellington.	R. R. Marples, Dunedin.
G. T. Burton, Auckland.	W. P. Mead, Wanganui.
R. N. Buttle, Auckland.	K. H. Miers, Christchurch.
B. W. Campbell, Dunedin.	E. M. Moore, Mrs., Dunedin.
O. R. Cartwright, Miss, Dunedin.	L. H. Munro, Clevedon.
J. D. Clark, Opotiki.	F. Murray, Clevedon.
W. C. Clark, Christchurch.	S. J. Murray, Auckland.
L. D. R. Cresswell, Christchurch.	F. R. Nurse, Miss, Christchurch.
D. E. Crockett, Christchurch.	C. H. Parkin, Little Barrier.
J. M. Cunningham, Masterton.	K. Paviour-Smith, Miss, Dunedin.
P. S. Dale, Christchurch.	J. G. Reid, Papakura.
M. P. Daniell, Te Kauwhata.	T. M. Roberts, Whangaroa.
E. W. Dawson, Christchurch.	R. V. Roberts, Wellington.
M. A. d'Anvergne, Mrs., Maraetai.	S. C. Rutherford, Awhitu.
J. C. Davenport, Auckland.	V. M. Rutherford, Awhitu.
W. H. Davidson, Dunedin.	J. W. St. Paul, Clevedon.
E. Dear, Kopane.	R. St. Paul, Minginui.
B. F. Duder, Clevedon.	R. B. St. Paul, Clevedon.
D.N.F.C., Dunedin Nat. Field Club.	R. J. Scarlett, Christchurch.
J. S. Edwards, Auckland.	H. L. Secker, Wellington.
R. A. Falla, Wellington.	D. J. Shaw, Clevedon.
R. J. Fenton, Rev., Clevedon.	T. Shout, Raetihi.
M. P. Finlayson, Dunedin.	R. B. Sibson, Mangere.
R. E. W. Fordham, Marakopa.	J. H. Sorensen, Wellington.
K. V. Fountain, Miss, Dunedin.	R. H. D. Stidolph, Masterton.
J. Gardner, Miss, Opoho.	J. Sutherland, Brothers Island.
O. E. Gibson, New Plymouth.	I. Tily, Mrs., Dunedin.
M. B. Gill, Auckland.	G. E. Thomas, Auckland.
A. J. Goodwin, Miss, Clevedon.	C. B. Trevarthen, Auckland.
R. L. Grant, Hastings.	E. G. Turbott, Auckland.
G. W. Gummer, Auckland.	D. A. Urquhart, Karaka.
L. Gurr, Nelson.	L. E. Walker, Mrs., Dunedin.
B. D. Heather, Auckland.	A. H. Watson, Tauranga.
G. T. Hill, Auckland.	J. A. Watters, Gore.
R. S. Hill, Auckland.	J. P. C. Watt, Awanui.
A. H. Hooper, Auckland.	M. E. Webber, Miss, Christchurch.
B. Iorns, Masterton.	A. S. Wilkinson, Levin.
B. M. Jenkins, Mrs., Dunedin.	C. K. Williams, Gisborne.
M. E. Johnson, Auckland.	J. D. Wilson, Auckland.
E. B. Jones, Wellington.	L. Winkle, Mangawhai.
King's College Bird Club, Auck.	K. A. Wodzieki, Wellington.
W. Knight, Paraparaumu Beach.	J. C. Yaldwyn, Wellington.
D. N. Lilburne, Hunua.	

The white-faced heron continues to be reported more frequently in both islands, as many at 12 having been recorded near Auckland. Up to seven whimbrel were seen in the Miranda district. The first crested tern for the mainland of New Zealand was recorded from Spirits Bay.

BROWN KIWI (*Apteryx australis*).—Minginui (45 miles S.-E. of Rotorua) on Waiau Stream, 23/10/50, 1 dead, partly eaten, apparently by wild dogs; heard each night in this area.—(R.St.P.) Whakatane watershed, 8-9/3/51, six to eight heard, 2 nights.—(T.S.) Marakopa Valley, within last 12 years, three killed on road within mile of Te Anga P.O.—(R.E.W.F.) Hawdon River, Canterbury, lower flats, 3/2/51, one seen by T. P. Cooper.—(R.J.S.)

CRESTED PENGUIN (*Eudyptes pachyrhynchus*).—Houghton Bay, Wellington, 5/12/49; sick bird with damaged foot observed among rocks.—(H.L.S.)

YELLOW-EYED PENGUIN (*Megadyptes antipodes*).—Corpses, New Brighton Beach, one, 25/4/51, two 19/4/51, one 19/8/51, one 2/9/51.—(D.E.C.) Beach, near Nuggets, 30/9/50, pair going through ceremonies, bowing, caressing, etc.—(I.T. and L.E.W.)

LITTLE BLUE PENGUIN (*Eudyptula minor*).—Near Whangaroa, Jan., 51, albino seen, snow white, in contrast bill and eye appeared very dark, bird passed within few yards of yacht; seen by myself and members of crew.—(M.E.J.) Great Barrier Is., 13-22/11/50, breeding in progress, many heard at night coming ashore to their nests behind boulders and in crevices around Whagapara Harbour.—(J.C.D. and B.D.H.)

WHITE FLIPPERED PENGUIN (*E. albosignata*).—Corpses, one 8/1/51, Brooklands Beach; one, 25/4/51, Waimari Beach; one 3/3/51, five 31/3/51, South Brighton Beach.—(D.E.C.)

DABCHICK (*Polioccephalus rufopectus*).—Muriwai Beach pools, Tarawera Road group, 9/7/51, five pairs and one trio.—(K.C.B.C.) Motuopa Bay, Lake Taupo, 29/10/50, two pairs, courtship seen.—(R.H.D.S.) Tongoro Lagoon, H.B., 17/2/51, one; Runanga Lake, 17/2/51, six.—(D.B.) Kourarau, Wairarapa, pair reared 2 chicks, 1950-51; one juvenile was not seen after 25/3/51, and the other after 18/8/51.—(J.M.C.) Tomahawk Lagoon, Dunedin, 19/8/51, one, also seen three previous days by others.—(L.E.W.)

CRESTED GREBE (*Colymbus cristatus*).—Four reported to have been seen frequently on Lake Gunn; also one seen on several occasions lower end of Lake Te Anau, March, 1951.—(K.V.F.)

DIVING PETREL (*Pelecanoides urinatrix*).—Little Barrier Is., 11/11/50, trip from Leigh, several hundred, mostly in first four miles; 25/11/50, return trip, two miles from L.B., 1 flying.—(W.P.M., J.W.St.P., and H.R.McK.) Great Barrier Is., 13/11/50, trip from Auckland, not seen until we approached Coromandel and then continuously until well past The Needles.—(J.C.D. and B.D.H.) Corpses: Muriwai, 29/1/51, 1 dead in 22 miles (H.R.McK.); 8/7/51, 1 fresh male; 2/9/51, two.—(N.M.)

WHITE-FACED STORM PETREL (*Pelagodroma marina*).—Leigh-Little Barrier Is., 11/11/50, 25 plus; 25/11/50, return trip, seven.—(W.P.M., J.W.St.P., H.R.McK.) Auckland-Great Barrier Is., 13/11/50, odd birds seen as we approached the Coromandel coast; return trip, 22/11/50, only two seen.—(J.C.D. and B.D.H.) Cape Colville-Great Barrier Is., 24/12/50, four; return trip to Waiheke Is., 31/12/50, seen all the way, c. 60 on calm water at one place.—(A.J.G.) Ohariu Bay, 24/12/49, old corpse.—(H.L.S.)

CAPE PIGEON (*Daption capensis*).—Corpses: Muriwai, 12/8/51, 1, (J.C.D.); 17/8/51, 1 (R.B.S.); Piha, 19/3/51, 2, one very fresh (G.E.T. and M.B.G.), 29/9/51, one (J.C.D. and N.M.); Awhitu Peninsula, 19/8/51, two (S.C.R. and V.M.R.) Lyall Bay, Wellington, 7/7/51, corpse, culmen 30mm., wing 255mm.; Ohariu Bay, Wellington, 22/8/51, corpse, culmen 30mm., wing 270mm.—(H.L.S.)

GIANT PETREL (*Macronectes giganteus*).—Waitemata-Noise Is., 9/12/50, one.—(N.M.) Corpses: Awhitu Peninsula, 3/9/51, one (S.C.R. and V.M.R.); Lyttelton Harbour, 10/3/51, two (E.W.D., K.H.M., W.C.C.); Menzies Bay, 7/5/51, one (E.W.D., M.E.V.)

PRIONS (*Pachyptila* spp.).—There was a heavy winter mortality (1951) of all species of prions; the records of various members will be published in a separate paper.—(J.C.D.)

BROAD-BILLED PRION (*P. vittata*).—On 24/7/51, schoolboys found a prion, weak but apparently uninjured, on a street about 200 yards from Castlecliff Beach. When released after a few hours' rest, it attempted to fly out to sea, but came down inside the line of breakers and was washed ashore again; next morning it was dead. This was following very rough weather—(W.P.M.) Ohariu Bay, Wellington, 2/1/48, corpse of juvenile bird, decomposed, culmen 35mm., basal breadth 17mm. (H.L.S.) Brooklands Beach, Christchurch, 8/1/51, one dead.—(D.E.C.)

SALVIN'S PRION (*P. salvini*).—Ohariu Bay, Wellington, 22/8/51, corpse, female with minute ovaries; culmen 30mm., basal breadth 16.5mm., tail 80mm., tarsus 33mm., wing 190mm; primaries and tail feathers frayed.—(H.L.S.)

DOVE PRION (*P. desolata*).—New Brighton Beach, 22/8/51, one dead.—(D.E.C.)

FAIRY PRION (*P. turtur*).—Muriwai, 27 and 29/1/51, seven in 22 miles and 1 at Maori Bay.—(H.R.McK.) Corpses: One, 25/4/51, Waimari Beach, two 10/9/51 New Brighton Beach.—(D.E.C.)

FLESH-FOOTED SHEARWATER (*Puffinus carneipes*).—Leigh-Little Barrier Is., 11/11/50, several hundreds, mostly near Leigh, but all across; return trip, 25/11/50, one 12 miles from L.B., 100 plus in last three miles to Leigh.—(W.P.M., J.W.St.P. and H.R.McK.) Auckland-Great Barrier Is., 13/11/50, many seen until we passed Waiheke, not seen again; return trip, 22/11/50, an occasional bird sighted up to Cape Colville, from Coromandel to Waiheke continuously seen in many hundreds; seas were extremely rough.—(J.C.D. and B.D.H.) Kaiaua, Firth of Thames, 28/12/50, four plus flying past point in E.-N.-E. gale; from the number and variety seen at this spot, the Thames Estuary must have been swarming with birds owing to the severe storm.—(H.R.McK.) Waiheke, 29/4/51, 10 plus off the N.-E. corner.—(K.C.B.C.) Muriwai Beach, 29/1/51, c. 100 flying close off-shore.—(H.R.McK.) Island Bay, Wellington, 5/2/49, one bird close to shore.—(H.L.S.)

BULLER'S SHEARWATER (*P. bulleri*).—Leigh-Little Barrier Is., 11/11/50, several hundreds, mostly in first four miles; return trip, 25/11/50, none. Kaiaua, 28/12/50, c. 8 flew past point to north in E.N.E. gale, identity certain. Muriwai, 29/1/51, 1 or 2 flying just outside breakers.—(H.R.McK.) Clevedon River-Ponui Is., 26/11/50, 2, not often seen inside Ponui Is. Cape Colville-Great Barrier, 24/12/50, many.—(A.J.G.) Corpses: Muriwai, 27/1/51, two (H.R.McK.); 8/7/51, two, recent (J.C.D. and N.M.). Karekare, 14/7/51, one, fresh (N.M.) Westshore Beach, Napier, 23/11/50, corpse; Ocean Beach, 3/2/51, another.—(D.B.) Petone Beach, 23/2/47, corpse (E.B.J.) New Brighton Beach, 17/5/51, corpse (D.E.C.)

SOOTY SHEARWATER (*P. griseus*).—Kaiaua, 28/12/50, 6 plus flew past point in E.N.E. gale, silvery patch under wing seen. Corpses: Clevedon, Mataitai Beach, 1 dead, dry, almost certainly a victim of storm on 28/12/50.—(H.R.McK.) Muriwai, 8/7/51, 1 recent (J.C.D. and N.M.); 15/7/51, 1 in 23 miles (D.A.U.); 17/8/51, 1 (R.B.S.) Piha, 4/6/51, 2, one just ashore (J.C.D. and M.B.G.); 19/8/51, 1 (M.B.G. and G.E.T.)—(unusual dates to find this bird.) Rakaia Mouth, 9/1/51, Waimari Beach, 17/4/51, South Brighton Beach, 24/5/51 (D.E.C.). Birdling's Flat, Canterbury, 2/2/50, at sea a huge flock (estimated at 500,000 approx.) flying south at mid-day.—(G.G.) Thousands seen off Warrington; masses on water and some flying north, some south, but main movement south, 11/10/50—(Miss C. White, D.N.F.C.) At sea, from Moeraki, in fishing launch about 10 birds seen 1-12/1/51.—(G. J. Billing, D.N.F.C.)

SHORT-TAILED SHEARWATER (*P. tenuirostris*).—Waikato Heads 29/4/51, 1 pair of wings found, quite fresh, evidently only a day or two old.—(D.A.U.) Lyall Bay, Wellington, 25/12/50, decapitated corpse; wing 274mm., rectrices 95mm. with tips worn, tarsus 50mm.—(H.L.S.)

FLUTTERING SHEARWATER (*P. gavia*).—Auckland-Great Barrier Is., 13/11/50, many loose flocks seen from Waiheke onwards; 22/11/50, return trip, continuously seen.—(J.C.D. and B.D.H.) Kaiaua, 28/12/50, 8 plus flew past point to north in heavy E.N.E. gale.—(H.R.McK.) Waitemata-Kawau, 20/1/51, many flocks.—(N.M.) Clevedon, 1951, frequently seen outside Waiheke and Ponui Is.; 27/3/51, off Rakino Is., flocks of 500 to 1900.—(A.J.G.) Muriwai, 27-29/1/51, 3 dead, 1 was a juvenile with very thin bill; 100 plus feeding close to breakers.—(H.R.McK.) Waiheke-Motutapu Channel, 22/5/50, two flocks of c. 500 each.—(J.S.E.) Ocean Beach, Napier, 3/2/51, two corpses.—(D.B.)

ALLIED SHEARWATER (*P. assimilis*).—Muriwai, 29/1/51, 1 dead (H.R.McK.)

BLACK PETREL (*Procellaria parkinsoni*).—Little Barrier Is., Nov., 1950, a few dead about the Summit.—(J.W.St.P.) 25/11/50, at sea, three single birds at intervals, on water and also flying, thought to be this species, compared with *P. carneipes*, blacker, less light on side of bill, more pointed wings, "shoulders" not hunched.—(W.P.M., J.W.St.P. and H.R.McK.)

WHITE-HEADED PETREL (*Pterodroma lessoni*).—Corpses: Muriwai, 27,29/1/51, 3 in 22 miles (H.R.McK.); 11/3/51, 1, June, 1951, 1 (R.N.B.); 12/8/51, 2, one still alive being pecked by black-backed gulls (J.C.D.); Awhitu Peninsula, 20/5/51, 2; August, 7 (S.C.R. and V.M.R.) Piha, 29/9/51, 2 (J.C.D. and N.M.); Ohariu Bay, Wellington, 26/11/48, fresh corpse, culmen 38mm., wing 310mm.; 11/10/51, fresh corpse, culmen 35mm., wing 315mm, tail 138mm (H.L.S.) South Brighton Beach, 22/8/51 corpse (D.E.C.)

MOTTLED PETREL (*P. inexpectata*).—Muriwai, 27/1/51, 2 dead in 22 miles.—(H.R.McK.)

COOK'S PETREL (*Pt. cookii*).—Little Barrier Is., 17/11/50, trip to Summit, 2 fairly recently dead containing eggs, 1 alive with broken wing, and old remains; no extra mortality in nesting colony after severe two-day storm.—(J.W.St.P.); return to Leigh, 25/11/50, one to five miles from L.B., several thousands; 5 to 6 miles, a few odd ones; then none; at 12 miles (3 miles out of Leigh), one and three.—(J.W.St.P., W.P.M. and H.R.McK.) Miranda, 10/2/51, one seen at c. 100 yards in rough sea and easterly gale.—(H.R.McK.)

WANDERING ALBATROSS (*Diomedea exulans*).—Corpses: Muriwai, 18/5/51, one (R.N.B.); 15/7/51, one adult, a non-breeding male, in full white plumage and one immature (D.A.U. and E.G.T.); Awhitu Peninsula, two in August, 1951 (S.C.R., V.M.R., G.T.H. and R.S.H.).

ROYAL ALBATROSS (*D. epomophora*).—Corpses: Muriwai, 18/5/51, one (R.N.B.); Awhitu Peninsula, two in August, 1951. (S.C.R. & V.M.R.)

BLACK-BROWED MOLLYMAWK (*Thalassarche melanophrys*).—Corpses: Awhitu Peninsula, August, 1951, one adult and one juvenile.—(S.C.R. and V.H.R.)

BULLER'S MOLLYMAWK (*Th. bulleri*).—Island Bay, Wellington, observed near land from shore on the following dates: 30/4/48, 4/1/49, 5/3/49, 12/6/49, 24/6/49; a bird on one occasion was seen being chased by a gull.—(H.L.S.) About six miles out to sea from Moeraki, in fishing launch, six seen Jan., 1951—(C. J. Billing, D.N.F.C.)

WHITE-CAPPED MOLLYMAWK (*Th. cauta*).—Great Barrier, 21/11/50, a non-breeding female was found, almost dead, on Kaitoke Beach; it was the Bounty Is. sub-species *Th. c. salvini*.—(J.C.D.) Island Bay, Wellington, seen from shore on following dates: 15/11/47, 20/2/48, 5/3/49, 17/3/51 and 30/4/48. The bird seen on 30/4/48 had a pure white head. This bird could have been either of the usual local race *Th. c.*

salvini or referable to another race. It is noteworthy, however, that the nuchal feathers of **Th. c. salvini** may turn white with wear.—(H.L.S.) New Brighton Beach, corpses, 29/4/51, one; 17/5/51, one; 10/9/51, two.—(D.E.C.)

WHITE-THROATED SHAG (*P. melanoleucos*).—Upper Waitotara, Dec., 1950, breeding; some young just flying; others still in nests.—(H.L.A.)

GREY-HEADED MOLLYMAWK (*Th. chrysostoma*).—Muriwai, 15/7/51, one freshly dead, an immature female.—(E.G.T.)

LITTLE BLACK SHAG (*Phalacrocorax sulcirostris*).—Maraetai, Clevedon, 23/7/51, 18 on wharf.—(R.J.F.); 26/7/51, nine swimming in close party.—(D.J.S.) Hamurana Springs, L. Rotorua, 8/11/50, four seen, three of them sitting on a punt with a black shag.—(M.J.S.B.)

SPOTTED SHAG (*Stictocarbo punctatus*).—Muriwai, 27, 29/1/51, 1 alive and four dead, three were lying close together 22 miles up beach, having been shot.—(H.R.McK.) Palliser Spit, 15/7/51, one diving off shore.—(R.H.D.S.) Katiki Beach, Otago, 29/1/51, two.—(L.E.W.) Near Nugget Point, 30/9/50, numbers in vicinity; Lake Forsyth, Banks Peninsula, one, man was going to shoot it.—(I.T., L.E.W.)

GANNET (*Morris serrator*).—30/12/50, many gannets scared off the water between 1 a.m. and 3 a.m. in darkness by a launch amongst the small islands off Gt. Barrier Is.—(R.V.R.) Kawhia (Te Maika), 18/2/51, one immature corpse; 21/2/51, one immature corpse.—(N.M.) Wellington Harb., 22/9/50, first gannet noted for some time; 20/10/50, gannets seen near the Floating Dock during the nesting season when these birds are usually absent from Cook Strait area; 10/4/51, small flock diving continuously; 18/5/51, two in harbour.—(R.V.R.) Menzies Bay, Aug., 1950, two (E.W.D.). Titahi Bay, 17/2/51, one.—(E.W.D., J.C.Y.). One mature and one imm. at sea off Sth. Brighton, 10/3/51.—(E.W.D., P.S.D., W.C.C.). Menzies Bay, 7/5/51, one.—(E.W.D.) Karatane, 14/5/51, three, one diving.—(J.P.C.W.) Allan's Beach, Otago Peninsula, 6/5/51, 20; one in immature plumage.—(B.J.M.)

WHITE HERON (*Casmerodius albus*).—Whangaroa, Northland, 20/6/51, one in harbour but chased away by gulls.—(T.M.R.) Kawa Kawa Bay, Clevedon, 23/3/51, one feeding on tide flat, seen only once so was probably moving north.—(M.J.M.) Whitianga, Mercury Bay, one came early May, 1951, still present up to 17/8/51, report by Mr. M. J. Thorn.—(H.R.McK.) Lake Rerewhakaitu, south of Rotorua, 10, 13, 14/5/51, one seen at east end of lake by Mr. Baden Prentice on fishing trips; it was very quiet.—(R.St.P.) 29/8/51, three near Napier.—(J.D.W.). Near Napier, 25/4/51, three; 5/6/51, two; 19/6/51, four; 11/5/51, one reported Oputama, near Mahia Pen., and one Tokomaru Bay "quite recently" (May).—(D.B.) Woodend, 1951, two.—(W.C.C.) Waimakariri Estuary, 12/5/51, one; Christchurch Estuary, 20/8/51, one under constant attack from black-backed gulls.—(D.E.C.) Kaiapoi district, near Christchurch, four seen during the winter months of 1951. They have been kept under observation, since 4/7/51, by Rima Hopkinson, Barrie Keetley, Nola Burgess, David Beggs, Alvin Todd, Ronald Marriott and others of Form I, Kaiapoi Borough School.—(E.W.D.); two, Kaiapoi, 16/9/51 (E.W.D., F.R.N.). Merton Swamp, 9/7/51, one.—(R.M.J.) Herbert, 23/2/51 to 26/4/51, one visited a farm at intervals; it was seen feeding in a creek and roosting in macrocarpa trees; first occurrence in this locality.—(S.I.A.)

WHITE-FACED HERON (*Notophox novaehollandiae*).—Manukau Harbour (a) Puketutu Causeway, 5/8/51, 1 (R.B.S.); (b) Karaka, a flock of varying numbers present during the winter; 12/5/51, 1; 31/5/51, 6; 26/7/51, 12; 26/8/51, 7.—(D.A.U.) Clevedon, 19/2/51 to 14/7/51, one seen frequently; Miranda, 24/6/51, one; Waiuku Channel, 6/5/51, three; 27/5/51, one.—(H.R.McK. et al.) One near Studholm Junction, five

Blueskin Bay, 30/1/51; Waikouaiti River, 27/3/51, one.—(I.T.) Waitati Bay, Otago, 19/11/50, four; Waikouaiti River, Otago, 27/3/51, close view of bird, very pale in colour, very little white, legs bright yellow.—(L.E.W.)

REEF HERON (*Demigretta sacra*).—Cornwallis, Manākau Harbour, 18/8/51, seven sitting on wharf; Shelly Park, Howick, 1/9/51, six; Howick, one or two present throughout year, three seen on 27/1/51 and 21/7/51.—(N.M.) Rocky foreshore, New Plymouth to Oakura and Waitara, several.—(O.E.G.) Waimakariri Est., 18/5/51, six feeding.—(D.E.C.) Menzies Bay, 5-11/5/51, one; Christchurch Est., 18/3/51, 8/4/51, 12/8/51. (E.W.D.)

BITTERN (*Botaurus poiciloptilus*).—Minginui, May, 1951, one has returned as usual for its annual moult.—(R.St.P.) Swamp, Cape Foulwind, 1/1/51, one.—(C.J.L.) Shag River, Dunback, 10-15/1/51, two.—(F.R.B.) Merton Swamps, 21/10/50, one.—(Mrs. J. Hickman, D.N.F.C.) Stirling, May, 1949, one.—(R.V.McL.) Hakataramea Valley, about seven miles from Kurow, 14/12/50, one; near Struan, Hakataramea Valley, 15-16/2/51, one.—(W.A.W.)

ROYAL SPOONBILL (*Platalea regia*).—One feeding on Wanganui Est. for three days, 13, 14 15/3/51.—(W.P.M.) Ashley River, Nov., 1950, one seen by R. G. Frean, Christchurch.—(E.W.D.)

PARADISE DUCK (*Tadorna variegata*).—Minginui, 7/10/50, two flying high; 26/5/51, one flying high.—(R.St.P.) Aorangi, Gisborne, a pair arrived here in the autumn of 1948. In February, 1950, I counted 110 in one paddock where they remained till the end of April. In 1951, 90 to 100 stayed in the next paddock from February-April.—(C.K.W.) Kopane, Manawatu, 26/3/51, eleven in flock.—(E.D.) Lake Guyon, 3/1/51, 50 plus; Lake Christopher, 2/1/51, four pairs.—(W.C.C.) North branch of Rouburn River, 27/12/49, four; upper reaches Olivine River, near Cow Saddle, N.W. Otago, 29/12/49, two pairs; Hidden Falls River, near junction with Hollyford, 6/1/50, twelve.—(W.A.W.) Lagoon, Titan Rocks, Garvie Mts., 3/1/50, pair; Old Man Range, 6/1/50, pair. In January, 1951, for the first time on numerous visits made to the Rouburn Valley over a period of years, no paradise ducks were seen on the river flats near the Rouburn Huts.—(E.M.M.)

GREY DUCK (*Anas poicilorhynca*).—Christchurch Est., 29/7/51, 100 plus; Lake Christopher, 2/1/51, 20 plus; Lake Guyon, 3/1/51, 70 plus.—(W.C.C.) Stirling, 19/4/51, numbers assembling on pond as usual in May, with mallards and resident mallard-domestic hybrids. Not so easily frightened off as in past winters; 11/5/51, c. 60 on pond, many more under willows.—(R.V.McL.)

GREY TEAL (*A. gibberifrons*).—Gladstone, Wairapa, 4/11/50, nests, 8 eggs (hatched 5/11/50); 8 and 9 eggs (both found deserted, 1 broken egg out of nest).—(J.M.C.)

SHOVELLER (*A. rhynchotis*).—Wanstead Swamp, 13/11/49, duck with seven ducklings.—(K.W.) Lake Tuakitoto, near Kaitangata, 30/9/50.—(I.T., L.E.W.) Stirling, 1949, three pairs; 1951, none on sanctuary.—(R.V.McL.)

BLACK TEAL (*Aythya novaeseelandiae*).—Muriwai Beach pools, Ratara Road group, 9/7/51, c. 30.—(K.C.B.C.) Hamurana Stream, L. Rotarua, 8/11/50, above the tearooms, c. 62, drake predominating.—(M.J.S.B.) Lake Waikaremoana, 4/3/51, eight.—(T.S.) Tiniroto Lake, East Coast, 25/11/50, at least four seen.—(K.W.) Monowai Canal, Dec., 1950, several.—(C.B.T. and J.S.E.)

BLUE DUCK (*Hymenolaimus malacorhynchus*).—Minginui, 22, 23/10/50, Waiau Stream, 6 and 2, two pairs of 5 and 2. These young birds were the prettiest I have seen, with their ivory bills, grey-white

necks and breasts, bluish-brown backs and black eyes. The drake swims about in a position where he can see up and down stream and when you come upon them he gives a piercing whistle and swims away a little to lure you while the duck makes for the nearest patch of driftwood to hide the little ones. The two broods were only a few days apart in age so that October must be the principal hatching month for the blue duck; 18, 19, 20, 21/4/51, 3, 2, 9, 2; 8, 9, 10/5/51, 2, 10, 2.—(R.St.P.) Waikare Stream, 7/3/51, at least six.—(T.S.) Edwards River, Arthur's Pass, 6-8/1/51, one.—(E.W.D.) Lower Hidden Falls Valley, W. Otago, 6/2/51, two; two in same valley about a mile from previous locality; calls upper N. branch Routeburn late p.m. 7/2/51.—(B.W.C.)

CANADA GOOSE (*Branta canadensis*).—Lake Guyon, 3/1/51, 100 plus; Lake Tennyson, 5/1/51, 150 plus.—(W.C.C.) Washdyke Lagoon, Timaru, 16/10/50, 2.—(J.M.C.)

BLACK SWAN (*Cygnus atratus*).—Waikopu Creek, Whitford, 29/7/51, 15.—(M.A.A. and C.D.B.) Shelly Park, Howick, 3/6/51, 49; Clarke's Beach, Manukau Harb., 16/6/51, 500 plus; Kawhia, 25/2/51, 200 plus.—(N.M.) Lake Guyon, 3/1/51, two; Lake Brunner, 3/6/51, 29; Christchurch Estuary, 29/7/51, two, 12/8/51, 36 plus.—(W.C.C.) Ashley Mouth, 19/5/51, c. 20 (E.W.D., J.C.Y.), flooded paddocks, Horotane Valley, Ch-ch., 22/5/51, 12 (E.W.D., J.C.Y.), Christchurch Est., common during winter, 1951.—(E.W.D.) Lake Ellesmere, Canterbury, 22/8/50, thousands; Tomahawk Lagoon, Otago, 11/5/51, 39.—(L.E.W.) Lake Waihola, 29/8/50, 150 to 200; 19/9/50, 250 to 300.—(I.T.) Lake Tuakitoto and Lake Waihola, 30/9/50, hundreds on both lakes.—(I.T., L.E.W.)

WHITE SWAN (*C. olor*).—Lake Pupuke, Takapuna, pair recently reared six young, which is unusual.—(R.B.S.) Two, Ashley Mouth, four, Waikuku, 19/5/51.—(E.W.D., J.C.Y.) Tomahawk Lagoon, Otago, 15/7/50, 7; 18/9/50, pair seen, one on nest in rushes; 11/5/51, six.—(L.E.W.) Lake Waihola, 29/8/50, 19/9/50, two.—(I.T.)

BUSH HAWK (*Falco novaeseelandiae*).—Minginui, Nov., 1950, to June, 1951, hardly any seen up to Feb., then 1 or 2 fairly often about bush workings and up to five in one day on hunting trips. The bush hawks are mostly up the rivers and they do great damage among the pigeons each year. On one hunting trip we saw several lots of feathers where pigeons had been struck down and eaten. One day we saw three bush hawks chasing one pigeon. One followed it through the trees while the other two kept above, ready to strike if the pigeon rose above the trees, or if it crossed a clear space. I have often seen one hawk chasing a pigeon and have seen the kill, but this was an example of team work. So far as we saw, the pigeon escaped by keeping low through dense growth, but it was a very lucky one.—(R.St.P.) Hunua, Aug., 1950, one seen, third in c. 15 years.—(D.N.L.) Ngaio, 18/5/51.—(E.B.J.) Wadestown, 19/5/51.—(R.V.E.) Governor's Bush, The Hermitage, Mt. Cook, 30/6/51, presumably the same bird seen near Governor's Bush, 1/7/51; White Horse Hill, 2 miles from Hermitage, another noted on dead tree, screeched incessantly and could be heard long after observer had passed from view, 8/7/51.—(R.M.J.) Base of Kirkliston Range, near Hakataramea Station, South Canterbury, 30/1/51, one.—(W.A.W.)

HARRIER (*Circus approximans*).—Howick, one seen chased by two red-billed gulls.—(N.M.) Minginui, Nov., 1950, to June, 1951, not plentiful, one to four most days, but none for several days at times.—(R.St.P.) Taupo, 12/7/50, feeding on fresh horse corpse.—(K.W.) Twice visits paid to a bird table, Maori Hill, Dunedin, 1 and 7/7/51, this was next door to the house where I.T. reported a young harrier perched on a chimney on 23/3/50, in Notornis, Jan., 1950.—(E.M.M.)

PHEASANT (*Phasianus colchicus*).—Howick, 15/8/51, two hens chased a red-billed gull, flying about 100 feet from the ground, round and round, eventually sending it out to sea; then the pheasants dropped to the ground again and resumed feeding.—(N.M.)

BROWN QUAIL (*Synoicus ypsilophorus*).—Dargaville-Waipoua Rd., 8/2/51, 2 along roadside; Hikurangi-Whangarei Road, 12/2/51, three; Ruakaka, 13/2/51, 3.—(N.M.) Minginui, Nov., 1950 to June, 1951, one or two at odd times, one flock of nine.—(R.St.P.)

CALIFORNIAN QUAIL (*Lophortyx californicus*).—Karekare, 14/7/51, eight feeding on lawn near boardinghouse.—(N.M.) Minginui, Nov., 1950 to June, 1951, summer, up to c. 30; autumn, up to c. 400; pairing noted, Nov.; many chicks Jan.; 20/1/51, on upper Whirinaki River, a pair with brood of day-old chicks, two miles inside bush.—(R.St.P.)

NORTH ISLAND WEKA (*Gallirallus greyi*).—Tuhunga Road, off Tiniroto-Gisborne highway, 25/11/50, at least four heard in a 50-acre patch of bush.—(K.W.)

BANDED RAIL (*Hypotaenidia philippensis*).—Whangaroa, Northland, 1950-51, certainly increasing.—(T.M.R.) Huia Stream, Waitakeres, 15/5/51, c. 12 feeding in open near the mouth.—(K.C.B.C.)

SPOTLESS CRAKE (*Porzana tabuensis*).—Ngahere, Grey Valley, Feb., 1951, one.—(R.J.S.)

MARSH CRAKE (*P. pusilla*).—Mangawhai, 15/2/51, first seen in district; feeding on tidal mudbank.—(L.W.) Forest Lakes, Otaki, 5/5/51 three seen by Mr. C. McArthur.—(C.J.L.)

PUKEKO (*Porphyrio poliocephalus*).—Lyal Bay, Wellington, 21/8/48, fresh corpse; Ohariu Bay, Wellington, 13/1/51, remains found, wing 300mm.; cf. Oliver, N.Z. Birds, 1930, who gives measurements of wing as 290mm.—(H.L.S.) Lake Ellesmere, near Birdling's Flat, 5/5/51, c. 200 seen by Mr. C. Holmes.—(D.E.C.) L. Waihola, 29/8/50, over 100; 30/9/50, 60 plus.—(I.T., L.E.W.)

SOUTH ISLAND PIED OYSTERCATCHER (*Haematopus finschi*). Puketutu Causeway, Manukau Harb., 12/1/51, three seen on rocks near Island, 4/2/51, 45 plus (J.C.D.); 300 plus wintered (R.B.S.) Miranda coast, 1950-51, lowest spring count, 4/11/50, 28, rising to 10/2/51, 253 plus, 18/3/51, c. 300; and on 24/6/51, 416.—(H.R.McK. et al.) Waitotara Estuary, 11/1/51, one, a very rough day; none seen on previous and subsequent fine days.—(W.P.M.) Lake Guyon, 4/1/51, nine; Waiau River at foot of Maling Pass, 5/1/51, five; Christchurch Estuary, 29/7/51, 800 plus.—(W.C.C.) Saltwater Creek, 26/3/51, flock of c. 600 feeding on mudflats; Waimakariri Estuary, c. 450 do.—(D.E.C.) Ashley Mouth, 19/5/51, c. 1,000 (E.W.D., J.C.Y.); 22/8/51, c. 200 (E.W.D., I.D.R.C.); 16/9/51, c. 200 (E.W.D. F.R.N.). Christchurch Estuary, 8/4/51 c. 500, 12/8/51 c. 300; South Brighton Beach, 4/1/51, 57.—(E.W.D.) Heathcote Est., Chch., 19/8/50, hundreds.—(L.E.W.) Rangitara River mouth, 18/1/49, two; five days earlier 12 in flight; Opihi River, Silverstream, 22/1/49, two.—(D.N.F.C.) Kyebrun, Central Otago, 26/8/50, five.—(M.A.E.B.) Owaka Heads, 12/8/50, about 150 associated with two black and 14 pied oystercatchers, 4 godwits and 1 Caspian tern.—(K.W.) Wickliff Bay, 9/8/50, two.—(Mrs. J. Hickman, D.N.F.C.) Dredge Flats, Dart Valley, 2/2/51, three.—(B.W.C.) Pounawea, Otago, 13/1/51, 52; Doctor's Point, Otago, 18/3/51, 50 plus; Taieri Mouth, Otago, 24/3/51, 95; Mudflats, Spit, Otago Heads, 8/4/51, very large flock.—(L.E.W.) Flock on Sandy Spit, Karatane, 10-19/5/51; birds collected shellfish in piles for the sun to open while they waited near by. When the shells opened they stuck their bills into the cracks and forced the shells open. Each bird appeared to have a separate pile. Called during the night.—(J.P.C.W.) Allan's Beach, Otago Pen., largest count 100, 6/5/51.—(B.J.M.) Taieri Mouth, 29/8/50, 52; 19/9/50, 9; 11/5/51, 43; Karatane, 92, 27/3/51.—(I.T.) Stirling, at pond, 12/11/49, saw two, not often seen.—(R.V.McL.) Ocean Beach, Invercargill, 5/2/51, c. 30.—(E.W.D.) Greenhills, Bluff, 5/2/51, c. 20 (E.W.D.)

NORTH ISLAND PIED OYSTERCATCHER (*H. reischeki*).—Great Barrier, 21/11/50, a pair with nest and 1 egg and a lone bird seen on Kaitoke Beach.—(J.C.D., B.D.H.) Miranda, 1 on 1/4/51, 2 on 23/9/51.—(R.B.S.) Maketu, 14/12/50, one by itself, black with a little white about the vent.—(A.H.W.)

BLACK OYSTERCATCHER (*H. unicolor*).—Russell, 11/2/51, three resting on small island next to Moturoa.—(N.M.) Karaka, one with wintering S.I.P.O.—(D.A.U.) Miranda, 1950-51, from 1 to 3 frequently seen.—(H.R.McK. et al.) Allan's Beach, Otago Pen., one seen twice during May and June, 1951, three 30/6/51.—(B.J.M.) Akatore River mouth, 19/9/50, 3, reported to have bred.—(I.T.) Katiki Beach, Otago, 18/1/51, 4; Mudflats, Spit, Otago Heads, 8/4/51, 3.—(L.E.W.) Kaka Point 12/6/51, three.—(M.P.F.)

BLACK OYSTERCATCHER (*Haematopus* ?sp.).—Porangahau Est., 15/11/50, pair keeping territory on northern side; also present in same territory in November, 1949.—(K.W.) Waitotara Est., 27/10/50, one only; 12/1/51, three together, of which one was immature.—(W.P.M.)

TURNSTONE (*Arenaria interpres*).—Manukau Harb.: (a) Puketutu, 31/10/50, 27; 31/1/51, 38; 4/3/51, 38; 18/3/51, 10; 31/3/51, 36; 1/4/51, c. 20; 31/8/51, 1. (N.M., H.R.McK., R.B.S.); (b), Karaka, maximum c. 75 on 8/4/51; 3 on 24/5/51; 4 on 8/7/51; Clevedon, 14/7/51, 4; the record for Clevedon.—(H.R.McK.)

LESSER GOLDEN PLOVER (*Pluvialis dominicus*). — Puketutu Causeway, Manukau Harb., 4/3/51, 18; 18/3/51, 4; 31/3/51, 5; 1/4/51, 11; 12/4/51, 6.—(N.M. and R.B.S.) Karaka, maximum c. 69 on 11/3/51; last seen, 55 on 8/4/51.—(D.A.U.) Clevedon, 6/1/51, 4; 4/3/51, 6, only one well coloured.—(F.M. and H.R.McK.) North Road, Clevedon, 24/2/51, 7 on a paddock.—(B.F.D.) Waitakaruru, 10/2/51, 15 flying; 1/4/51, 17; and at Piako, 15.—(H.R.McK. et al.) Waitotara Estuary, the best count this year was six on 12/1/51, but as the birds were scattered feeding, there may have been more about.—(W.P.M.)

BANDED DOTTEREL (*Charadrius bicinctus*).—Great Barrier, 21/11/50, c. 10 pairs seen on Kaitiki Beach, at least two with nests and eggs.—(J.C.D. and B.D.H.) Muriwai, 11/3/51, 9 seen at stream mouth.—(J.C.D.) Bethells, 15/10/50, a pair with nest and 1 egg.—(N.M.) Puketutu, 1/4/51, 16 (N.M.); 1500 plus feeding on mud at low tide (A.H.H.) Whitford, 29/7/51, 200 plus in field half-mile from coast.—(M.A.d'A. and C.D.B.) Kaiaua, Miranda coast, 1950-51, breeding in usual numbers, 20 plus pairs; Piako, 29/4/51, 1000 plus in flock.—(H.R.McK. et al.) Miranda, 1 nest, 2 eggs on 3/12/50; Kawhia, 21/2/51, 150 plus.—(N.M.) Tauranga Harbour, near station, Jan., 1951, one only.—(A.S.W.) Chateau Golf Links, 21/10/50, three.—(T.S.) When camped at Waitotara Estuary for six days from 10/1/51, there were only about 12 banded dotterels on the estuary and lagoon during the first three days. Then a flock of about 40 arrived and were still there when we left, scattered feeding on the estuary and lagoon area.—(W.P.M.) Waiwakaiho River mouth, Bellblock Beach, on shingle flats.—(O.E.G.) Porangahau Estuary, 30/9/51.—(R.L.G.) Ohau Estuary, 4/3/51, c. 100.—(A.S.W.) Waikanae River Estuary, 2/2/51, only one.—(B.I.) Lake Wairarapa north, 18/2/51, 12; Lake Onoke north, 25/2/51, 1; Palliser Spit, 13/5/51, none; 15/7/51, 1; 1/9/51, c. 3; 23/9/51, c. 5, not yet breeding.—(J.M.C.) Outlet of Lake Forsyth, 2 pair, one with nest and three eggs, 20/11/51; New Brighton Beach, 8/4/51, six.—(W.C.C.) South Brighton Beach, 17/5/51, 107 scattered on beach; Waimari Beach, 25/4/51, 30 in sandhills.—(D.E.C.) Ashley Mouth, 22/8/51, two (E.W.D., I.D.R.C.); 16/9/51, three pairs, one chick ringed No. 4706 (E.W.D., F.R.N.) Rangitata River mouth, 18/1/49, seven; Opihi River, near Hanging Rock, 20/1/49, two pairs.—(D.N.F.C.) Kyebrun, Central Otago, 20/8/50, six.—(M.A.E.B.) Allan's Beach, Otago Peninsula, flock of c. 70 during May and June, 1951.—(B.J.M.)

GEOFFREY'S SAND PLOVER (*C. leschenaulti*)—Miranda, 4/11/50, two, one with grey shoulder tabs and one with fading reddish tabs, dark eye patches at a minimum; 28/12/50, one; 14/1/51, one showing a little new colour; 10/2/51, one; 26/8/51 and 8/9/51, one.—(H.R.McK. and R.B.S. et al.)

NEW ZEALAND DOTTEREL (*Pluviorhynchus obscurus*)—Ruakaka, 13/2/51, one.—(N.M.) Great Barrier Is., 21/11/50, at least 12 were seen on Kaitoki Beach.—(J.C.D. and B.D.H.) Whatipu, 21/1/51, two seen together in the sand dunes.—(J.C.D. and G.W.G.) Miranda, 7/11/50, 3 adults and one running chick (R.H.D.S.); 18/3/51, 1 adult and 2 juv., 1/4/51, 2; 24/6/51, 2, pale. (H.R.McK. et al.)

WRYBILL (*Anarhynchus frontalis*)—Ruakaka, 13/2/51, two; Puketutu, 11/1/51, c. 100; 4/2/51, 222; 18/2/51, 300; 12/4/51, 310; 14/4/51, 346; c. 200 on 20/8/51; c. 120 on 31/8/51.—(J.C.D., N.M. and R.B.S.) Ruakaka, 12 summered 1950-51, maximum count 250 on 31/3/51.—(D.A.U.) Shelly Park, Howick, 22/7/51, 24 on far side of river, first record here.—(N.M.) Waikokopu Creek, Whitford, 29/7/51, 150 plus feeding on tide flat, first record here; 11/8/51, c. 200.—(M.A.d'A. and C.D.B.) Miranda, 4/11/50, two; 30/12/50, three (N.M.); 1951 winter tally, c. 1430 (J.C.D., H.R.McK. et al.) Kawhia, 21/2/51, c. six feeding at low tide; 23/2/51, three feeding by themselves.—(N.M.) Waitotara Estuary, six on 10/1/51 and 12/1/51.—(W.P.M.) Palliser Spit, 23/9/51, one, full band.—(R.H.D.S.) Washdyke Lagoon, Timaru, 16/10/50, several.—(J.M.C.)

SPUR-WINGED PLOVER (*Lobibyx novaehollandiae*)—Wendon, Gore, female, 15/5/51.—(M.J.M.)

KNOT (*Calidris canutus*)—Miranda coast, 1950-51, summer, several thousands; 24/6/51, Kaiaua to Thames, 630 plus.—(J.C.D. and H.R.McK. et al.) Maketu, 14/12/50, two amongst a flock of godwit, not often seen on the east coast.—(A.H.W.) Kawhia, 25/2/51, several feeding with godwit.—(N.M.)

AMERICAN PECTORAL SANDPIPER (*C. melanotos*)—Miranda, one on several dates from 1/10/50 to 10/2/51.—(H.R.McK. et al.) Porangahau Estuary, 16/11/50, two associated with c. 50 godwits.—(K.W.)

SIBERIAN PECTORAL SANDPIPER (*C. acuminata*)—Miranda, 1/10/51, four; 4/11/50, 14; 3/12/50, 10; 14/1/51, nine; 10/2/51, 18.—(H.R.McK. et al.) Ahuriri Lagoon, H.B., 7/4/51, nine.—(D.B.)

RED-NECKED STINT (*Pisobia ruficollis*)—Miranda, one seen on 4/11/50, 23/12/50 and 10/2/51.—(H.R.McK. et al.)

CURLEW SANDPIPER (*Erolia testacea*)—Miranda, one seen several times from 4/11/50 to 10/2/51 (H.R.McK. et al.); 8/11/50, two feeding with sandpipers and a red-necked stint.—(R.H.D.S.)

BAR-TAILED GODWIT (*Limosa lapponica*)—Mangonui, 9/2/51, c. 25. on small island off coast; Ruakaka, 13/2/51, c. 20.—(N.M.) Great Barrier, 21/11/50, three very pale birds feeding on Kaitoki Beach.—(J.C.D.) Whatipu, 20/1/51, four.—(G.W.G. and J.C.D.) Puketutu, Manukau Harb., 1/4/51, c. 2000; 14/4/51, c. 30.—(N.M.) Howick, 27/12/50, c. 100 on far side of creek; 11/3/51, c. 600 plus flew over toward the Manukau at about 2 p.m.; then 500 plus in same direction about 4 p.m. (low tide), some knot among them; 15/7/51, c. 300 resting above high water mark on far side, no attempt to fly away as they usually do.—(N.M.) Waikokopu Creek, Whitford, 29/7/51, four.—(M.A.d'A. and C.D.B.) Mataitai, Clevedon, 1950-51 summer population c. 1500 (H.R.McK.), 9/10/50, many while performing aerobatics hit the water with a splash (F.M.) Miranda coast, summer population several thousands; 24/6/51, from Kaiaua to Thames, winter population c. 1290, no evidence of shooting this year.—(J.C.D., H.R.McK. et al.) Sulphur

Point, Tauranga, 9/5/51, c 26 all in winter plumage, may be total winter population for the harbour.—(A.H.W.) Ohiwa Harbour, 12/8/51, 500 plus on sand-bar on east side of harbour, probably the full winter population.—(P.H.B.) Kawhia, 21/2/51, 14; Rakanui, 25/2/51, 200 plus.—(N.M.) Tauranga Harbour, Jan., 1951, c. 500 near station; Maketu, 25.—(A.S.W.) Porangahau Estuary, 30/9/51, c 27.—(R.L.G.) Ohau Estuary, 12/6/51, six.—(A.S.W.) Okarito, 7/5/51, two.—(G.G.) Christchurch Estuary, 8/4/51, nine and 50 plus, winter plumage.—(E.W.D.) Ashley Mouth, 16/9/51, twelve, winter plumage.—(E.W.D., F.R.N.) Brooklands Beach, 18/5/51, seven feeding among pied oystercatchers.—(D.E.C.) Heathcote-Avon Estuary: 6/2/51, about 3000 godwits circled widely over south end of sandspit on seaward side of estuary; they were in three flocks, one of about 2000 and two of about 500 each. 23/3/51, about 1000 godwits still around; 26/3/51, no godwits remaining in estuary. In 1950 some birds (as recorded previously) returned during winter. During 1951 none were seen by me in estuary between 22/3/51 and 1/10/51. On latter date, about 500 godwits returned to estuary. 5/10/51, flock of godwits appears to have increased in number. As tide waters rise in the estuary the area of dry sand rapidly decreases in size. Black-backed gulls have first priority near coast, pied oystercatchers second and godwits third; godwits are first to move as each area becomes crowded.—(G.G.) Heathcote Estuary, 19/8/50, 27 plus; 16/12/50, large flocks (L.E.W.); 26/2/51, 30 plus (W.C.C.); 26/1/51, hundreds (I.T., L.E.W.). Heathcote-Avon Estuary, 19/2/50, large numbers, probably 2,000; 21/2/50, disappeared; 15/8/50, c. 100 in winter plumage.—(G.G.) Washdyke Lagoon, Timaru, 16/10/50, 19.—(J.M.C.) Allan's Beach, Otago Peninsula, May and June, largest count 50, 17/6/51.—(R.R.M.) Blue-skin Bay, 30/1/51, 20 plus.—(I.T.)

WHIMBREL (*Numenius phaeopus variegatus*).—Clevedon, one seen often from 28/10/50 to 18/3/51. On 5/2/51, a second whimbrel was harrying the other, its rump feathers up and tail down towards ground at severe angle. Miranda, 28/12/50, one; 10/2/51, seven in a party, five identified as *N. p. variegatus*, sizes varied considerably; 1/4/51, call heard at Piako by F.M., L.H.M. and J.G.R.; 26/5/51, one; 24/6/51, two.—(R.B.S., H.R.McK., et al.)

PIED STILT (*Himantopus himantopus*).—Tara, near Mangawhai, four pairs nested in swampy area, first egg 26/8/51; have nested for four years in same place; birds in district increasing.—(L.W.) Mangonui, 9/2/51, c. 30 on small island off shore; Paihia, 12/2/51, c. 12 in bay next to Paihia.—(N.M.) Te Arai South, Rodney County, 22/10/50, 3 nests in wet field, 4, 3 and 3 eggs, one clutch chipping; Great Barrier, 21/11/50, three seen on Kaitoki Beach.—(J.C.D. and B.D.H.) Mangawhai, 1/9/51, four nests with eggs.—(S.J.M.) Waikeke Is., 1/1/51, 25 plus including several young at Awaawaroa Bay.—(A.J.G.) Rotorua Is., E. of Waiheke Is., 5/11/50, c six reported on a dam (H.R.McK.) Waiuku, 11/8/51, four on pools formed by rain in paddock; 11/8/51, at the Waitangi Falls, seven (N.M.); 6/5/51, in the channel, one seen doing aerobatics hit and splashed the water several times.—(H.R.McK.) Shelly Park, Howick, 25/11/50, none; 27/1/51, two; 4/3/51, 14; 26/3/51, 32; 14/7/51, two; 1/9/51, none.—(N.M.) Miranda, 1950-51, breeding was fairly successful; 28/12/50, a late nest, 4 eggs; 14/1/51, a small chick of this nest seen, it was observed until flying. Kaiaua-Thames, 29/4/51, c. 4,450; 24/6/51, c. 6930; 26/8/51, nest four eggs at Kaiaua.—(H.R.McK. and R.B.S.) L. Tarawera, 26/2/50, six observed on a shallow lagoon about 400 yards below the outlet, four adults and two adolescents.—(M.J.S.B.) Kawhia, 19/2/51, 150 plus.—(N.M.) Tauranga Harbour, near Station, Jan., 1951, counted 139; Maketu, 70.—(A.S.W.) Porangahau Estuary, 30/9/51, c. 4.—(R.L.G.) Kopane, Manawatu, 11/7/51, two, first of season; 16/7/51, four; 23/7/51, eleven.—(E.D.) Ohau Estuary, 4/3/51 c. 12.—(A.S.W.) Waikanae River Estuary, 2/2/51, eight.—(B.I.) Lake Wairarapa, north, 18/2/51, only four seen; Lake Onoke, north, 25/2/51, only two.—(J.M.C.) Flat Point, 16/5/51, seven; 29/8/51, six.—(J.H.S.)

Karori, 1/9/49, perhaps two birds passing south-west at 2000; from the way calls were made it appeared that the birds on meeting the crest of a hill veered east and west along its ridges until the correct direction was found.—(H.L.S.) Ashley River, 25/11/50, common; two nests each with four eggs, 18/12/50; Christchurch Estuary, 12/8/51, numerous; Woodend, numerous, 19/5/51, three 16/9/51.—(E.W.D.) Heathcote-Avon Estuary, west end, 3/2/50, c. 250; 15/8/50, c. 20.—(G.G.) Rangitata River mouth, 18/1/49, 16; Opihi River, near Hanging Rock bridge, 20/1/49, 22.—(D.N.F.C.) Kyeburn, C. Otago, 26/8/50, 17 in one flock; swamp near Patearoa, c. 40.—(M.A.E.B.) Merton Swamps, 17, Kartiki Beach, six or seven, 21/10/50.—(D.N.F.C.) Stirling, 10/2/49, for three weeks have been varying number up to 15 feeding on pond; 12/4/49, very noisy at twilight. Watched two adults join three young; all marched along in line yelping like fife-and-drum band; suddenly stopped, turned right, put heads under wings; gone next day; a flood ensued; no more stilts seen till spring; 11/9/49, one at pond; 13/9/49, pair seen; 18/9/49, two pairs; 22/2/51, three, came occasionally till 25/4/51 and 28/4/51, heard at night.—(R.V.McL.) Allan's Beach, Otago Peninsula, largest count 50, 3/6/51.—(R.R.M.) Lake Waihola, 29/8/50, eight; Akatore, 19/9/50, seven; Taieri Mouth five, and Akatore seven, 21/11/50; Shag River lagoons four, and Blueskin Bay 20 plus, 30/1/51.—(I.T.) Lake Tuakitoto, c. 40; Lake Waihola three and Port Molyneux two, 30/9/50.—(I.T., L.E.W.) Near Akatore, Otago, 31/12/50, four; Waitati Bay, Otago, 10/1/51, 50; Taieri mouth, Otago, 24/3/51, 35; foreshore, Otago Harbour, 30/3/51, 32, 10/5/51, 102.—(L.E.W.) Kew, Invercargill, 5/2/51, two; Goodwood, 6/2/51, two; South Timaru, 6/2/51, numerous; 19/1/51, 28 plus; Warrington, 19/1/51, 50 plus.—(E.W.D.)

BLACK STILT (*H. himantopus*).—Miranda, one with some whitish patches, seen often from 18/3/51 to 24/6/51.—(H.R.McK. et al.) Kawhia, 19/2/51, one.—(N.M.)

BLACK-FRONTED TERN (*Chlidonias albigularis*).—Palliser Spit, 13/5/51, c. 21; 15/7/51, c. 18; 1/9/51, none; 23/9/51, none.—(J.M.C.) Ohariu Bay, Wellington, 7/1/50, decapitated corpse, wing 225mm.—(H.L.S.) Ashley River, 18/12/50, nest with two eggs.—(E.W.D.) Ashley mouth, 19/5/51, one (E.W.D., J.C.Y.); 22/8/51, 30 plus (E.W.D., I.D.R.C.); 16/9/51, six (E.W.D., F.R.N.) On way to Mt. Peel, 19/1/49, three or four solitary birds over fields; Opihi River, near Hanging Rock, 20/1/49, c. 20 adults and young on shingle island, two adults dived at intruders; between Geraldine and Burke's Pass, 22/6/49, few over rivers; between Geraldine and Mt. Peel, 24/1/49, ten over fields.—(D.N.F.C.)

CASPIAN TERN (*Hydroprogne caspia*).—Mangawhai, breeding ground in the sandhills has been in occupation for at least 50 years. In October, 1950, estimate of 1500 nests in three groups, unfortunately a great deal of damage is done each year by sightseers, etc.—(L.W.) Howick, up to four seen regularly; 11/3/51, two quarrelling, screeching and flying at one another in mid-air, looked like immature birds.—(N.M.) Puketutu, Manukau Harbour, 1/4/51, 64.—(N.M.) Clevedon River mouth, 14/1/51, 70 best tally for season.—(A.J.G.) L. Tarawera, 26/2/50, four observed during the day, fishing well off shore.—(M.J.S.B.) Tauranga Harbour, near Station, Jan., 1951, 39.—(A.S.W.)

FAIRY TERN (*Sterna nereis*).—Mangawhai sand hills, six in flight 10/1/51; Mangawhai Estuary, one pair 17/7/51 and one pair 28/8/51; these birds were once fairly plentiful but are still not uncommon; they nest on the Mangawhai sand hills.—(L.W.)

WHITE-FRONTED TERN (*S. striata*).—Mohaka River Estuary, 25/11/50, 200-300 displaying and copulating on spit but only two nests with one egg each.—(K.W.) Ngawe Point, Palliser, apparently did not nest 1950-51; Mataikona, 1/1/51, small colony (eggs and chicks) reported remains seen later; Castlepoint, 16/12/50, c. 20.—(J.M.C.) Castlepoint, 15/5/51, c. 60 including c. 20 immature.—(J.H.S.) Flat Point, 16/5/51,

c. 40; Pahau, 17/5/51, remains old nests seen.—(J.H.S.) Palliser Spit, 13/5/51, c. 190, only one immature; 15/7/51, none; 1/9/51, two; 23/9/51, none.—(J.M.C.) Birdling's Flat, Canterbury, 2/2/50, c. 1000 (many being that season's chicks) were flocked on the shingle beach.—(G.G.) Rangitara River mouth, 18/1/50, c. 50 young flying.—(D.N.F.C.) Diamond Harbour, 30/11/50, approx. nine nesting pairs with eggs and young.—(W.C.C.) Six at sea between Lyttelton and South Brighton, 10/3/51; six, Christchurch Estuary, 8/4/51; ten, Ashley Mouth, 19/5/51; 40 plus, Ashley Mouth, 22/8/51; c. 40 Ashley Mouth, 16/9/51; common, Menzies Bay, 5-11/5/51.—(E.W.D.) Okarito, 7/5/51, numerous.—(G.G.) Spit, Otago Heads, 8/4/51, thousands on mole.—(L.E.W.) Two at The Nuggets chased by skuas, 30/9/50; hundreds, Christchurch Estuary, 26/1/51; 241, Kartiki Beach, 27/3/51.—(I.T., L.E.W.)

CRESTED TERN (*Thalassius bergii*).—Spirits Bay, Northland, March, 1951, one dead, dry, at high tide mark.—(H.R.McK.)

BLACK-BACKED GULL (*Larus dominicanus*).—Waikanae River Estuary, 2/2/51, c. 950.—(B.I.) Palliser Spit, first chicks hatched c. 1/12/50, practically all hatched by 1/1/51.—(J.M.C.) Diamond Harb., 30/11/50, two nests with three eggs each on rock ledges only few feet from sea.—(W.C.C.) Heathcote-Avon Estuary, 23/9/50, flying excitedly about east end of estuary, probably courting.—(G.G.) Lake Tuakitoto, 50; appeared to be nesting on a rocky islet, Nuggets, 30/9/50; several thousands, mouth of Kaikorai Stream, 17/5/51.—(I.T., L.E.W.) Over 100 nesting on island, Moeraki Kaik, 21/10/50; one seen, Margaret Glacier, Dart Valley, at height of 6000ft., 31/1/51.—(B.W.C.)

RED-BILLED GULL (*L. novaehollandiae*).—Whangaroa, Jan.-March, 1951, thousands, mostly young of the season, died all along the coast, apparently starved as very light; heavily infested with vermin; school fish were lacking all the season.—(T.M.R.) Howick, 24/4/51, one chasing a sparrow with a crust of bread and darting again and again at a wire-netting fence when the sparrow flew through and it was unable to follow.—(N.M.) Porangahau Estuary, 18/11/50, southern spit, about 50 birds in attendance on 21 nests, five of which contained a single egg.—(K.W.) Waikanae River Estuary, 2/2/51, c. 1150.—(B.I.) Ngawe Point, Palliser, 23/12/50, c. 100 chicks c. 10 days old; Mataikona, 1/1/51, small colony (eggs and chicks) reported, remains seen later.—(J.M.C.) Saltwater Creek Estuary, 26/3/51, c. 100.—(D.E.C.) Christchurch Estuary 29/7/51, 40 plus.—(W.C.C.)

BLACK-BILLED GULL (*L. bulleri*).—Matiatia, Waiheke Is., 27/3/51, one alone seen clearly, the black bill and plain upper wing noted.—(A.J.G.) Clevedon, 1/7/51, five; 14/7/51, four.—(H.R.McK.). Miranda, 1950-51 summer, a few odd ones; 1951 winter, up to 650.—(H.R.McK., R.B.S., et al.); Lake Tarawera, 26/2/50, three small parties, 9, 5 and 4, seen on lake well out from the shore.—(M.J.S.B.) Motuopa Bay, Lake Taupo, 29/10/50, two.—(R.H.D.S.) Porangahau Estuary, 18/11/50, southern spit, about 100 birds on a separate hummock, about 20 yards away from red-billed gull colony. There were 37 nests, 11 with one and one with two eggs.—(K.W.) Ada River, near Ada Station Homestead, 2/1/51, c. 150.—(W.C.C.) Lake Pearson, 14/1/51, two.—(E.W.D.) Ashley Mouth, 19/5/51, c. 40 mostly juv., 3 with rings.—(E.W.D., J.C.Y.); 22/8/51, c. 40, with one red-billed gull.—(E.W.D., I.D.R.C.) Rangitara River Mouth, 18/1/49, hundreds; Opihi River, near Hanging Rock, 20/1/49, small flocks; occasional small flocks followed tractor on farms.—(D.N.F.C.)

SOUTHERN SKUA (*Catharacta skua lonnbergi*).—Single bird eight miles north of Moturaratiti Island, Cook Strait, 1/9/51.—(C.A.F.)

ARCTIC SKUA (*Stercorarius parasiticus*).—29/4/51, four seen on circumnavigation of Waiheke.—(K.C.B.C.) Kaiaua, 28/12/50, 10 plus flying past point in E.N.E. gale; no doubt more further out, but could

not be identified in rain and spray; all flying north. Muriwai Beach, 29/1/51, one chasing terns outside breakers.—(H.R.McK.) Near Coromandel, 22/5/50, two circled boat.—(J.S.E.)

POMARINE SKUA (*S. pomarinus*)—Kaiaua, 28/12/50, one flew past point in E.N.E. gale; seen fairly clearly and compared with Arctic species, but identity not certain.—(H.R.McK.)

NATIVE PIGEON (*Hemiphaga novaeseelandiae*).—Great Barrier, 13-22/11/50, c. 10 seen during our stay.—(B.D.H. and J.C.D.) Cascades, Waitakere, 7/7/51, one; Karekare, 14/7/51, one.—(N.M.) Moumoukai, Clevedon, 1951, a pleasing increase, up to 14 seen in flock.—(J.W.St.P.) Minginui area, numbers seen: Nov., 1950, hardly any; Dec.-Jan., none to six; Feb., none to nine; March, April, up to 30; May, up to 11; June, up to five, but seen only four days in thirty. Feeding noted: Late Jan., Feb., tawa fruit; March, tawa, rimu and kahikatea; April, the same, but miro in high country; May, mainly kahikatea; June, gone to supplement jack country.—(R.St.P.) Fletcher's Bay, Coromandel, May, 1950, one.—(J.S.E.) Kairimu Road, Marakopa, only odd ones flying over for years but this year (1951) two wintered quite close to house, still there 20/9/51. (R.E.W.F.) Western Hutt Road, Belmont, 5/8/51, one in flight.—(R.H.D.S.) Okute Valley, 10/12/51, two.—(D.E.C.) Christchurch Bot. Gardens, 3/4/51, 23/6/51, one.—(E.W.D.) Geraldine Hill bush, 16/1/49, three; Te Moana Gorge, 17/1/49, two; cottage owner latter place reported 40 feeding on berries of 20ft. high laurel hedge when berries are ripe; Lynn Creek, Mt. Peel, 19/1/49, three; Mt. Peel Forest, 24/1/49, count for day, six.—(D.N.F.C.) On wineberry tree outside bedroom window, Bragg's Bay, Stewart Island, 16, 5/2/50; on wineberry, Milford Sound, four, 9/2/51.—(Mrs. A. Prentice, D.N.F.C.) Woodside, 23/9/50, c 20 plus.—(D.N.F.C.) Hidden Falls Valley, W. Otago, 6/2/51, one.—(B.W.C.) On edge of bush, Maungatua, at 2000ft., 2/6/51, one.—(R. Gilbreath, D.N.F.C.) During the past year numbers up to seven have been recorded in gardens in Maori Hill, Dunedin.—(I.T.) Waipori Gorge, 1/4/51, c 36.—(I.T., L.E.W.) Swampy Hill, Dunedin, 6/5/51, about a dozen in bush, largest number ever seen in this area which is frequently visited.—(E.M.M.) Barrier Hut, Pyke Valley, 31/12/49, four.—(W.A.W.)

ROCK PIGEON (*Columba livia*).—Nesting under several wharves, Wellington Harbour, 20/10/50.—(R.V.R.) Rangitata, Canterbury, 14/12/50, large flock in paddock.—(L.E.W.)

LACE-NECK DOVE (*Streptopelia chinensis*).—Reported in Mt. Albert crater, 8/4/50, on ornamental cherry trees and high upon the bare rocks. The beautiful fan tail showing the white bar was clearly seen several times during the morning.—(R.V.R.)

KAKA (*Nestor meridionalis*).—On 12/11/50, R.H.McK. and I found a nesting hole in a puriri near the Te Waikohare Track, Little Barrier Island. I was passing here most days during the next fortnight and the bird would come out if its mate flew near calling, or if there was any disturbance against the trunk of the tree. On two occasions the sitting bird was seen to be fed by its mate high up in a nearby tree. Owing to a bend in the hole it was not possible to see down to the nest, but from the time taken by the bird to come out, it is presumed that the nest was some distance below the entrance. There was no sign of food being carried to chicks up to the time we left.—(W.P.M.) Great Barrier 13-22/11/50, very common in all the bush-clad areas, parties of seven and eight were seen aerobating in the deep valleys on the upper slopes of Mt. Hobson (2038ft.).—(J.C.D. and B.D.H.) Moehau, Oct., 1949, flock of c. 30; Jan., 1950 and Jan., 1951, odd ones; 27/8/51, two.—(G.J.B. and R.N.B.) Whenuahiti, Coromandel Peninsula, June, 1951, two seen and heard, report by Mr. M. S. Thorn.—(H.R.McK.) North Road, Clevedon, 13/5/51, one in patch of bush on Duder's Point, seen and heard.—(B.F.D.) Moumoukai, Clevedon, 1950-51, very scarce, position disquieting.—(J.W.St.P.) Minginui, Nov., 1950-June, 1951, usually two seen

about the bush workings; others heard. More plentiful in the higher beech country and some in the rata-tawa belt between the high country and the timber valley. 4/2/50, a nest found in a hollow half of a matai with forked trunk, 30 feet from the ground, entrance five feet above the nest, a 2in. crack in trunk from top down to nest. The bushmen felling the tree did not know of the nest and the kaka flew out when it hit the ground. The two eggs were broken and were not found until the log had been hauled 20 chains to the skids. The two kaka stayed around and called for two days, after which they remained in the locality. 8/2/51, a nest 25 feet up in a matai, containing two half-grown young. The bushmen saw the parents flying about close to them but did not suspect a nest when felling the tree on the afternoon of the 7th. On the morning of the 8th when the log was rolled over a faint screech was heard. The tractor driver chopped away some wood and got out the surviving chick, having his fingers well bitten while doing so. It was very weak, so was taken to camp and was fed on tawa berries and konini which the bushmen brought for it each evening. At the end of a week it would eat almost any kind of food and seemed to need quite a quantity of water to drink.—(R.St.P.) Near Mt. Moehau, Coromandel, May, 1950, heard several times.—(J.S.E.) Whakatane watershed, 8/3/51, one heard.—(T.S.) Heard in bush near Alabaster Pass, N.W. Otago, 30/12/49; three flying above Pyke River, near Upper Pyke Hut, 31/12/49.—(W.A.W.) Three, Beansburn Valley, W. Otago; four in forest, Hidden Falls Valley; others heard, 3-6/2/51.—(B.W.C.) Titiroa Range, Dec., 1950, heard several times.—(B.T. and J.S.E.)

KEA (*N. notabilis*).—Edwards River, Arthur's Pass, 6-8/1/51, several heard, one seen.—(E.W.D.) Temple Basin, 7/7/51, seven; Ada Valley, 30/12/50, one.—(W.C.C.) Two, Cow Saddle, one heard north branch Routeburn, one slopes Mt. Hedon, Dart Valley, 31/1/51.—(B.W.C.) Two near North Col, north branch of Routeburn Valley, 28/12/49; one seen and others heard near Cow Saddle, N.W. Otago, 29/12/49; two near Hidden Falls Hut, Hollyford Valley, 6/1/50.—(W.A.W.)

RED-FRONTED PARAKEET (*Cyanoramphus novaeseelandiae*).—Whakapapa Valley, near Chateau, 25/6/50, two seen.—(T.S.)

YELLOW-FRONTED PARAKEET (*C. auriceps*).—On 12/11/50, when with R.H.MeK. on Te Waikohare Track, Little Barrier Is., he saw a parakeet come out from a nesting hole in a tree alongside the track. We presumed this would be the usual red-fronted parakeet and passing there almost daily during the next fortnight I did not see the bird actually coming from the hole again. But on one occasion a yellow-fronted parakeet was perched on a branch level with the nesting hole and only a few feet from it, looking towards the hole. Actually I did not recognise it as a yellow-fronted till it turned to look down at me, which illustrates the difficulty in differentiating between the two species when seen under the usual bush conditions.—(W.P.M.) Minginui, Nov., 1950-Jan, 1951, seen almost daily in very small numbers in lower areas, but more plentiful in high beech country where up to 10 seen in one day.—(R.St.P.)

PARAKEET (*Cyanoramphus* ? sp.).—South of Pokako, 14/11/50, heard.—(R.H.D.S.) Heard near Barrier Hut, Pyke Valley, N.W. Otago, 31/12/49.—(W.A.W.)

KAKAPO (*Strigops habroptilus*).—In early February, 1951, Mr Jack Ede, at the head of the Tutoko Valley, Milford Sound district, heard the booming call of a kakapo on four successive nights, but did not see the birds. He also found what appeared to be a nest of the kakapo "scratched in the ground at the base of a tree." It was about 11in. in diameter and 6in. deep, with four tracks leading away from it in different directions; about 2in. wide and quite clear near the nest, but they faded away as they travelled further away. There were no feathers in the nest." The cry is described as "a booming call which

came from the vicinity of the nest—answered in a similar manner by another bird further down the valley. The calls averaged from fourteen to sixteen distinct booms starting quietly and reaching a crescendo at the seventh boom." George Guy tells me that Mrs. Guy some years ago "saw a kakapo when travelling by night from Lake Manapouri to Doubtful Sound. The bird appeared in the pathway, in dense bush, and when a torch was shone on it, the bird slowly walked off."—(R.J.S.)

EASTERN ROSELLA (*Platycercus eximius*).—Bethell's, November, 1950, eight in flock.—(R.V.R.) Karekare, 14/7/51, c. 20; Huia, 18/8/51, two.—(N.M.) Orere, Clevedon, 30/8/50, two reported seen.—(H.R.McK.) During the past year this bird has neither been heard nor seen in the garden nor on walks in the Reservoir Reserve, Dunedin; several reports received that the birds have been netted for cage birds.—(I.T.) July, 1951, has returned to Maori Hill Reservoir Reserve, from which area it had disappeared after caging operations over a year ago.—(E.M.M.)

SHINING CUCKOO (*Chalcites lucidus*).—Mangawhai, first heard this season, 2/10/51.—(L.W.) Great Barrier, 13-22/11/50, well dispersed throughout the island.—(J.C.D. and B.D.H.) Muriwai, visit only on 27, 28, 29/1/51, a full song each day, downward notes added.—(H.R.McK.) Remuera, Auckland, 23/11/50, at 11 p.m. heard to call continuously, twice 95 upward notes and once about c. 45.—(J.C.D.) Clevedon, 3/1/51, last song; 23/2/51, one seen.—(H.R.McK.) North Road, Clevedon, 7, 10/3/51, latest calls heard.—(A.J.G.) Moumoukai, Clevedon, 18/9/50, first song.—(J.W. St.P.) Minginui, Nov., 1950, to June, 1951; Nov., up to eight seen per day, singing in parties; Dec., up to ten seen; Jan., less seen, but many heard; Feb., only odd ones seen, song decreasing; March, none.—(R.St.P.) Waikanae, 30/9/51, first call of season.—(W.K.) First heard in Wadestown, 22/9/50.—(R.V.R.) Brothers Is, Cook Strait, 24/9/51, one found near tower in bedraggled state (thick drizzle), released next morning, flew away strongly northwards.—(J.S.) Nelson City, 20/9/51, heard first song, 10.30 hours, bright sunny day, also reliable report of having heard song in Nelson City on 18/9/51.—(L.G.) Anderson's Bay, Dunedin, 25/9/50, heard.—(M.A.E.B.) Kaka Point, 20/10/50, first calls.—(M.P.F.) Lower Taieri Gorge, 3/3/51, one seen.—(J.G.) One heard calling near river flats in the lower part of the Hidden Falls River, 29/12/49; heard calling near Barrier Hut, Pyke Valley, 31/12/49.—(W.A.W.) Report from Sullivan Dam, Leith Valley, heard 8.30 a.m. and again 3 p.m. 25/9/50; our garden, Maori Hill, 27/9/50; heard garden 13 days in October; daily in November and occasionally seen; recorded 16 days in December; heard daily January 1 to 7 and not heard again; one seen in garden, 14/3/51.—(I.T.) Glenleith, Leith Valley, 26/9/50, calls 4.5 p.m.—(W.H.D.) Opoho, 14/10/50, heard.—(Miss I. McArthur, D.N.F.C.) Howick, Auckland, 6/10/50, heard.—(Miss I. T. Currie, D.N.F.C.) Leith Valley, five calls and about 15 second later the end notes, 2/1/51.—(W.H.D.) Stirling, 30/1/51, one seen during strong northerly on hawthorn tree in shelter of house; seemed to be devouring the green berries; only one seen this summer; none heard.—(R.V.McL.)

LONG-TAILED CUCKOO (*Eudynamis taitensis*).—Howick, 15/4/51, one in macrocarpa tree in garden.—(N.M.) Minginui, 26/10/50, first call; Nov., 1 or 2 seen each day; Dec., up to four; Jan., down to one some days; Feb., very few; March, none.—(R.St.P.) Whakatane watershed, Urewera, 8/3/51, one heard.—(T.S.) Twice seen and commonly heard, Arthurs Pass, 9-13/1/51.—(E.W.D.) Westland, Feb., 1951, common from Lewis Pass to Fox Glacier.—(G.G.) Morrison's Creek, Leith Valley, 18/11/50, one heard and seen.—(J.G.) One calling on west side of Alabaster Pass, N.W. Otago, 30/12/49.—(W.A.W.) Titiroa Ranges, Dec., 1950, heard occasionally.—(C.B.T. and J.S.E.)

MOREPORK (*Ninox novaeseelandiae*).—Little Barrier Is., 11/11/50, bird sitting on nest in hollow puriri near house; 21/11/50, one egg hatched; part of a rat seen in nest; C.H.P. reported second egg had

hatched later.—(C.H.P., W.P.M., J.W.St.J. and H.R.McK.) Mt. Moehau, May, 1950, several heard.—(J.S.E.) Hunua, 9/4/51, none recorded last two years.—(D.N.L.) Kopane, Palmerston North, in this district the morepork feeds mainly on the introduced house sparrow. I have seen the owls actually taking sparrows in the grey dawn.—(E.D.) First record this year of morepork in Wadestown, 17/3/51, as it was mobbed by small birds.—(R.V.R.) Lake Brunner, 3/6/51, heard.—(W.C.C.) Recorded Maori Hill on following dates: 10 and 16/6/50, 3/7/50, 24/7/50, 6 and 26/8/50, 12/9/50, 17, 18, 19/10/50, 1/6/51.—(I.T.) Calling, Lake Te Anau, near Glow Worm Caves, 26/3/51, and Milford Sound 27/3/51.—(K.V.F.) Two calling at junction of Pyke and Olivine rivers, N.W. Otago, 30/12/49; heard at Upper Pyke Hut, Pyke Valley, 31/12/49, 1/1/50.—(W.A.W.) Titiroa Ranges and district, Dec., 1950, heard occasionally.—(C.T.B. and J.S.E.)

LITTLE OWL (*Athene noctua*).—Commonly heard at night in parks and gardens in central Christchurch.—(E.W.D.) Hilderthorpe, N. Otago, 14/3/50, harrier seen eating remains of one; 5/4/51, one heard, seen 10.45 a.m., not so plentiful.—(C.W.McL.) Recorded garden, Maori Hill on following dates: 26/7/50, 27/8/50, 21/4/51, 3, 7/5/51.—(I.T.) About 5 p.m. on tree in Museum Reserve, Dunedin, 25/6/51.—(K.P.S.)

KINGFISHER (*Halcyon sanctus*).—Great Barrier, 13-22/11/50, very common all over the island, even in the deep bush and at the top of Mt. Hobson (2038ft.).—(J.C.D. and B.D.H.) Minginui, Nov., 1950–June, 1951, Jan., up to nine seen each day, then declining to odd occurrences; Jan., feeding beetles to young; they come here to breed but must winter in lower warmer country.—(E.St.P.) Nest on Flat Is., Hauraki Gulf, 6/1/51, 150 feet above sea level on steep cliff face; three young, within a few feet of the nest of a blue penguin and a petrel. Nest containing three young and number of dead lizards six feet above high water in Katherine Bay, Great Barrier Is.; kingfishers abound in Katherine Bay.—(R.V.R.) Fletcher's Bay, Coromandel, May, 1950, several seen with skinks.—(J.S.E.) Christchurch Botanical Gardens, 23/6/51, three (E.W.D.) At Owaka, one chased by six sparrows, Feb., 1951.—(Mrs. J. A. Moore, D.N.F.C.) Between Portobello and Macandrew's Bay, 9/6/51, six.—(Miss I. McArthur, D.N.F.C.) Stirling, sometimes seen; 14/4/51, seen in orchard fighting with blackbird, driven off, not seen again.—(R.V.McL.) Spit, Otago Heads, 8/4/51, two birds on wires, one a very bright yellow instead of buff.—(L.E.W.)

RIFLEMAN (*Acanthisitta chloris*).—Minginui, 14/12/50, four, the only ones seen in lower country; up to 24 seen on trips to high country.—(R.St.P.) Nelson Reservoir to Dun Mt., 15/4/51, 15 seen, 30 plus others heard.—(R.B.St.P.) Lake Guyon, 3/1/51, very common in beech forest.—(W.C.C.) Seen once and often heard, Edwards River, Arthur's Pass, 6-8/1/51.—(E.W.D.) Geraldine Hill bush, 16/1/49, eight; Waihi Gorge bush, 21/1/49, 22 (in numbers one to four, some young).—(D.N.F.C.) Dunedin: A pair searching for a nesting site in garden late July and August, 1951. Last season was the first for several years that a nest had not been located in the garden.—(E.M.M.) Numerous along the edge of the beech forest beside the river at Cascade Creek, 29/3/51.—(K.V.F.) Several heard in bush in lower part of Hidden Falls River, 29/12/49.—(W.A.W.)

N.Z. PIPIT (*Anthus novaeseelandiae*).—North Road, Clevedon, 24/11/50, nest, two eggs; 12/12/50, nest, three eggs.—(A.J.G.) Miranda, 30/12/50, nest, two eggs.—(N.M.) Karori Cemetery, Wellington, 23/10/50, one on tombstone.—(A.A.B.) Lake Brunner, 2/6/51, six.—(W.C.C.) Hae Hae Te Moana Gorge, 17/1/49, five in two miles.—(D.N.F.C.)

FERNBIRD (*Bowdleria punctata*).—Ruskapu, 13/2/51, one seen at least two others heard, in lupin on sand dunes.—(N.M.) Great Barrier, 14/11/50, heard and seen in Kaitoki swamp.—(J.C.D. and B.D.H.) Minginui, Nov., 1950, to June, 1951, one to four often seen; 12/5/51, 12

seen; as usual with this species, many more heard than seen.—(R.St.P.) Tauranga Harbour (near), Jan., 1951, many; also in swamp in hills, 12 miles away.—(A.S.W.) Lake Taupo, eastern, 30/10/50, several heard.—(R.H.D.S.) Wairarapa Lake, N. end, 23/10/49, heard.—(R.H.D.S.) Mr. James A. Sinclair, Deadman's, Lake Te Anau, states that the fernbird is present at the barrier in Pike Valley. He has seen three specimens which frequent one locality and are very tame. Reported to me by H. D. Skinner.—(R.A.F.)

GREY WARBLER (*Pseudogerygone igata*).—c. 20, Reservoir Reserve, Dunedin, 13/6/50; c. 30 of a mixed flock of grey warblers and brown creepers. Birds began to disappear before counts of the separate species could be taken, but they appeared to be about even, 17/6/50. On 12/4/51, heard and seen in nine localities in the Reservoir Reserve, c. 50 plus.—(I.T.) Stirling, 18/2/51, young one in orchard; 0/4/51, half-song or squeaks; 19/4/51, full song.—(R.V.McL.) Three or four singing near flats in lower part of Hidden Falls River, 29/12/49.—(W.A.W.)

YELLOW-BREASTED TIT (*Petroica m. macrocephala*).—Nelson Reservoir to Dun Mt., 15/4/51, six seen (one female), ten others heard.—(R.B.St.P.) Lake Brunner, 2/6/51, three; Anti-Crow Hut, Upper Waimakiriri River, 11/5/51, six plus.—(W.C.C.) Okuku Pass, Mt. Grey, 18/8/51, one.—(D.E.C.) Te Moana Gorge, 17/1/49, seven males, songs heard in other localities; Waihi Gorge, 21/1/49, two females and three males, one of latter young; Lynn Creek, Mt. Peel, 19/1/49, two males, one female.—(D.N.F.C.) Hilderthorpe, N. Otago, 9/1/49, one seen within 3 ft., had pure white breast.—(C.W.McL.) Song, Ross Creek, Wakari Road, 16/8/50; song, Reservoir Reserve, 12/9/50; male flew in front of car on road, Akatore to Waihola, 21/11/50; song, Glen Leith, 6/1/51; song from trees in garden, 9/3/51; female seen in garden, 20/3/51; three males seen, Reservoir Reserve, 12/4/51.—(I.T.) Trotter's Gorge, 11 males, one female, 27/3/51; Waipori Gorge, seven males, one female, 1/4/51.—(I.T., L.E.W.) Lower Taieri Gorge, four males, 3/3/51.—(K.V.F., I.T., L.E.W.) On flats in lower part Hidden Falls Valley, 29/12/49.—(W.A.W.) Titiroa Range, Dec., 1950, frequently seen.—(C.B.T., J.S.E.)

PIED TIT (*P. m. toitoi*).—Waitakeres, 26/3/51, one heard at Dreamland.—(R.E.S.) 15/5/51, two singing, Huia Valley.—(K.C.B.C.) 7/7/51, one at Cascades.—(N.M.) Minginui, Nov., 1950 to June, 1951, up to seven daily at bush workings; up to 28 on trips to deep bush and high country.—(R.St.P.) Mt. Moehau, May, 1950, one seen at 1200 ft.—(J.S.E.) South of Atiamuri, in pinus insignis forest, 30/10/50, male heard.—(R.H.D.S.)

ROBIN (*Miro australis*).—Minginui, Nov., 1950, to June, 1951, two usually follow the bushmen or logging tractors. Many in the higher beech country and in the dark gullies of the wilder parts. In the first hour's walk up a tributary of Waiau, I saw 18.—(R.St.P.) Bush near Tauranga, Jan., 1951, one seen.—(A.S.W.) Nelson Reservoir to Dun Mt., 15/4/51, two seen, two others heard.—(R.B.St.P.) Travers River, 7/1/51, two.—(W.C.C.) Rahu Saddle, Feb., 1951.—(G.G.) Plentiful Piano Flat, Waikaia Valley, and in the bush between the valley and Titan Rocks, 1 to 7/1/50.—(E.M.M.) Titiroa Ranges, Dec., 1950, five or six seen in three days; present in beech forest along Waiau River, very friendly.—(C.B.T. and J.S.E.)

FANTAIL (*Rhipidura fuliginosa*).—Howick, one pied flew into house during day and stayed in all night, appeared not to be able to see in dark or half-light as it was handled easily then. Slept with head tucked into wing, 30/5/51.—(N.M.) North Road, Clevedon, 9/11/50, one pied flew round and round a shining cuckoo, then pursued it when it flew.—(A.J.G.) Minginui, 1950-51, plentiful; none of the black phase seen here.—(R.St.P.) Nelson Reservoir to Dun Mt., 15/4/51, 20 seen, only one black.—(R.B.St.P.) Very common about Aranui School, Christ-

church, June, July, 1951.—(W.C.C.) Geraldine district, Jan., 1949, count over nine days, 27 pied, seven black, three black seen together in garden.—(D.N.F.C.) Count, June 1, 1950 to May 31, 1951: Garden, Maori Hill, pied 101, black 32; outside of garden, pied 75, black 15.—(I.T.)

WHITEHEAD (*Mohoua o. albicilla*).—Little Barrier Is., 12-25/11/50, three nests found, one with three added eggs, one with four eggs, only one of which hatched, chick leaving nest safely; one with two eggs, which hatched, but the chicks died in the nest when 12 days old.—(C.H.P., J.W.St.P., W.P.M. and H.R.McK.) Minginui, Nov., 1950 to June, 1951, up to 40 seen daily; in high back country up to 400 daily. From 23/1/51, seen feeding flying young.—(R.St.P.)

BROWN CREEPER (*Finschia novaeaeelandiae*).—34 in flock before a snow fall Anti-Crow Hut, Upper Waimakiriri River, 12/5/51.—(W.C.C.) Reservoir Reserve, Maori Hill, mixed flock creepers and grey warblers, c. 30, 17/6/50; pair Ross Creek Track, 16/8/51; heard three localities, Reservoir Reserve, 12/9/50; Akatore, 19/9/50; calls, Botanic Gardens, 20/10/50; six times during March, 1951, heard or seen, sometimes in little flocks in garden.—(I.T.) Pair, Woodside Glen, repeated song as if nesting, 23/9/50; flock, bush, Fish Hatcheries, Waitata, 21/4/51.—(D.N.F.C.) Lower Taieri Gorge, 3/3/51, small flock.—(O.R.C.)

SILVER EYE (*Zosterops lateralis*).—Moumoukai, Clevedon, 1951, winter, several flocks of up to 50, a good increase from last year.—(J.W.St.P.) Minginui, Nov., 1950, to June, 1951: Nov. to Feb. small lots in lower country, average 10 per day. March, some increase in lower area and flocks in high country, up to 250 birds seen in one day. In April and May the lower increased to a maximum of 300 in one day though many remained in the high country. On June 23rd the bushmen reported many hundreds which had apparently been driven from the heights by a cold S.E. gale. In greater numbers and in better condition than in the winter of 1950.—(R.St.P.) Thirty-five plus in flock, Lake Brunner, 3/6/51.—(W.C.C.) Hilderthorpe, N. Otago, feeding at table, 28/3/49, two; 10/4/49, five or six; 12/4/49, 30; June, 1949, 200 to 300; 21/7/49, 30; 10/9/49, many; 20/4/50, 12; 7/7/50, 200 to 300; 23/7/50, 12; 7/8/50, 20 to 30.—(C.W.McL.) June, 1950, the largest number of silver-eyes ever known constantly about the garden; rough estimates (accurate counts impossible) on and about bird tables and nearby trees, 50 to 100; largest consumption of syrup ever recorded, 16½ quart bottles being the peak on June 24, 1950. Absence of berries on native trees and shrubs. (June, 1951, peak so far this year, 3½ bottles).—(I.T.)

TUI (*Prothemadera novaeaeelandiae*).—Little Barrier Is., 15/11/50, one out of nest; up to 24/11/50, large young about to leave nest in high macrocarpa. Many birds feeding chicks in nest. The greatest number of breeding birds apparently start laying at about the same time.—(C.H.P., J.W.St.P., W.P.M. and H.R.McK.) 25/5/51, one at Middlemore; mid-Sept., two on kowhai on Mangere Mt.; uncommon immediately to the south of Auckland.—(K.C.B.C.) Howick, 30/6/51, seven in garden, usually two or three.—(N.M.) North Road, Clevedon, 27/1/51, two eating laurel berries in garden.—(A.J.G.) Minginui, 1950-51, fairly regular, up to 40 seen in one day, average about 20. On deer-stalking trips to Whirinaki, up to 500 seen in one day. Strangely enough, none in autumn on the Waiau Stream or its tributaries, though plentiful in the lower bush and on top of the ranges. Feeding notes: Oct., on kowhai; Jan., chasing and catching cicadas; March, April, May, on fruit of rimu and kahikatea. The kahikatea crop was the heaviest for years, the branches bending with the weight of fruit. On June 24th practically all of this temporary population, numbering many thousands, moved off south-west. A bush hawk appeared in Feb. but left owing to its being chased frequently by parties of 15 to 20 tuis.—(R.St.P.) Kairimu Road, Marakopa, at least five pairs on farm (includes three small patches of bush), breeds.—(R.E.W.F.) Lake Brunner, 2/6/51, two.—(W.C.C.) Okute Valley, one; albino tui observed in bush behind Akaroa township by C.

Holmes, 10/5/51.—(D.E.C.) Geraldine Hill bush, 16/1/49, two.—(D.N.F.C.) Maori Hill, 21/2/51, young tui flew upwards about 3ft. and caught white butterfly passing over its head, brushed it against bough before eating it.—(I.T.) On the edge of the bush, Swampy Hill, Dunedin, 6/5/51, 10 to 12, the largest record in this area which is often visited.—(E.M.M.) Pounawea, Otago, 13/1/51, four feeding on rata.—(L.E.W.)

BELLBIRD (*Anthornis melanura*).—Little Barrier Is., 14/11/51, no young out of nests; 15/11/51, three chicks left nest in garden hedge; 18/11/51, three other young seen and from then many seen.—(C.H.P., J.W.St.P., W.P.M. and H.R.McK.) Minginui, 1950-51, average of c. 25 seen each day, maximum from Nov., c 40, rising to 60 in April and 100 in June. Hunting trips up Whirinaki River, March and April, max. 500 seen in one day. Great numbers feeding on rimu and kahikatea fruit March, April, May. Did not leave in June as they do not migrate so much as the pigeon and tui. Fledglings left the nest in January.—(R.St.P.) Lake Tarawera, 26/2/50, very abundant especially around the landing and at the outlet.—(M.J.S.B.) Mokoia Is., Rotorua, 4/2/51, one auditory record.—(M.J.S.B.) South of Atiamuri, in *pinus insignis* forest, 30/10/50, heard.—(R.H.D.S.) First one seen in my garden at Castlecliff this winter on 21/5/51; not many about Castlecliff this winter.—(W.P.M.) Nelson Reservoir to Dun Mt., 15/4/51, ten seen, 100 plus heard.—(R.B.St.P.) Anti-Crow Hut, Upper Waimakiriri River, 10/5/51, nine; Lake Brunner, 2/6/51, three; Lake Guyon, 3/1/51, two.—(W.C.C.) Okute Valley, 10/12/51, six.—(D.E.C.) Geraldine district, Jan., 1949, frequently heard or seen.—(D.N.F.C.) During June, 1950, counts of five to eight common, once c. 15. First young in garden, November 14. February, 1951, notes, male, female and young frequent; March, songs of females as frequent as those of males.—(I.T.) Stirling, 14/6/50, two at honey-water jar, one chased other away all day. This went on for a few days, till 20/6/50, when big one allowed other to drink. June, 1951, one comes, first visit, 5/5/51.—(R.V.McL.) Two bellbirds noted on a power wire, Blacks Road, Opoho; one had a straw in its beak and the other pecked at the straw. Did some snuggling together, caressing and playing with the straw. The one with the straw flew away and was followed by the other bird.—(W. W. McArthur, D.N.F.C.) Pounawea, Otago, 13/1/51, several feeding on ratas.—(L.E.W.)

BLUE-WATTLED CROW (*Calleas wilsoni*).—Upper Ness Valley, Clevedon, 6/1/51, one seen, calling quietly as if to a mate; rarely seen here.—(R.J.F.) Moumoukai, Clevedon, 1951, winter, several heard each time I go through bush, especially about Mt. Kohukohunui; 1/7/51, Mr. F. Gibbons heard three or four singing in the Hunua Bush.—(J.W.St.P.) Pongakawa district, 3½ miles from Rotoehu, my nephew tells me they are still to be heard here; he recently recorded seven in the one day.—(A.H.W.) Near Tauranga, Jan., 1951, two reported seen in bush area.—(A.S.W.) Mr. R. D. Larritt, Frankton Junction, reports, 1/8/51:—“While deer-stalking on the Marapara Mountain (between lakes Rotoiti and Rotoehu next to Hongi's Track) I noticed two birds. I stood still and watched them, being keen to get a good look since I had not seen their like before. They were in size about half-way between a black-bird and a pigeon; their colour was a steely grey not unlike the colour of the ink that I am using to write this letter, perhaps a little darker. Their beaks were as far as I can remember (it was a dull day in virgin bush last Easter) were black and an inch to one and a half-inches long. They came right down to within about 8 or 10 feet and we had a good look at their movements in the wineberry and supplejack. They were quite quick in their movements. They would run quickly along a branch or twig and seemed to hop from branch to branch; when running along a branch they reminded me of a parrot but the movement was much quicker. They made no sound and left us after they had had a good look. I believe they may be a very rare native crow.”—(R.A.F.) Toa Toa district, Bay of Plenty, 1951, specimen brought to Opotiki.—(J.D.C.)

GREENFINCH (*Chloris chloris*).—Minginui, Nov., 1950, to June, 1951, Nov. to March, very small lots; April, starting to flock; May, flocks of c 200 and up to c 450 seen in one day; June, up to 600.—(R.St.P.) Maori Hill, six or seven feeding on chickweed, 27/8/50; calling commenced 1/11/50; February-May. This season daphne berries ripened and fell. This is the first time for years that greenfinches have not stripped the shrub before the berries had time to ripen.—(I.T.)

CHAFFINCH (*Fringilla coelebs*).—Clevedon, April-May, 1951, feeding on seeds of flax (*Phormium tenax*) by holding seed in bill and shaking the wings from it.—(H.R.McK.) Minginui, Nov., 1950 to June, 1951, mostly on pumice flats, especially among the self-sown hawthorn trees; Nov. to Jan., small lots, but up to 50 seen in one day; Feb., flocking commenced, lots of 60 to 100 in old timber workings; March-June, flocks of c. 200 and up to 500 seen in one day.—(R.St.P.) Nelson City, last song heard 4/4/51; first full song for season heard 4/8/51; singing general by 8/8/51.—(L.G.)

REDPOLL (*Carduelis cabaret*).—Waerenga, during August and September, 1949, huge flocks of redpolls frequented young grass to feed on small "water grass" growing on ridges, but none remained by November.—(M.P.D.) Minginui, Nov., 1950 to June, 1951, very patchy in numbers; a few large flocks in autumn.—(R.St.P.) Atiamuri, 30/1/49, on the high ground before reaching the bridge a small flock of c 11 seen.—(M.J.S.B.) Northland, Wellington, last seen 16/10/50; four seen day 1 returned from H.B., 24/8/51; maximum, 25, on 24/9/51; last seen, 1/10/51.—(A.A.B.) Thousands on Balmacewan golf links, 18/6/50; lawn had just been sown.—(I.T.)

GOLDFINCH (*C. carduelis*).—Howick, from 26/1/51, to c 9/4/51, flocks of from 4 to 50 juveniles came every morning to feed on scabious, cornflower, golden rod and zinnia seeds. At first they had no red on heads, but by 3/3/51, they were starting to attain the scarlet colour of adult birds.—(N.M.) Moumoukai, Clevedon, July, 1951, flocks of up to 100 seen, only odd small lots earlier.—(J.W.St.P.) Minginui, Nov., 1950, to June, 1951, usually scarce, but odd flocks of up to 200, once 500.—(R.St.P.) Nelson City, first full song heard, short bursts of song from bird sitting on telegraph wires. Winter flocks have not broken up yet; 17/9/51. Birds are paired up by now, 5/10/51.—(L.G.)

SPARROW (*Passer domesticus*).—Takapuna, 16/7/51, G. E. Minhinnick reported seeing sparrow chasing a red-billed gull.—(N.M.) Minginui, Nov., 1950 to June, 1951, 100 to 500 seen daily. Flocks noted to be living in open country far from houses. Seen to take cicadas. Increasing fast.—(R.St.P.) Dunedin, 28/6/51, buff-coloured sparrow seen, ordinary dark markings replaced by cinnamon, edges of wings and upper side of tail cream coloured, eye light brown. Earlier in breeding season, peculiar young one seen being fed at bird table, probably same bird.—(L.E.W.)

YELLOWHAMMER (*Emberiza citrinella*).—Little Barrier Island, 11-25/11/51, two or more males about Titoki Point acting as if they had mates sitting.—(H.R.McK.) Eskdale, still singing, full song, 10/2/51; Nelson City, first full song heard 15/9/51.—(L.G.)

CIRL BUNTING (*E. cirius*).—Tauranga, Jan., 1951, three seen.—(A.S.W.) Seen Hawke's Bay.—(D.B.) Lincoln, Canterbury, 22/8/50, male.—(L.E.W.)

SONG THRUSH (*Turdus ericetorum*).—Howick, 14/5/51, first full song, dawn.—(N.M.) North Road, Clevedon, 1/5/51, song commenced.—(A.J.G.) Mokoia Is., Rotorua, 4/2/51, at least four in full song. Last song recorded on Rotorua mainland was 2/1/51.—(M.J.S.B.) Nelson City, first full song, 07.30 hours, 21/3/51.—(L.G.) Hilderthorpe, N. Otago, drinking milk from cat's saucer.—(C.W.McL.) Last recorded song, Maori Hill, 12/1/51; first recorded song, 25/3/51.—(I.T.) Stirling,

very plentiful; 21/6/50, song; 5/5/51, full song; several singing throughout May and June.—(R.V.McL.)

BLACKBIRD (*T. merula*).—Howick, 18/6/51, in full song.—(N.M.) North Road, Clevedon, 5/5/51, first song.—(A.J.G.) Orakei Korako, 30/1/49, around tea rooms six blackbirds in full song; singing ceased in Rotorua in last week in Dec. Mokoia Is., Rotorua, 4/2/51, at least four singing lustily; last song Rotorua mainland, 17/12/50; a male observed with billful of earthworms.—(M.J.S.B.) Nelson City, first full song heard, still lacking the volume of later season song, but stronger than sub-song and consisting of all the phrases of full song. Delivered from telegraph pole, 17.45 hours 22/7/51; singing is general in the evenings but still weak, 30/7/51.—(L.G.) First recorded song, Maori Hill, 24/7/50. Full, loud, eight-phrase song sung repeatedly at twilight, 12/5/51. On 31/3/51, great chasing about garden; this happens yearly about this date; full song repeatedly about sunset, 19/5/51.—(I.T.) Report of blackbird heard singing during the night in the Taieri, 25/9/50, and heard singing Morningside, 12/6/51.—(Miss T. Jackson, D.N.F.C.) Blackbird with white neck, Ravensbourne, 18/6/51.—(Mrs Campbell, D.N.F.C.) Stirling, very plentiful; 14/4/51, chasing; one attacked kingfisher; 2/6/51, weak song at sunset while at least four thrushes singing loudly; 23/6/51, distinct song at dawn; 30/6/51, quiet song several times while two thrushes sang loudly, sunset.—(R.V.McL.) Dunedin, nesting in garden, and on 10/10/50 hen bird flew out of nest and dropped a dead baby bird at my feet, stood and watched till I picked it up, when she immediately began to feed. First whisper song heard in garden on 23/5/51 and louder whisper song on 29/5/51. On 25/3/51, a male bird executed a dance round and round a currant bush, after a brief rest resumed the dance, repeated several times.—(L.E.W.)

HEDGE SPARROW (*Prunella modularis*).—Hora Hora Islet, off Waikeke, 14/10/50, one singing in taupata scrub.—(H.R.McK.) Minginui, Nov., 1950 to June, 1951, up to six in one day in summer and up to 35 in autumn.—(R.St.P.) Song recorded 17/6/50, but daily singing not commenced until end of July. January 13, 1951, last recorded song of season.—(I.T.) Stirling, 28/6/51, one singing; 1/7/51, two sang alternately in orchard, mid-day.—(R.V.McL.)

SKYLARK (*Alauda arvensis*).—Nelson City, lark singing on wing, mid-morning, 11/8/51; cold but sunny, quite long bursts of song as it flew overhead and continued until out of sight. This was not territorial soaring flight, but rather song on the wing as the bird moved from one locality to another; similar occurrence 15/8/51.—(L.G.) Increasing again at Waitahuna, where they had for some time been scarce, Jan., 1951.—(Miss A. Clancy, D.N.F.C.) Stirling, 18/4/51, singing.—(R.V.McL.)

STARLING (*Sturnus vulgaris*).—Minginui, Nov., 1950 to June, 1951, Nov. up to 250; Dec., 400; Jan., 300; Feb. 9; March, 70. but two lots of 800 and 500; April, 30, but one lot of c 4,000; May, 80, but two lots of c 1000 and one of c 4000; June, 80 and one lot of c 1000. The large flocks seemed to be distinct from the usual population from March to June. In March and April they fed on kahikatea and rimu berries and on grass-grubs on the flats. When they moved the roar of their wings was audible for a quarter of a mile.—(R.St.P.) Dunedin, starling seen to pick up a sparrow; shook it, jerking it about five times; when released the sparrow fell to the ground where it lay with wings outspread as if stunned. It was some minutes before it recovered sufficiently to be able to fly to cover; 16/6/51.—(Miss J. Stewart, D.N.F.C.) Stirling, 22/2/51, large flock flying west across farm; 5/5/51, hundreds on belt of hawthorn trees, towards sunset, singing; flew on towards night roost.—(R.V.McL.) Interest in sparrows' old nests resumed 3/9/50. Maori Hill; young paid first visits to bird table, 22/12/50. May, 1951, starling for the first time seen eating apples from the trees in garden.—(I.T.)

MYNA (*Acridotheres tristis*).—Howick, 17/3/51, eight in garden; now plentiful.—(N.M.) Lake Rotorua, 8/11/50, single bird seen at the six-mile peg, Rotorua-Tauranga main highway.—(M.J.S.B.) 2/1/51, on a drive from Te Karaka to Hexton counted 150 on the road.—(C.K.W.) Feilding, 1951, at least five pairs in business area.—(E.D.)

WHITE-BACKED MAGPIE (*Gymnorhina hypoleuca*).—Karaka, 23/7/51, one, casual.—(D.A.U.) Ruatoria district, they arrived in this district in about 1933; now numerous and appear well settled.—(C.K.W.) Geraldine district, heard from dawn till dusk, so common that only flocks over 20 were recorded. On Jan. 17, near Te Moana, a flock of 30 to 40; Jan. 28, near Kakahu, flock of 28.—(D.N.F.C.) Stirling, none in district.—(R.V.McL.) For some years usual to see about half-dozen at Morrison's, near Dunback; seen 10-15/1/51.—(F.R.B.)

BLACK-BACKED MAGPIE (*G. tibicen*).—Between Dannevirke and Woodville, 13/11/50, 1; Martinborough, 5/9/51, one.—(J.M.C.)

MAGPIE (*Gymnorhina* ? sp.).—Parakai, near Helensville, 9/7/51, one with black band across upper back.—(R.E.S.)

HERONS IN CANTERBURY.

By H. Barker, Kaiapoi.

A waste of mud, water and rushes and a flooded gateway, and on either of the gate's two supporting posts, a heron—pale greyish blue, almost white—very upright, tall and slender, with head well back and long necks curved forward.

They regarded my approach with more curiosity than alarm, and from a distance of about thirty yards I was able to watch them as they stood motionless like two exquisite figures in porcelain. This was my first view of these unusual visitors to my farm at Kaiapoi, and incidentally the only time they allowed me to approach them closely. We have been long familiar with both the blue and white herons, but these birds were obviously different, and it was some time before I was able to identify them as the white-fronted heron (*matuku moana*) a species of which I had never heard before in this district.

This was February 25, 1951, and for several days these birds were a familiar sight as they waded, fishing in the shallow waters of the swamp; and then they were joined by two more. After that, progress was rapid, and by March 15 there were eleven. The whole family was interested by this time, and competition was keen in establishing a new count, which was not hard to do as they kept on coming. By the end of March there were thirty-six—but that seemed to be the limit.

Owing to the very wet summer, the fields were all more or less flooded, and everywhere there were countless thousands of frogs (they even invaded the house at night if the doors were left open) and every pool and every puddle was soon swarming with tadpoles, so even when our thirty-six herons were joined by three white ones, the food supply was almost unlimited. The herons seemed to make our swamp their headquarters, and in the mornings could be seen flying off in groups to prove the surrounding country. Their habits seemed quite regular. In the morning, just before sunrise, they would come up from the swamp and settle along the tops of a row of tall pines close to the house. There, in the first rays of the sun, they would make a leisurely toilet, preening their feathers and sunning themselves for about half an hour, then, in small groups, would drift off to an adjacent pond to wade and play for an hour or so before moving further afield. In the evening the same order was observed, and we used to watch about half an hour before sunset for them coming in to perch on the tree-tops in the last sunlight,