

A LIST OF THE CHEWING LICE (INSECTA: MALLOPHAGA) FROM BIRDS IN NEW ZEALAND

by

R.L.C. PILGRIM

Department of Zoology, University of Canterbury
Christchurch, New Zealand

and

R.L. PALMA

National Museum of New Zealand
Wellington, New Zealand

Jointly published as
SUPPLEMENT TO NOTORNIS VOLUME 29

and

NATIONAL MUSEUM OF NEW ZEALAND
MISCELLANEOUS SERIES No. 6

by

Ornithological Society of New Zealand
and
National Museum of New Zealand

SEPTEMBER 1982

A LIST OF THE CHEWING LICE (INSECTA: MALLOPHAGA) FROM BIRDS IN NEW ZEALAND

By R. L. C. PILGRIM and R. L. PALMA

ABSTRACT

Authenticated records of chewing (feather) lice, largely from unpublished observations, are listed from birds in the New Zealand zoogeographic subregion, including the Ross Dependency in Antarctica. From a total of 388 host taxa (including 38 introductions), lice are reported from 250 hosts (including 27 introductions). These records comprise 267 lice species and subspecies; a further 138 records are listed at generic level only, but all these do not necessarily represent separate species. Although some bird groups have been collected from frequently, there are many gaps in our knowledge, even for these hosts. With some orders (e.g. Passeriformes) and families of birds, little or no attention has been paid to their ectoparasites. An appeal is made for assistance in building up collections.

INTRODUCTION

The New Zealand bird fauna is broadly characterised by the endemic order Aptygiformes and, in comparison with other faunas, the exceptionally high proportion of taxa in the orders Procellariiformes and Sphenisciformes and in the family Phalacrocoracidae (Pelecaniformes) (Pilgrim, 1980). Since the genera of chewing (feather) lice are, in general terms, associated with particular groups of hosts, the louse fauna shows correspondingly a disproportionate representation of lice on the foregoing host groups. The louse genus *Apterygon*, which is found on all members of the Aptygiformes, is the sole endemic genus.

Some endemic genera of host birds, such as *Notornis*, *Strigops*, *Nestor*, *Cyanoramphus*, *Hymenolaimus*, have lice peculiar to them; the latter constitute one of the major characteristics distinguishing the New Zealand louse fauna from those of other regions. Among some orders, e.g. Anseriformes and Charadriiformes, many host species have a wide distribution in one or both hemispheres and their lice are thus common to many countries. However, on endemic hosts within these orders the lice may also be endemic: Martens & Palma (1981) showed that *Quadraceps cedemajori*, *Q. novaeelandiae* and *Q. dominella* are confined to New Zealand endemic plovers.

Lice from Passeriformes are inadequately known in New Zealand, even from introduced birds. Some native hosts in this order carry lice which are referable to widespread passerine louse genera such as *Menacanthus*, *Myrsidea*, *Brueelia*, *Philopterus*, but undescribed species are probably present.

The phylogenetic relations of lice often parallel those of their hosts and may help both to elucidate the relationships of the hosts and to distinguish closely related host taxa which are otherwise poorly defined. Only those lice which are positively established on a host can be considered in such interpretations. A louse taxon can be regarded as positively established on a certain host taxon when both taxa have been found associated on several occasions. Secondary infestations may result in well-established parasite species, but they often occur on hosts which are

clearly only distantly related and cannot then be used as evidence of host phylogeny. Outstanding examples of secondary infestations include a louse species belonging in a penguin-louse genus (*Austrogoniodes*) but parasitising a duck in Australia (Clay, 1971a), and a louse genus belonging in a marsupial-louse family (Boopidae) but established on the cassowary in Papua (Clay, 1971b). Less obvious and therefore more difficult to interpret is the genus *Rallicola*, which parasitises all species and subspecies of kiwis, as well as several other avian orders including the Gruiformes (Clay, 1953).

Wise (1977) compiled a list of the Mallophaga from the New Zealand subregion, including only those species which had been previously published. Lists of lice from limited areas within the subregion have been published by Clay, 1964 (Campbell Island); Clay & Moreby, 1970 (Macquarie Island); Horning, Palma & Pilgrim, 1980 (Snares Islands); Watson, 1967 (Macquarie Island); and Watt, 1971 (Kermadec Islands). Our aim is to provide a preliminary checklist of the lice known to occur on birds in the New Zealand subregion, incorporating numerous unpublished observations made on collections which have become available to us. The area defined in this survey is that shown in the 1970 *Checklist* of the Ornithological Society of New Zealand (maps 1-2, including Macquarie Island). The numbers in the list and the names of the hosts are those used in the *Checklist* as amended and added to in the Ornithological Society of New Zealand's *Amendments and Additions* (1980).

We include only those lice species of which we have seen specimens (except where noted). Unconfirmed literature reports, misidentifications, junior synonyms, stragglers and contaminants (see below) have been excluded. In all cases the list includes records only from hosts taken within the subregion. Explanatory notes, referred to in square brackets, have been added when more detailed explanations are necessary; the notes appear after the list.

Wherever lice are recorded in this paper as a generic name followed by "sp", it means either that we have had insufficient material or opportunity to compare our specimens with the types or other authenticated specimens, or that the genus requires revision (e.g. *Ancistrona*, *Longimenopon*). Sometimes, new species are almost certainly involved, and these will be the subject of future papers.

The abbreviation *s. l.* (= *sensu lato*) is used to denote populations with a range of measurement and morphological variation at present considered different from the named species but not sufficiently distinct to warrant the erection of separate taxa.

The two following categories of reports are deliberately excluded from the list:

1. *Stragglers*: Lice which are found occasionally on a bird but which appear not to be regular parasites of that host, even though the occurrence is from field collecting. For example, predatory birds may, at least temporarily, acquire lice from their prey, and birds breeding on the ground may become infested by lice from other species breeding nearby.
2. *Contaminants*: Lice which have transferred to a bird through careless handling and contact with another host species or even through placing the bird in a bag, on a museum tray, etc., previously occupied by another host species. We can vouch for a recent instance in which an Erect-crested Penguin yielded several lice of two species characteristic of the Pukeko — inquiries revealed that the penguin had in fact been caged overnight with a Pukeko! This is an accident arising from careless mishandling and is a situation to be rigorously avoided.

Many birds have been omitted from the list because they are not known to have had lice collected from them in the subregion; we expect that lice will be found on most when opportunities for examining them arise. As well, even for the hosts that are included, we do not suggest that the list of lice be regarded as complete since in many cases further lice species are known to occur on the hosts in other parts of the world.

In New Zealand, the major collections of bird lice are located in the Entomology Division, DSIR, Auckland; the National Museum of New Zealand, Wellington; and the "R.L.C. Pilgrim Collection" housed in the Department of Zoology, University of Canterbury, Christchurch (ultimately to be deposited in the National Museum of New Zealand). Smaller, less comprehensive collections are maintained at several other museums and by Government departments.

We hope that this paper will draw attention to our interest in this group and will encourage ornithologists to collaborate with us in obtaining specimens, leading to a fuller representation of the mallophagan fauna. Suggestions for collecting lice are given in Appendix 1.

This paper includes literature reports and hitherto unpublished identifications up to December 1981.

LIST OF BIRDS AND THEIR LICE

(*denotes a host introduced to New Zealand by human agency)

Order APTERYGIFORMES
Family APTERYGIDAE

- 1.a. **Apteryx australis mantelli** Bartlett, 1850 **North Island Brown Kiwi**
Apterygon mirum Clay, 1961 [Note 1]
Rallicola (Apterocola) sp. [Note 2]
- 1.b. **Apteryx australis australis** Shaw & Nodder, 1813 **South Island Brown Kiwi**
Apterygon dumosum Tandan, 1972 s. l.
Apterygon sp.
Rallicola (Apterocola) gadowi Harrison, 1915
- 1.c. **Apteryx australis lawryi** Rothschild, 1893 **Stewart Island Brown Kiwi**
Apterygon dumosum Tandan, 1972
Rallicola (Apterocola) gadowi Harrison, 1915 s. l.
2. **Apteryx owenii** Gould, 1847 **Little Spotted Kiwi**
Apterygon dumosum Tandan, 1972 s. l.
Rallicola (Apterocola) gadowi Harrison, 1915 s. l.
Rallicola (Apterocola) piligrimi Clay, 1972
3. **Apteryx haastii** Potts, 1872 **Great Spotted Kiwi**
Apterygon hintoni Clay, 1966
Rallicola (Apterocola) gracilentus Clay, 1953

Order SPHENISCIFORMES
Family SPHENISCIDAE

- | | |
|--|---|
| <p>4. Aptenodytes forsteri Gray, 1844
<i>Austrogoniodes mawsoni</i> Harrison, 1937</p> <p>5. Aptenodytes patagonicus Miller, 1778
<i>Nesiotinus demersus</i> Kellogg, 1903</p> <p>6. Megadyptes antipodes (Hombron & Jacquinot, 1841)
<i>Austrogoniodes concii</i> (Kéler, 1952) s. l.
<i>Austrogoniodes</i> sp.</p> <p>7. Pygoscelis adeliae (Hombron & Jacquinot, 1841)
<i>Austrogoniodes antarcticus</i> Harrison, 1937</p> <p>10.a. Eudyptula minor iredalei Mathews, 1911
<i>Austrogoniodes waterstoni</i> (Cummings, 1914)</p> <p>10.b. Eudyptula minor variabilis Kinsky & Falla, 1976
<i>Austrogoniodes waterstoni</i> (Cummings, 1914)</p> <p>10.c. Eudyptula minor albosignata Finsch, 1874
<i>Austrogoniodes waterstoni</i> (Cummings, 1914)</p> <p>White-flipped Blue Penguin</p> <p>10.d. Eudyptula minor minor (Forster, 1781)
<i>Austrogoniodes waterstoni</i> (Cummings, 1914)</p> <p>Southern Blue Penguin</p> <p>10.e. Eudyptula minor chathamensis Kinksy & Falla, 1976
<i>Austrogoniodes waterstoni</i> (Cummings, 1914)</p> <p>Chatham Island Blue Penguin</p> <p>12.a. Eudyptes chrysocome chrysocome (Forster, 1781) [Note 3]
<i>Austrogoniodes cristati</i> Kéler, 1952
<i>Austrogoniodes hamiltoni</i> Harrison, 1937
<i>Austrogoniodes macquariensis</i> Harrison, 1937</p> <p>Rockhopper Penguin</p> <p>13.a. Eudyptes chrysolophus chrysolophus Brandt, 1837
<i>Austrogoniodes cristati</i> Kéler, 1952
<i>Austrogoniodes macquariensis</i> Harrison, 1937 s. l.
<i>Austrogoniodes</i> sp.</p> <p>Macaroni Penguin</p> <p>13.b. Eudyptes chrysolophus schlegeli Finsch, 1876
<i>Austrogoniodes cristati</i> Kéler, 1952
<i>Austrogoniodes hamiltoni</i> Harrison, 1937
<i>Austrogoniodes macquariensis</i> Harrison, 1937</p> <p>Royal Penguin</p> <p>14. Eudyptes pachyrhynchus Gray, 1845 [Note 4]
<i>Austrogoniodes concii</i> (Kéler, 1952) s. l.
<i>Austrogoniodes cristati</i> Kéler, 1952</p> <p>Fiordland Crested Penguin</p> | <p>Emperor Penguin</p> <p>King Penguin</p> <p>Yellow-eyed Penguin</p> <p>Adelie Penguin</p> <p>Northern Blue Penguin</p> <p>Cook Strait Blue Penguin</p> <p>White-flipped Blue Penguin</p> <p>Southern Blue Penguin</p> <p>Chatham Island Blue Penguin</p> <p>Rockhopper Penguin</p> <p>Macaroni Penguin</p> <p>Royal Penguin</p> <p>Fiordland Crested Penguin</p> |
|--|---|

- 14.1 **Eudyptes robustus** Oliver, 1953 [Note 5] Snares Crested Penguin
Austrogoniodes concii (Kéler, 1952) *s. l.*
Austrogoniodes cristati Kéler, 1952

14.2 **Eudyptes sclateri** Buller, 1888 [Note 6] Erect-crested Penguin
Austrogoniodes concii (Kéler, 1952) *s. l.*
Austrogoniodes cristati Kéler, 1952
Austrogoniodes hamiltoni Harrison, 1937 [Note 7]
Austrogoniodes struthaeus Harrison, 1915 [Note 7]

Order PODICIPEDIFORMES
Family PODICIPEDIDAE

15. ***Podiceps cristatus australis*** Gould, 1844 **Southern Crested Grebe**
Aquanirmus sp.

16. ***Podiceps rufopectus*** Gray, 1843 **New Zealand Dabchick**
Aquanirmus australis Kettle, 1974

Order PROCELLARIIFORMES
Family DIOMEDEIDAE

- 18.a. **Diomedea exulans exulans** Linnaeus, 1758 **Wandering Albatross**

 - Austromenopon affine* (Piaget, 1890)
 - Docophoroides brevis* (Dufour, 1835)
 - Episbates pederiformis* (Dufour, 1835)
 - Harrisoniella hopkinsi* Eichler, 1952
 - Paraclisis hyalina* (Neumann, 1911)

18.b. **Diomedea exulans chionoptera** Salvin, 1895 [Note 8] **Snowy Albatross**

 - Austromenopon affine* (Piaget, 1890)
 - Docophoroides brevis* (Dufour, 1835)
 - Harrisoniella hopkinsi* Eichler, 1952
 - Paraclisis hyalina* (Neumann, 1911)
 - Perineus concinnoides* Kéler, 1957

- 19.a. *Diomedea epomophora sanfordi* (Murphy, 1917) **Northern Royal Albatross**

 - Austromenopon affine* (Piaget, 1890)
 - Docophoroides brevis* (Dufour, 1835)
 - Episbates pederiformis* (Dufour, 1835)
 - Harrisoniella hopkinsi* Eichler, 1952
 - Paraclisis hyalina* (Neumann, 1911)

- 19.b. **Diomedea epomophora epomophora** Lesson, 1825 Southern Royal Albatross

 - Austromenopon affine* (Piaget, 1890)
 - Docophoroides brevis* (Dufour, 1835)
 - Episbates pederiformis* (Dufour, 1835)
 - Harrisoniella hopkinsi* Eichler, 1952
 - Paraclisis hyalina* (Neumann, 1911)
 - Perineus concinnoides* Kéler, 1957

- 20.a. Diomedea melanophrrys melanophrrys Temminck, 1828** **Black-browed Mollymawk**
- Austromenopon navigans* (Kellogg, 1896)
 - Docophoroïdes harrisoni* Waterston, 1917
 - Docophoroïdes simplex* (Waterston, 1914)
 - Harrisoniella ferox* (Giebel, 1867)
 - Paraclisis diomedaeae* (J. C. Fabricius, 1775)
 - Perineus circumfasciatus* Kéler, 1957
- 20.b. Diomedea melanophrrys impavida Mathews, 1912** **New Zealand Black-browed Mollymawk**
- Austromenopon pinguis* (Kellogg, 1896)
 - Docophoroïdes simplex* (Waterston, 1914)
 - Harrisoniella ferox* (Giebel, 1867)
 - Paraclisis diomedaeae* (J. C. Fabricius, 1775)
 - Perineus circumfasciatus* Kéler, 1957
 - Saemundssonia* sp.
- 21. Diomedea chrysostoma Forster, 1785** **Grey-headed Mollymawk**
- Austromenopon navigans* (Kellogg, 1896)
 - Austromenopon pinguis* (Kellogg, 1896)
 - Docophoroïdes simplex* (Waterston, 1914) s. l.
 - Paraclisis diomedaeae* (J. C. Fabricius, 1775)
 - Perineus circumfasciatus* Kéler, 1957
 - Saemundssonia* sp.
- 22. Diomedea chlororhynchos Gmelin, 1789** **Yellow-nosed Mollymawk**
- Perineus circumfasciatus* Kéler, 1957
- 23. Diomedea bulleri Rothschild, 1893** **Buller's Mollymawk**
- Austromenopon bulleri* Price & Clay, 1972
 - Austromenopon navigans* (Kellogg, 1896)
 - Austromenopon pinguis* (Kellogg, 1896)
 - Docophoroïdes harrisoni* Waterston, 1917
 - Harrisoniella ferox* (Giebel, 1867)
 - Paraclisis diomedaeae* (J. C. Fabricius, 1775)
 - Perineus circumfasciatus* Kéler, 1957
- 24.a. Diomedea cauta cauta Gould, 1841** **White-capped Mollymawk**
- Austromenopon navigans* (Kellogg, 1896)
 - Austromenopon pinguis* (Kellogg, 1896)
 - Docophoroïdes harrisoni* Waterston, 1917
 - Harrisoniella ferox* (Giebel, 1867)
 - Paraclisis diomedaeae* (J. C. Fabricius, 1775)
 - Perineus circumfasciatus* Kéler, 1957
- 24.b. Diomedea cauta salvini (Rothschild, 1893)** **Salvin's Mollymawk**
- Austromenopon navigans* (Kellogg, 1896)
 - Austromenopon pinguis* (Kellogg, 1896)
 - Docophoroïdes harrisoni* Waterston, 1917
 - Harrisoniella ferox* (Giebel, 1867)
 - Paraclisis diomedaeae* (J. C. Fabricius, 1775)
 - Perineus circumfasciatus* Kéler, 1957

- 24.c. **Diomedea cauta eremita** (Murphy, 1930) Chatham Island Mollymawk
Docophoroides harrisoni Waterston, 1917
Paraclisis diomedae (J. C. Fabricius, 1775)
Perineus circumfasciatus Kéler, 1957
26. **Phoebetria palpebrata** (Forster, 1785) Light-mantled Sooty Albatross
Austromenopon pinguis (Kellogg, 1896)
Docophoroides simplex (Waterston, 1914) s. l.
Paraclisis diomedae (J. C. Fabricius, 1775) s. l.
Perineus circumfasciatus Kéler, 1957 [Note 9]
Saemundssonia sp.

Family PROCELLARIIDAE

27. **Macronectes halli** Mathews, 1912 Northern Giant Petrel
Austromenopon ossifragae (Eichler, 1949)
Docophoroides murphyi (Kellogg, 1914)
Paraclisis obscura (Rudow, 1869)
Perineus sp.
Saemundssonia gaini (Neumann, 1913)
- 27.1 **Macronectes giganteus** (Gmelin, 1789) Southern Giant Petrel
Austromenopon ossifragae (Eichler, 1949)
Docophoroides murphyi (Kellogg, 1914)
Paraclisis miriceps (Kellogg & Kuwana, 1902) s. l.
Paraclisis obscura (Rudow, 1869)
Perineus sp.
Saemundssonia gaini (Neumann, 1913)
28. **Fulmarus glacialisoides** (Smith, 1840) Antarctic Fulmar
Ancistriona sp.
Austromenopon brevifimbriatum (Piaget, 1880)
Perineus nigrolimbatus (Giebel, 1874)
Saemundssonia bicolor (Rudow, 1870)
29. **Thalassoica antarctica** (Gmelin, 1789) Antarctic Petrel
Ancistriona sp.
Austromenopon brevifimbriatum (Piaget, 1880)
Pseudonirmus lugubris (Taschenberg, 1882)
Saemundssonia antarctica (Wood, 1937)
- 30.a. **Daption capense australe** Mathews, 1913 Snares Cape Pigeon
Ancistriona sp.
Pseudonirmus gurlti (Taschenberg, 1882)
Saemundssonia stammeri Timmermann, 1959
- 30.b. **Daption capense capense** (Linnaeus, 1758) Cape Pigeon
Ancistriona sp.
Austromenopon brevifimbriatum (Piaget, 1880)
Pseudonirmus gurlti (Taschenberg, 1882)
Saemundssonia stammeri Timmermann, 1959

- | | | |
|-------|--|-----------------------|
| 31.a. | Pagodroma nivea nivea (Forster, 1777)
<i>Pseudonirmus charcoti</i> (Neumann, 1907)
<i>Saemundssonia antarctica</i> (Wood, 1937) | Lesser Snow Petrel |
| 31.b. | Pagodroma nivea major Bonaparte, 1856
<i>Pseudonirmus charcoti</i> (Neumann, 1907) | Greater Snow Petrel |
| 32. | Pterodroma macroptera gouldi (Hutton, 1869)
<i>Ancistriona</i> sp.
<i>Austromenopon popellus</i> (Piaget, 1890)
<i>Halipeurus (Halipeurus) procellariae</i> (J. C. Fabricius, 1775)
<i>Naubates heteroproctus</i> Harrison, 1937 [Note 10]
<i>Trabeculus schillingi</i> Rudow, 1866 s. l. | Grey-faced Petrel |
| 33. | Pterodroma lessonii (Garnot, 1826)
<i>Ancistriona</i> sp.
<i>Austromenopon popellus</i> (Piaget, 1890)
<i>Halipeurus (Halipeurus) procellariae</i> (J. C. Fabricius, 1775)
<i>Naubates</i> sp. [Note 11]
<i>Saemundssonia</i> sp.
<i>Trabeculus schillingi</i> Rudow, 1866 s. l. | White-headed Petrel |
| 34.a. | Pterodroma externa cervicalis (Salvin, 1891)
<i>Ancistriona</i> sp.
<i>Austromenopon popellus</i> (Piaget, 1890)
<i>Halipeurus (Halipeurus) noctivagus</i> Timmermann, 1960
<i>Trabeculus</i> sp. | White-naped Petrel |
| 34.b. | Pterodroma externa externa (Salvin, 1875)
<i>Austromenopon popellus</i> (Piaget, 1890)
<i>Halipeurus (Halipeurus) kermadecensis</i> (Johnston & Harrison, 1912) s. l.
<i>Trabeculus hexakon</i> (Waterston, 1914) s. l. | Juan Fernandez Petrel |
| 37. | Pterodroma inexpectata (Forster, 1844)
<i>Ancistriona</i> sp.
<i>Austromenopon popellus</i> (Piaget, 1890)
<i>Austromenopon</i> sp.
<i>Halipeurus (Halipeurus) consimilis</i> Timmermann, 1960
<i>Longimenopon</i> sp.
<i>Naubates pterodromi</i> Bedford, 1930 [Note 12]
<i>Trabeculus schillingi</i> Rudow, 1866 s. l. | Mottled Petrel |
| 37.1 | Pterodroma mollis mollis (Gould, 1844)
<i>Austromenopon popellus</i> (Piaget, 1890)
<i>Halipeurus (Halipeurus) procellariae</i> (J. C. Fabricius, 1775) s. l.
<i>Trabeculus schillingi</i> Rudow, 1866 | Soft-plumaged Petrel |
| 38. | Pterodroma brevirostris (Lesson, 1831)
<i>Ancistriona</i> sp.
<i>Austromenopon popellus</i> (Piaget, 1890) s. l.
<i>Bedfordiella unica</i> Thompson, 1937
<i>Saemundssonia pterodromae</i> Timmermann, 1959 | Kerguelen Petrel |

39. **Pterodroma neglecta** (Schlegel, 1863) **Kermadec Petrel**
Austromenopon popellus (Piaget, 1890)
Halipeurus (Halipeurus) kermadecensis (Johnston & Harrison, 1912)
Longimenopon sp.
Trabeculus fuscoclypeatus (Johnston & Harrison, 1912)
40. **Pterodroma magentae** (Giglioli & Salvadori, 1868) **Chatham Island Taiko**
Ancistrona sp.
Austromenopon popellus (Piaget, 1890) s. l.
Halipeurus (Halipeurus) procellariae (J. C. Fabricius, 1775)
41. **Pterodroma longirostris** (Stejneger, 1888) **Stejneger's Petrel**
Ancistrona sp.
Halipeurus (Halipeurus) leucophryna Timmermann, 1960
Longimenopon sp.
Trabeculus hexakon (Waterston, 1914) s. l.
42. **Pterodroma pycrofti** Falla, 1933 **Pycroft's Petrel**
Austromenopon popellus (Piaget, 1890)
Halipeurus (Halipeurus) leucophryna Timmermann, 1960
Longimenopon sp.
Trabeculus hexakon (Waterston, 1914) s. l.
- 43.b. **Pterodroma leucoptera** subspecies
Halipeurus (Halipeurus) sp.
Naubates damma Timmermann, 1961
Trabeculus hexakon (Waterston, 1914) s. l.
44. **Pterodroma cookii cookii** (Gray, 1843) **Cook's Petrel**
Ancistrona sp.
Halipeurus (Halipeurus) turtur Edwards, 1961
Naubates damma Timmermann, 1961
Trabeculus hexakon (Waterston, 1914) s. l.
45. **Pterodroma nigripennis** (Rothschild, 1893) [Note 13] **Black-winged Petrel**
Ancistrona sp.
Austromenopon sp.
Halipeurus (Halipeurus) sp. [Note 14]
Longimenopon sp.
Trabeculus hexakon (Waterston, 1914) s. l.
- 45.1 **Pterodroma axillaris** (Salvin, 1893) **Chatham Island Petrel**
Halipeurus (Halipeurus) theresae Timmermann, 1969
Trabeculus hexakon (Waterston, 1914) s. l.
46. **Halobaena caerulea** (Gmelin, 1789) **Blue Petrel**
Ancistrona sp.
Austromenopon stammeri Timmermann, 1963
Longimenopon sp.
Naubates clypeatus (Giebel, 1874)
Saemundssonia pterodromae Timmermann, 1959

47. **Pachyptila vittata vittata** (Forster, 1777) **Broad-billed Prion**
Ancistriona sp.
Austromenopon stammeri Timmermann, 1963
Longimenopon sp.
Naubates prioni (Enderlein, 1908)
Saemundssonia desolata Timmermann, 1959
48. **Pachyptila salvini salvini** (Mathews, 1912) **Lesser Broad-billed Prion**
Ancistriona sp.
Austromenopon stammeri Timmermann, 1963
Longimenopon sp.
Naubates prioni (Enderlein, 1908)
Saemundssonia desolata Timmermann, 1959
- 49.a. **Pachyptila desolata desolata** (Gmelin, 1789) [Note 15] **Antarctic Prion**
Ancistriona sp.
Longimenopon sp.
Naubates prioni (Enderlein, 1908)
- 49.b. **Pachyptila desolata banksi** (Smith, 1840) **South Georgian Prion**
Austromenopon stammeri Timmermann, 1963
Naubates prioni (Enderlein, 1908)
Saemundssonia desolata Timmermann, 1959
- 49.c. **Pachyptila desolata alter** (Mathews, 1912) **Auckland Island Prion**
Ancistriona sp.
Austromenopon stammeri Timmermann, 1963
Longimenopon sp.
Naubates prioni (Enderlein, 1908)
Saemundssonia desolata Timmermann, 1959
50. **Pachyptila belcheri** (Mathews, 1912) **Narrow-billed Prion**
Ancistriona sp.
Austromenopon stammeri Timmermann, 1963
Longimenopon sp.
Naubates prioni (Enderlein, 1908)
Saemundssonia desolata Timmermann, 1959
51. **Pachyptila turtur** (Kuhl, 1820) **Fairy Prion**
Ancistriona sp.
Austromenopon stammeri Timmermann, 1963
Longimenopon sp.
Naubates prioni (Enderlein, 1908)
Saemundssonia desolata Timmermann, 1959
- 52.a. **Pachyptila crassirostris crassirostris** (Mathews, 1912) **Fulmar Prion**
Ancistriona sp.
Austromenopon stammeri Timmermann, 1963
Naubates prioni (Enderlein, 1908)
Saemundssonia desolata Timmermann, 1959

- | | | |
|-------|--|--------------------------------|
| 52.c. | Pachyptila crassirostris pyramidalis Fleming, 1939 | Chatham Fulmar Prion |
| | <i>Austromenopon stammeri</i> Timmermann, 1963 | |
| | <i>Longimenopon</i> sp. | |
| | <i>Naubates prioni</i> (Enderlein, 1908) | |
| | <i>Saemundssonia desolata</i> Timmermann, 1959 | |
| 53. | Procellaria cinerea Gmelin, 1789 [Note 16] | Grey Petrel |
| | <i>Ancistrona</i> sp. | |
| | <i>Austromenopon popellus</i> (Piaget, 1890) | |
| | <i>Naubates fuliginosus</i> (Taschenberg, 1882) | |
| | <i>Trabeculus hexakon</i> (Waterston, 1914) | |
| 54. | Procellaria parkinsoni Gray, 1862 | Black Petrel |
| | <i>Austromenopon popellus</i> (Piaget, 1890) | |
| | <i>Naubates fuliginosus</i> (Taschenberg, 1882) | |
| | <i>Trabeculus hexakon</i> (Waterston, 1914) | |
| 55. | Procellaria westlandica Falla, 1946 | Westland Black Petrel |
| | <i>Ancistrona</i> sp. | |
| | <i>Austromenopon popellus</i> (Piaget, 1890) | |
| | <i>Naubates fuliginosus</i> (Taschenberg, 1882) | |
| | <i>Trabeculus hexakon</i> (Waterston, 1914) | |
| 56. | Procellaria aequinoctialis steadi Mathews, 1912 | White-chinned Petrel |
| | <i>Ancistrona</i> sp. | |
| | <i>Austromenopon popellus</i> (Piaget, 1890) | |
| | <i>Naubates fuliginosus</i> (Taschenberg, 1882) | |
| | <i>Trabeculus hexakon</i> (Waterston, 1914) | |
| 58. | Puffinus carneipes hullianus Mathews, 1912 | Flesh-footed Shearwater |
| | <i>Ancistrona</i> sp. | |
| | <i>Austromenopon paululum</i> (Kellogg & Chapman, 1899) | |
| | <i>Halipeurus (Halipeurus) gravis priapulus</i> Timmermann, 1961 | |
| | <i>Naubates harrisoni</i> Bedford, 1930 | |
| | <i>Saemundssonia</i> sp. | |
| | <i>Trabeculus hexakon</i> (Waterston, 1914) s. l. | |
| 59.a. | Puffinus pacificus pacificus (Gmelin, 1789) | Wedge-tailed Shearwater |
| | <i>Ancistrona</i> sp. | |
| | <i>Austromenopon paululum</i> (Kellogg & Chapman, 1899) | |
| | <i>Halipeurus (Halipeurus) mirabilis</i> Thompson, 1940 | |
| | <i>Naubates harrisoni</i> Bedford, 1930 | |
| | <i>Saemundssonia puellula</i> Timmermann, 1965 | |
| | <i>Trabeculus hexakon</i> (Waterston, 1914) s. l. | |
| 59.b. | Puffinus pacificus cuneatus Salvin, 1888 | |
| | <i>Austromenopon paululum</i> (Kellogg & Chapman, 1899) | |
| | <i>Halipeurus (Halipeurus) mirabilis</i> Thompson, 1940 | |
| | <i>Trabeculus hexakon</i> (Waterston, 1914) s. l. | |

- 59.c. **Puffinus pacificus chlororhynchus** Lesson, 1831
Austromenopon paululum (Kellogg & Chapman, 1899)
Halipeurus (Halipeurus) mirabilis Thompson, 1940
Trabeculus hexakon (Waterston, 1914) s. l.
60. **Puffinus bulleri** Salvin, 1888 **Buller's Shearwater**
Ancistrona sp.
Austromenopon paululum (Kellogg & Chapman, 1899)
Halipeurus (Halipeurus) thompsoni Edwards, 1961
Naubates harrisoni Bedford, 1930
61. **Puffinus griseus** (Gmelin, 1789) **Sooty Shearwater**
Ancistrona sp.
Austromenopon paululum (Kellogg & Chapman, 1899)
Halipeurus (Halipeurus) diversus (Kellogg, 1896)
Trabeculus hexakon (Waterston, 1914) s. l.
62. **Puffinus tenuirostris** (Temminck, 1835) **Short-tailed Shearwater**
Ancistrona sp.
Austromenopon paululum (Kellogg & Chapman, 1899)
Halipeurus (Halipeurus) diversus (Kellogg, 1896)
Trabeculus hexakon (Waterston, 1914) s. l.
- 62.2 **Puffinus puffinus puffinus** (Brunnich, 1764) **Manx Shearwater**
Halipeurus (Halipeurus) diversus (Kellogg, 1896)
Trabeculus aviator (Evans, 1912)
63. **Puffinus gavia gavia** (Forster, 1844) **Fluttering Shearwater**
Austromenopon paululum (Kellogg & Chapman, 1899)
Halipeurus (Halipeurus) spadix Timmermann, 1961
Saemundssonia sp.
64. **Puffinus huttoni** Mathews, 1912 **Hutton's Shearwater**
Austromenopon paululum (Kellogg & Chapman, 1899)
Halipeurus (Halipeurus) spadix Timmermann, 1961
Naubates harrisoni Bedford, 1930 s. l.
Trabeculus flemingi Timmermann, 1959
- 65.b. **Puffinus assimilis kermadecensis** Murphy, 1927 **Kermadec Little Shearwater**
Ancistrona sp.
Austromenopon paululum (Kellogg & Chapman, 1899)
Halipeurus (Halipeurus) placodus Edwards, 1961
Trabeculus mirabilis (Kellogg, 1896) s. l.
- 65.c. **Puffinus assimilis haurakiensis** Fleming & Serventy, 1943 **North Island Little Shearwater**
Ancistrona sp.
Austromenopon paululum (Kellogg & Chapman, 1899)
Halipeurus (Halipeurus) mundae Edwards, 1961
Trabeculus mirabilis (Kellogg, 1896) s. l.

- 65.d. **Puffinus assimilis elegans** Giglioli & Salvadori, 1869
Subantarctic Little Shearwater
Halipeurus (Halipeurus) mundae Edwards, 1961
Trabeculus mirabilis (Kellogg, 1896) s. l.
- Family HYDROBATIDAE
67. **Oceanites oceanicus oceanicus** (Kuhl, 1820) **Wilson's Storm Petrel**
Philoceanus robertsi (Clay, 1940)
68. **Garrodia nereis** (Gould, 1841) **Grey-backed Storm Petrel**
Philoceanus garrodiiae (Clay, 1940)
Saemundssonia nereis Timmermann, 1956
- 69.a. **Pelagodroma marina maoriana** Mathews, 1912 [Note 17]
White-faced Storm Petrel
Austromenopon enigki Timmermann, 1963
Halipeurus (Synnautes) pelagicus (Denny, 1842) s. l.
Longimenopon sp.
Saemundssonia marina Timmermann, 1956
- 69.b. **Pelagodroma marina albiclinis** Murphy & Irwing, 1951
Kermadec Storm Petrel
Ancistrone sp. [Note 18]
Halipeurus (Synnautes) pelagicus (Denny, 1842) s. l.
70. **Fregetta tropica** (Gould, 1844) **Black-bellied Storm Petrel**
Austromenopon sp.
Halipeurus (Synnautes) pelagicus (Denny, 1842) s. l.
Philoceanus fasciatus (Carriker, 1958)
Saemundssonia sp.
71. **Fregetta grallaria** (Vieillot, 1818) **White-bellied Storm Petrel**
Halipeurus (Synnautes) pelagicus (Denny, 1842) s. l.
- Family PELECANOIDIDAE
- 72.a. **Pelecanoides urinatrix urinatrix** (Gmelin, 1789) **Northern Diving Petrel**
Austromenopon elliotti Timmermann, 1954
Halipeurus (Halipeurus) falsus pacificus Edwards, 1961
Pelmatocerandra setosa (Giebel, 1876)
- 72.b. **Pelecanoides urinatrix exsul** Salvin, 1896 **Subantarctic Diving Petrel**
Austromenopon elliotti Timmerman, 1954
Halipeurus (Halipeurus) falsus pacificus Edwards, 1961
Pelmatocerandra setosa (Giebel, 1876)
- 72.c. **Pelecanoides urinatrix chathamensis** Murphy & Harper, 1916
Southern Diving Petrel
Austromenopon elliotti Timmermann, 1954
Halipeurus (Halipeurus) falsus pacificus Edwards, 1961
Pelmatocerandra setosa (Giebel, 1876)

Order PELECANIFORMES
Family PHAETHONTIDAE

74. **Phaethon rubricauda roseotincta** Mathews, 1926 Red-tailed Tropic Bird
Austromenopon becki (Kellogg, 1906) *s. l.*
Saemundssonia hexagona (Giebel, 1874)
- 74.1 **Phaethon lepturus dorothae** Mathews, 1913 White-tailed Tropic Bird
Saemundssonia upoluensis (Rudow, 1870)

Family PELECANIDAE

75. **Pelecanus conspicillatus conspicillatus** Temminck, 1824 Australian Pelican
Colpocephalum eucarenum Burmeister, 1838
Pectinopygus australis Thompson, 1948

Family SULIDAE

76. **Sula bassana serrator** Gray, 1843 Australasian Gannet
Eidmanniella pustulosa (Nitzsch, 1866)
Pectinopygus bassani (O. Fabricius, 1780)
77. **Sula leucogaster plotus** (Forster, 1844) Brown Booby
Eidmanniella albescens (Piaget, 1880)
Pectinopygus garbei (Pessôa & Guimarães, 1935)
78. **Sula dactylatra personata** Gould, 1846 Masked Booby
Pectinopygus annulatus (Piaget, 1880)

Family PHALACROCORACIDAE

79. **Phalacrocorax carbo novaehollandiae** Stephens, 1826 Black Shag
Eidmanniella pellucida (Rudow, 1869)
Pectinopygus gyricornis (Denny, 1842)
80. **Phalacrocorax varius varius** (Gmelin, 1789) Pied Shag
Eidmanniella pellucida (Rudow, 1869)
Pectinopygus varius Timmermann, 1964
81. **Phalacrocorax sulcirostris** (Brandt, 1837) Little Black Shag
Eidmanniella subrotunda (Piaget, 1880)
Pectinopygus setosus (Piaget, 1880)
82. **Phalacrocorax melanoleucus brevirostris** Gould, 1837 Little Shag
Eidmanniella subrotunda (Piaget, 1880)
Pectinopygus dispar (Piaget, 1880)
- 83.a. **Leucocarbo carunculatus carunculatus** (Gmelin, 1789) King Shag
Pectinopygus sp.
- 83.b. **Leucocarbo carunculatus chalconotus** (Gray, 1845) Stewart Island Shag
Eidmanniella pellucida (Rudow, 1869)
Pectinopygus sp.

- 83.c. **Leucocarbo carunculatus onslowi** Forbes, 1893 Chatham Island Shag
Pectinopygus sp.

84.a. **Leucocarbo campbelli ranfurlyi** Ogilvie-Grant, 1901 Bounty Island Shag
Pectinopygus sp.

84.b. **Leucocarbo campbelli colensoi** Buller, 1888 Auckland Island Shag
Pectinopygus carunculatus Timmermann, 1964

84.c. **Leucocarbo campbelli campbelli** (Filhol, 1878) Campbell Island Shag
Pectinopygus carunculatus Timmermann, 1964 s. l.

85. **Leucocarbo albiventer purpurascens** (Brandt, 1837) Macquarie Island Shag
Pectinopygus turbinatus (Piaget, 1890)

86.a. **Stictocarbo punctatus punctatus** (Sparrman, 1786) Spotted Shag
Eidmanniella pellucida (Rudow, 1869)
Pectinopygus punctatus Timmermann, 1964

86.b. **Stictocarbo punctatus steadi** (Oliver, 1930) Blue Shag
Eidmanniella pellucida (Rudow, 1869)
Pectinopygus punctatus Timmermann, 1964 s. l.

86.c. **Stictocarbo punctatus featherstoni** Buller, 1873 Pitt Island Shag
Pectinopygus sp.

Order CICONIIFORMES
Family ARDEIDAE

90. **Ardea novaehollandiae novaehollandiae** Latham, 1790 **White-faced Heron**
Ardeicola piligrimi Tandan, 1972
Ciconiphilus decimfasciatus (Boisduval & Lacordaire, 1835)

92. **Egretta alba modesta** (Gray, 1831) **White Heron**
Ardeicola expallidus Blagoveshtchensky, 1940
Ciconiphilus decimfasciatus (Boisduval & Lacordaire, 1835)

93. **Egretta garzetta immaculata** Gould, 1846 **Little Egret**
Ardeicola expallidus Blagoveshtchensky, 1940
Ciconiphilus decimfasciatus (Boisduval & Lacordaire, 1835)

95. **Bubulcus ibis coromandus** (Boddaert, 1783) **Cattle Egret**
Ciconiphilus decimfasciatus (Boisduval & Lacordaire, 1835)

97. **Botaurus stellaris poiciloptilus** (Wagler, 1827) **Australasian Bittern**
Ardeicola stellaris (Denny, 1842) *s. l.*
Ciconiphilus decimfasciatus (Boisduval & Lacordaire, 1835)

Family THRESKIORNITHIDAE

- | | | |
|------|---|------------------------------|
| 99. | Plegadis falcinellus (Linnaeus, 1758) | Glossy Ibis |
| | <i>Ardeicola raphaelius</i> (Nitzsch, 1866) | |
| | <i>Colpocephalum leptopygos</i> Nitzsch, 1874 | |
| | <i>Ibidoecus bisignatus</i> (Nitzsch, 1866) | |
| | <i>Plegadiphilus plegadis</i> (Dubinin, 1938) | |
| 100. | Threskiornis molucca (Cuvier, 1829) | Australian White Ibis |
| | <i>Ibidoecus dianae</i> Tandan, 1958 | |
| | <i>Plegadiphilus threskiornis</i> Bedford, 1939 | |
| 101. | Platalea leucorodia regia Gould, 1838 | Royal Spoonbill |
| | <i>Ardeicola plataleae</i> (Linnaeus, 1758) | |
| | <i>Eucolpocephalum femorale</i> (Piaget, 1880) | |
| | <i>Ibidoecus plataleae</i> (Denny, 1842) | |

Order ANSERIFORMES
Family ANATIDAE

- | | | |
|-------|---|--------------------------|
| *103. | Cygnus olor (Gmelin, 1789) | Mute Swan |
| | <i>Anatoecus icterodes oloris</i> Zlotorzycka, 1970 | |
| | <i>Ornithobius bucephalus</i> (Giebel, 1874) | |
| *104. | Cygnus atratus (Latham, 1790) | Black Swan |
| | <i>Anatoecus dentatus magnicornutus</i> Zlotorzycka, 1970 | |
| | <i>Anatoecus icterodes oloris</i> Zlotorzycka, 1970 | |
| | <i>Anatoecus</i> sp. | |
| | <i>Ciconiphilus pectiniventris</i> (Harrison, 1916) | |
| | <i>Holomenopon leucoxanthum</i> (Burmeister, 1838) | |
| | <i>Ornithobius fuscus</i> Le Souëf, 1902 | |
| | <i>Trinoton nigrum</i> Le Souëf, 1902 | |
| *105. | Branta canadensis (Linnaeus, 1758) | Canada Goose |
| | <i>Ciconiphilus pectiniventris</i> (Harrison, 1916) | |
| | <i>Holomenopon leucoxanthum</i> (Burmeister, 1838) | |
| | <i>Ornithobius goniopleurus</i> Denny, 1842 | |
| | <i>Trinoton querquedulae</i> (Linnaeus, 1758) s. l. | |
| 107. | Tadorna variegata (Gmelin, 1789) | Paradise Shelduck |
| | <i>Anaticola magnificus</i> Ansari, 1955 | |
| | <i>Anatoecus dentatus</i> (Scopoli, 1763) s. l. | |
| | <i>Anatoecus icterodes</i> (Nitzsch, 1818) s. l. | |
| | <i>Holomenopon tadornae</i> (Gervais, 1844) | |
| | <i>Trinoton querquedulae</i> (Linnaeus, 1758) s. l. | |
| *108. | Anas platyrhynchos platyrhynchos Linnaeus, 1758 | Mallard |
| | <i>Anaticola crassicornis</i> (Scopoli, 1763) | |
| | <i>Anatoecus dentatus</i> (Scopoli, 1763) | |
| | <i>Anatoecus icterodes</i> (Nitzsch, 1818) s. l. | |
| | <i>Holomenopon leucoxanthum</i> (Burmeister, 1838) | |
| | <i>Trinoton querquedulae</i> (Linnaeus, 1758) s. l. | |

- | | | |
|--------|---|-----------------------------|
| 109. | Anas superciliosa superciliosa Gmelin, 1789
<i>Anaticola crassicornis</i> (Scopoli, 1763) s. l.
<i>Anatoecus dentatus</i> (Scopoli, 1763) s. l.
<i>Anatoecus icterodes</i> (Nitzsch, 1818) s. l.
<i>Trinoton querquedulae</i> (Linnaeus, 1758) s. l. | Grey Duck |
| 110. | Anas gibberifrons gracilis Buller, 1869
<i>Anaticola crassicornis</i> (Scopoli, 1763) s. l.
<i>Anatoecus dentatus</i> (Scopoli, 1763) s. l.
<i>Anatoecus icterodes</i> (Nitzsch, 1818) s. l.
<i>Trinoton querquedulae</i> (Linnaeus, 1758) s. l. | Grey Teal |
| 111.a. | Anas aucklandica chlorotis Gray, 1845
<i>Anaticola</i> sp.
<i>Anatoecus icterodes</i> (Nitzsch, 1818) s. l. | Brown Teal |
| 111.b. | Anas aucklandica aucklandica (Gray, 1844)
<i>Holomenopon leucoxanthum</i> (Burmeister, 1838) | Auckland Island Teal |
| 113. | Anas rhynchos variegata (Gould, 1856)
<i>Anaticola crassicornis</i> (Scopoli, 1763) s. l.
<i>Anatoecus icterodes</i> (Nitzsch, 1818) s. l.
<i>Holomenopon</i> sp.
<i>Trinoton querquedulae</i> (Linnaeus, 1758) s. l. | New Zealand Shoveler |
| 114. | Hymenolaimus malacorhynchos (Gmelin, 1789)
<i>Acidoproctus gottwaldhirschi</i> Eichler, 1958
<i>Anatoecus icterodes</i> (Nitzsch, 1818) s. l. | Blue Duck |
| 116. | Aythya novaeseelandiae (Gmelin, 1789)
<i>Anaticola crassicornis</i> (Scopoli, 1763) s. l.
<i>Anatoecus icterodes</i> (Nitzsch, 1818) s. l.
<i>Holomenopon</i> sp. [Note 19]
<i>Trinoton querquedulae</i> (Linnaeus, 1758) s. l. | New Zealand Scaup |

Order FALCONIFORMES
Family ACCIPITRIDAE

- 119.a. **Circus approximans gouldi** Bonaparte, 1850 **Australasian Harrier**
Colpocephalum turbinatum Denny, 1842
Degeeriella fusca (Denny, 1842)
Nosopon lucidum (Rudow, 1869)

Family FALCONIDAE

120. **Falco novaeseelandiae** Gmelin, 1788 **New Zealand Falcon**
Degeeriella rufa rufa (Burmeister, 1838)
Nosopon sp.

121. **Falco cenchroides cenchroides** Vigors & Horsfield, 1827 **Nankeen Kestrel**
Colpocephalum zerafae Ansari, 1955
Degeeriella rufa rufa (Burmeister, 1838)
Laemobothrion (*Laemobothrion*) *tinnunculi* (Linnaeus, 1758)

Order **GALLIFORMES**
Family PHASIANIDAE

*122	Alectoris chukar (J. E. Gray, 1830) <i>Cuclogaster heterographus</i> (Nitzsch, 1866) <i>Goniocotes pusillus</i> (Nitzsch, 1866) <i>Goniodes dispar</i> Burmeister, 1838	Chukor
*123	Perdix perdix perdix (Linnaeus, 1758) <i>Amrysidea perdicis</i> (Denny, 1842) <i>Lipeurus maculosus maculosus</i> Clay, 1938	Grey Partridge
*125.	Synoicus ypsilophorus (Bosc, 1792) <i>Cuclogaster synoicus</i> (Clay, 1938) <i>Goniodes retractus</i> Le Souëf, 1902	Brown Quail
*126.	Colinus virginianus (Linnaeus, 1758) <i>Goniodes</i> sp. <i>Oxylipeurus clavatus</i> (McGregor, 1917)	Bobwhite Quail
*127	Lophortyx californica brunnescens Ridgway, 1884 <i>Goniodes stefani</i> Clay & Hopkins, 1955 <i>Lagopoecus docophoroides</i> (Piaget, 1880) <i>Oxylipeurus ellipticus</i> (Kéler, 1958)	California Quail
*128	Phasianus colchicus Linnaeus, 1758 <i>Amrysidea perdicis</i> (Denny, 1842) <i>Goniocotes chrysocephalus</i> Giebel, 1874 <i>Goniodes colchici</i> Denny, 1842 <i>Lipeurus maculosus maculosus</i> Clay, 1938 <i>Menacanthus</i> sp. <i>Oxylipeurus mesopelios colchicus</i> Clay, 1938	Pheasant
*129	Pavo cristatus Linnaeus, 1758 <i>Amrysidea minuta</i> Emerson, 1961 <i>Colpocephalum tausi</i> (Ansari, 1951) <i>Goniodes pavonis</i> (Linnaeus, 1778)	Peafowl

Family MELEAGRIDIDAE

*	Meleagris gallopavo Linnaeus, 1758 <i>Chelopistes meleagridis</i> (Linnaeus, 1758) <i>Menacanthus stramineus</i> (Nitzsch, 1818) <i>Oxylipeurus polytrapezius polytrapezius</i> (Burmeister, 1838)	Turkey
---	--	--------

Order **GRUIFORMES**
Family RALLIDAE

131.a.	Rallus philippensis assimilis Gray, 1843 <i>Pseudomenopon scopulacorne</i> (Denny, 1842) <i>Rallicola ortygometrae</i> (Schrank, 1781) s. l.	Banded Rail
134.a.	Gallirallus australis greyi (Buller, 1888) <i>Pseudomenopon pilgrimi</i> Price, 1974 <i>Rallicola harrisoni</i> Emerson, 1955	North Island Weka

- 134.b. **Gallirallus australis australis** (Sparrman, 1786) **Western Weka**
Pseudomenopon piligrimi Price, 1974
Rallicola harrisoni Emerson, 1955
- 134.c. **Gallirallus australis hectori** (Hutton, 1873) **Buff Weka**
Pseudomenopon piligrimi Price, 1974
Rallicola harrisoni Emerson, 1955
- 134.d. **Gallirallus australis scotti** (Ogilvie-Grant, 1905) **Stewart Island Weka**
Rallicola harrisoni Emerson, 1955 *s. l.*
135. **Porzana pusilla affinis** (Gray, 1846) **Marsh Crake**
Fulicoffula sp.
Incidifrons sp.
Rallicola sp.
136. **Porzana tabuensis plumbea** (Gray, 1829) **Spotless Crake**
Fulicoffula sp.
Rallicola tabuensis Emerson, 1966
137. **Porphyrio porphyrio melanotus** Temminck, 1820 **Pukeko**
Pseudomenopon concretum (Piaget, 1880)
Rallicola lugens (Giebel, 1874)
138. **Notornis mantelli** Owen, 1848 **Takahe**
Rallicola takahe Holloway, 1956
141. **Fulica atra australis** Gould, 1845 **Australasian Coot**
Fulicoffula lurida (Nitzsch, 1818)
Incidifrons fulicae (Linnaeus, 1758)
Pseudomenopon pilosum (Scopoli, 1763)
Rallicola fulicae (Denny, 1842)

Order CHARADRIIFORMES
Family HAEMATOPODIDAE

142. **Haematopus ostralegus finschi** Martens, 1897 **South Island Pied Oystercatcher**
Actornithophilus grandiceps (Piaget, 1880)
Quadraceps auratus (Haan, 1829)
Saemundssonia haematopi (Linnaeus, 1758)
143. **Haematopus unicolor** Forster, 1844 **Variable Oystercatcher**
Actornithophilus grandiceps (Piaget, 1880)
Quadraceps auratus (Haan, 1829)
Quadraceps ridgwayi (Kellogg, 1906)
Saemundssonia haematopi (Linnaeus, 1758)
144. **Haematopus chathamensis** Hartert, 1927 **Chatham Island Oystercatcher**
Actornithophilus grandiceps (Piaget, 1880)
Quadraceps auratus (Haan, 1829)

Family CHARADRIIDAE

145. ***Vanellus miles novaehollandiae*** Stephens, 1819 **Spur-winged Plover**
Actornithophilus hoplopteri (Mjöberg, 1910) *s. l.*
Austromenopon aegialitidis (Durrant, 1906) *s. l.*
Quadraceps renschi Timmermann, 1954
147. ***Pluvialis fulva*** (Gmelin, 1789) **Least Golden Plover**
Actornithophilus ochraceus (Nitzsch, 1818) *s. l.* [Note 20]
Quadraceps charadrii (Linnaeus, 1758) [Note 21]
Saemundssonia conica conica (Denny, 1842)
148. ***Charadrius obscurus*** (Gmelin, 1789) **New Zealand Dotterel**
Actornithophilus ochraceus (Nitzsch, 1818) *s. l.*
Quadraceps dominella Timmermann, 1953
- 150.a. ***Charadrius bicinctus bicinctus*** Jardine & Selby, 1827 **Banded Dotterel**
Quadraceps cedemajori Timmermann, 1969
Saemundssonia platygaster (Denny, 1842) *s. l.*
- 150.b. ***Charadrius bicinctus exilis*** Falla, 1978 **Auckland Island Banded Dotterel**
Quadraceps cedemajori Timmermann, 1969
Quadraceps novaeelandiae Timmermann, 1953
155. ***Thinornis novaeelandiae*** (Gmelin, 1789) **New Zealand Shore Plover**
Quadraceps novaeelandiae Timmermann, 1953
Saemundssonia chathamensis Timmermann, 1977
156. ***Anarhynchus frontalis*** Quoy & Gaimard, 1830 **Wrybill**
Quadraceps cedemajori Timmermann, 1969

Family SCOLOPACIDAE

157. ***Numenius madagascariensis*** (Linnaeus, 1766) **Far-eastern Curlew**
Lunaceps numenii numenii (Denny, 1842)
- 158.a. ***Numenius phaeopus variegatus*** (Scopoli, 1786) **Asiatic Whimbrel**
Austromenopon phaeopodis (Schrank, 1802)
Lunaceps numenii oliveri (Johnston & Harrison, 1912) [Note 22]
Saemundssonia scolopaciphaeopodis scolopaciphaeopodis (Schrank, 1803)
161. ***Limosa limosa melanurooides*** Gould, 1846 **Asiatic Black-tailed Godwit**
Actornithophilus spinulosus (Piaget, 1880)
Austromenopon limosae Timmermann, 1954
Carduiceps cingulatus (Denny, 1842)
Lunaceps limosella Timmermann, 1954
Saemundssonia thompsoni Timmermann, 1951

Family RECURVIROSTRIDAE

183. **Himantopus himantopus leucocephalus** Gould, 1837 **Pied Stilt**
Austromenopon himantopi Timmermann, 1954
Quadraceps hemichrous (Nitzsch, 1866)
Quadraceps semifissus (Nitzsch, 1866)
Saemundssonia platygaster (Denny, 1842) *s. l.*

184. **Himantopus novaezealandiae** Gould, 1841 **Black Stilt**
Quadraceps hemichrous (Nitzsch, 1866)

Family GLAREOLIDAE

188. **Glareola maldivarum** J. R. Forster, 1795 **Oriental Pratincole**
Actornithophilus sp.
Austromenopon sp.
Quadraceps ellipticus (Nitzsch, 1866) s. l.

Family STERCORARIIDAE

189. **Stercorarius skua lonnbergi** (Mathews, 1912) Southern Great Skua
Austromenopon fuscofasciatum (Piaget, 1880)
Haffneria grandis (Piaget, 1880) s. l.
Saemundssonia stresemanni Timmermann, 1949

189.1 **Stercorarius maccormicki** Saunders, 1893 Antarctic Skua
Austromenopon fuscofasciatum (Piaget, 1880)
Haffneria grandis (Piaget, 1880) s. l.
Quadraceps alpha (Kellogg, 1914)
Saemundssonia stresemanni Timmermann, 1949

191. **Stercorarius parasiticus** (Linnaeus, 1758) Arctic Skua
Austromenopon fuscofasciatum (Piaget, 1880)
Haffneria grandis (Piaget, 1880) s. l.
Quadraceps normifer (Grube, 1851)
Saemundssonia cephalus (Denny, 1842)

Family LARIDAE

- | | | |
|------|--|----------------------------|
| 193. | Larus dominicanus Lichtenstein, 1823 | Southern Black-backed Gull |
| | <i>Actornithophilus piceus</i> (Denny, 1842) s. l. | |
| | <i>Austromenopon transversum</i> (Denny, 1842) | |
| | <i>Quadraceps ornatus fuscolaminulatus</i> (Enderlein, 1908) | |
| | <i>Quadraceps punctatus</i> (Burmeister, 1838) s. l. | |
| | <i>Saemundssonia lari</i> (O. Fabricius, 1780) s. l. | |
| 194. | Larus novaehollandiae scopulinus Forster, 1844 | Red-billed Gull |
| | <i>Actornithophilus piceus</i> (Denny, 1842) s. l. | |
| | <i>Austromenopon transversum</i> (Denny, 1842) | |
| | <i>Quadraceps punctatus</i> (Burmeister, 1838) s. l. | |
| | <i>Saemundssonia lari</i> (O. Fabricius, 1780) s. l. | |
| 195. | Larus bulleri Hutton, 1871 | Black-billed Gull |
| | <i>Austromenopon transversum</i> (Denny, 1842) | |
| | <i>Quadraceps punctatus</i> (Burmeister, 1838) s. l. | |
| | <i>Saemundssonia lari</i> (O. Fabricius, 1780) s. l. | |

Family STERNIDAE

199. **Hydroprogne caspia** (Pallas, 1770) Caspian Tern
Austromenopon atrofulvum (Piaget, 1880) s: l.
Quadraceps caspius (Giebel, 1874)

199.1 **Sterna albostriata** (Gray, 1845) Black-fronted Tern
Quadraceps sp.

201. **Sterna vittata bethunei** Buller, 1896 [Note 24] **Antarctic Tern**
Austromenopon atrofulvum (Piaget, 1880) s. l.
Quadraceps sellatus (Burmeister, 1838) s. l.
Saemundssonia lockleyi Clay, 1949 s. l.
202. **Sterna paradisaea** Pontoppidan, 1763 **Arctic Tern**
Quadraceps houri Hopkins, 1949
Saemundssonia lockleyi Clay, 1949 s. l.
204. **Sterna albifrons sinensis** Gmelin, 1789 **Eastern Little Tern**
Quadraceps nychthererus (Burmeister, 1838)
Saemundssonia melanocephalus (Burmeister, 1838)
205. **Sterna striata** (Gmelin, 1789) **White-fronted Tern**
Austromenopon atrofulvum (Piaget, 1880) s. l.
Quadraceps sellatus (Burmeister, 1838) s. l.
Saemundssonia sterna (Linnaeus, 1758)
206. **Sterna fuscata** (Linnaeus, 1766) **Sooty Tern**
Austromenopon atrofulvum (Piaget, 1880) s. l.
Quadraceps birostris (Giebel, 1874)
Saemundssonia albemarlensis (Kellogg & Kuwana, 1902)
208. **Anous minutus minutus** Boie, 1844 **White-capped Noddy**
Actornithophilus ceruleus (Timmermann, 1954) s. l.
Austromenopon atrofulvum (Piaget, 1880) s. l.
Quadraceps hopkinsi hopkinsi Timmermann, 1952
Saemundssonia lobaticeps remota Timmermann, 1951
210. **Procelsterna cerulea albivitta** Bonaparte, 1856 **Grey Ternlet**
Actornithophilus ceruleus (Timmermann, 1954)
Austromenopon atrofulvum (Piaget, 1880) s. l.
Quadraceps hopkinsi apophoretus Timmermann, 1969

Order COLUMBIFORMES
Family COLUMBIIDAE

- 211.a. **Hemiphaga novaeseelandiae novaeseelandiae** (Gmelin, 1789) **New Zealand Pigeon**
Hohorstiella sp.
Patellinirmus novaeseelandiae Tendeiro, 1972
- 211.b. **Hemiphaga novaeseelandiae chathamensis** (Rothschild, 1891) **Chatham Island Pigeon**
Patellinirmus harrisoni Tendeiro, 1972
- *212. **Columba livia** Gmelin, 1789 **Rock Pigeon**
Bonomiella columbae Emerson, 1957
Campanulotes bidentatus compar (Burmeister, 1838)
Colpocephalum turbinatum Denny, 1842
Columbicola columbae columbae (Linnaeus, 1758)
Hohorstiella lata (Piaget, 1880)

Order PSITTACIFORMES
Family CACATUIDAE

214. **Strigops habroptilus** Gray, 1845 Kakapo
Heteromenopon (Keamenopon) kea (Kellogg, 1907)
- Family NESTORIDAE
- 216.a. **Nestor meridionalis septentrionalis** Lorenz, 1896 North Island Kaka
Forficuloeucus meinertzhageni Guimarães, 1974
Heteromenopon (Keamenopon) kea (Kellogg, 1907)
Neopsittaconirmus kea (Kellogg, 1907)
- 216.b. **Nestor meridionalis meridionalis** (Gmelin, 1788) South Island Kaka
Forficuloeucus meinertzhageni Guimarães, 1974
Heteromenopon (Keamenopon) kea (Kellogg, 1907)
Neopsittaconirmus kea (Kellogg, 1907)
217. **Nestor notabilis** Gould, 1856 Kea
Colpocephalum pilgrimi Price, 1967
Forficuloeucus meinertzhageni Guimarães, 1974 s. l.
Heteromenopon (Keamenopon) kea (Kellogg, 1907)
Neopsittaconirmus kea (Kellogg, 1907)
- Family PLATYCERCIDAE
220. **Cyanoramphus unicolor** (Lear, 1831) Antipodes Island Parakeet
Forficuloeucus sp.
Heteromenopon (Keamenopon) sp.
- 221.b. **Cyanoramphus novaezelandiae novaezelandiae** Sparrman, 1787 Red-crowned Parakeet
Forficuloeucus sp.
- 221.c. **Cyanoramphus novaezelandiae chathamensis** Oliver, 1930 Chatham Island Red-crowned Parakeet
Forficuloeucus sp.
- 221.d. **Cyanoramphus novaezelandiae hochstetteri** (Reischek, 1889) Reischek's Parakeet
Forficuloeucus sp.
- 222.a. **Cyanoramphus auriceps auriceps** (Kuhl, 1820) **Yellow-crowned Parakeet**
Forficuloeucus sp.
- 222.b. **Cyanoramphus auriceps forbesi** Rothschild, 1893 Chatham Island Yellow-crowned Parakeet
Forficuloeucus sp.
223. **Cyanoramphus malherbi** (Souance, 1857) Orange-fronted Parakeet
Forficuloeucus sp.

Order CUCULIFORMES
Family CUCULIDAE

227. **Chrysococcyx lucidus lucidus** (Gmelin, 1788) **Shining Cuckoo**
Cuculicola kui Kettle, 1980

Order STRIGIFORMES
Family STRIGIDAE

230. **Tyto alba delicatula** (Gould, 1837) **Australian Barn Owl**
Strigiphilus aitkeni Clay, 1966
231. **Ninox novaeseelandiae novaeseelandiae** (Gmelin, 1788) **Morepork**
Kurodaia cryptostigmatica (Nitzsch, 1861)
Strigiphilus vapidus Clay, 1977
- 232.b. **Sceloglaux albifacies albifacies** (Gray, 1844) **Laughing Owl**
Strigiphilus sp.
- *233. **Athene noctua** (Scopoli, 1769) **Little Owl**
Strigiphilus cursitans (Nitzsch, 1861)

Order CORACIIFORMES
Family ALCEDINIDAE

236. **Halcyon sancta vagans** (Lesson, 1830) **New Zealand Kingfisher**
Alcedoecus alatoclypeatus (Piaget, 1885) s. l.

Family CORACIIDAE
238. **Eurystomus orientalis pacificus** (Latham, 1801) **Broad-billed Roller**
Capriella sp.

Order PASSERIFORMES
Family XENICIDAE

- 239.b. **Acanthisitta chloris chloris** (Sparrman, 1787) **South Island Rifleman**
Menacanthus sp.
-
- Family ALAUDIDAE

- *243. **Alauda arvensis arvensis** Linnaeus, 1758 **Skylark**
Brueelia sp.
Philopterus sp.

Family MOTACILLIDAE

- 246.a. **Anthus novaeseelandiae novaeseelandiae** (Gmelin, 1789) **New Zealand Pipit**
Philopterus sp.
- 246.c. **Anthus novaeseelandiae steindachneri** Reischek, 1889 **Antipodes Pipit**
Philopterus sp.

Family PRUNELLIDAE

- *249. **Prunella modularis occidentalis** (Hartert, 1910) **Hedgesparrow**
Philopterus sp.

Family MUSCICAPIDAE

252. **Mohoua albicilla** (Lesson, 1830) **Whitehead**
Myrsidea sp.
Philopterus sp.
254. **Gerygone igata** (Quoy & Gaimard, 1830) **Grey Warbler**
Myrsidea sp.
Philopterus sp.
Ricinus sp.
- 256.a. **Rhipidura fuliginosa placabilis** Bangs, 1921 **North Island Fantail**
Philopterus sp.
- 256.b. **Rhipidura fuliginosa fuliginosa** (Sparrman, 1787) **South Island Fantail**
Menacanthus sp.
Philopterus sp.
- 257.a. **Petroica macrocephala toitoi** (Lesson, 1828) **Pied Tit**
Philopterus sp.
- 257.b. **Petroica macrocephala macrocephala** (Gmelin, 1789) **Yellow-breasted Tit**
Philopterus sp.
- 257.d. **Petroica macrocephala dannefaerdi** (Rothschild, 1894) **Black Tit**
Philopterus sp.
- 258.b. **Petroica (Miro) australis australis** (Sparrman, 1788) **South Island Robin**
Menacanthus eurysternus (Burmeister, 1838)
- *261. **Turdus philomelos clarkei** Hartert, 1909 **Song Thrush**
Brueelia sp.
Menacanthus eurysternus (Burmeister, 1838)
Philopterus turdi (Denny, 1842)
- *262. **Turdus merula merula** Linnaeus, 1758 **Blackbird**
Brueelia merulensis (Denny, 1842)
Brueelia sp.
Menacanthus eurysternus (Burmeister, 1838)
Myrsidea thoracica (Giebel, 1874)
Philopterus turdi (Denny, 1842)

Family ZOSTEROPIDAE

263. **Zosterops lateralis lateralis** (Latham, 1801) **Silvereye**
Menacanthus eurysternus (Burmeister, 1838)

Family MELIPHAGIDAE

264. **Notiomystis cincta** (Du Bus, 1839) **Stitchbird**
Myrsidea sp.
- 265.a. **Anthornis melanura obscura** Falla, 1948 **Three Kings Bellbird**
Brueelia sp.
Myrsidea sp.
- 265.b. **Anthornis melanura melanura** (Sparrman, 1786) **Bellbird**
Brueelia sp.
Menacanthus eurysternus (Burmeister, 1838)
Myrsidea sp.
266. **Prosthemadera novaeseelandiae novaeseelandiae** (Gmelin, 1788) [Note 25] **Tui**
Brueelia sp.
Menacanthus eurysternus (Burmeister, 1838)
Myrsidea sp.

Family EMBERIZIDAE

- *268. **Emberiza citrinella caliginosa** Clancey, 1940 **Yellowhammer**
Brueelia delicata (Nitzsch, 1866)
Myrsidea sp.

Family FRINGILLIDAE

- *271. **Carduelis chloris chloris** (Linnaeus, 1758) **Greenfinch**
Brueelia sp.
Myrsidea sp.
- *272. **Carduelis carduelis britannica** (Hartert, 1903) **Goldfinch**
Brueelia sp.

Family PLOCEIDAE

- *274. **Passer domesticus domesticus** (Linnaeus, 1758) **House Sparrow**
Brueelia cyclothorax (Burmeister, 1838)
Menacanthus eurysternus (Burmeister, 1838)

Family STURNIDAE

- *275. **Sturnus vulgaris vulgaris** Linnaeus, 1758 **Starling**
Brueelia nebulosa (Burmeister, 1838)
Menacanthus eurysternus (Burmeister, 1838)
Sturnidoecus sturni (Schrank, 1776)
- *276. **Acridotheres tristis** (Linnaeus, 1766) **Indian Myna**
Menacanthus eurysternus (Burmeister, 1838)
Sturnidoecus sp.

Family CALLAEIDAE

- 277.a. **Philesturnus carunculatus rufusater** (Lesson, 1828) **North Island Saddleback**
Brueelia sp.
- 277.b. **Philesturnus carunculatus carunculatus** (Gmelin, 1789) **South Island Saddleback**
Brueelia sp.
278. **Heteralocha acutirostris** (Gould, 1837) **Huia**
Rallicola sp.
- 279.a. **Callaeas cinerea wilsoni** (Bonaparte, 1851) **North Island Kokako**
Brueelia sp.
Philopterus sp.
- 279.b. **Callaeas cinerea cinerea** (Gmelin, 1788) **South Island Kokako**
Brueelia sp.
Philopterus sp.

Family CRACTICIDAE

- *280.a. **Gymnorhina tibicen tibicen** (Latham, 1801) **Black-backed Magpie**
Brueelia semiannulata (Piaget, 1883)
Myrsidea sp.
Philopterus sp.
- *280.b. **Gymnorhina tibicen hypoleuca** (Gould, 1837) **White-backed Magpie**
Brueelia semiannulata (Piaget, 1883)
Myrsidea sp.

Family CORVIDAE

- *282. **Corvus frugilegus frugilegus** Linnaeus, 1758 **Rook**
Colpocephalum fregili Denny, 1842

EXPLANATORY NOTES

1. Tandan (1972:68) recorded several specimens of *Apterygon mirum* Clay, 1961 and two males as *Apterygon* sp. *incertae sedis* from his host, suggesting a wrong host record as one of the possible explanations for an apparently abnormal situation. Evidence now available confirms Tandan's suggestion that the host for the above two males was misidentified. It is *Apteryx australis australis*.
2. Clay (1972:73) recorded 42 specimens of *Rallicola gadowi* Harrison, 1915 from this host. Re-examination of the skins from which those lice were collected shows that the hosts were misidentified; they are *Apteryx australis australis*. Further samples of *Rallicola* (*Aptericola*) from *Apteryx australis mantelli* do not include *R. (A.) gadowi*.
3. Clay (1967:154) recorded *Austrogoniodes concii* from this host in "New Zealand — Christchurch", but this record appears to be based on specimens taken from a yearling female, New Brighton, March 1941. The type host of *A. concii* is *Eudyptes chrysocome moseleyi*, type locality Tristan da Cunha. Lice from nine specimens of *Eudyptes chrysocome chrysocome* from the New Zealand subregion have been examined without finding any *A. concii*. Hence, we hold Clay's record in suspense pending confirmation.
4. Clay (1967:154) recorded *Austrogoniodes hamiltoni* Harrison, 1937 and *Austrogoniodes macquariensis* Harrison, 1937 from this host. Our many subsequent samples of *Austrogoniodes* from *E. pachyrhynchus* do not include these two species. We have reason to believe that there is a misinterpretation of the host identity in Clay (1967).
5. Watson (1967:71) recorded *Austrogoniodes hamiltoni* Harrison, 1937 and *Austrogoniodes macquariensis* Harrison, 1937 from this host (then named *Eudyptes pachyrhynchus atratus*). However, we have been unable to find these lice species from this bird again, despite the large number of *Austrogoniodes* collected from The Snares (Horning, Palma & Pilgrim, 1980:8). We regard Watson's records as stragglers on this host.
6. Harrison (1937:15) recorded one female *Austrogoniodes waterstoni* (Cummings, 1914) from this host. We regard this record as a straggler on this host as, in our experience, *A. waterstoni* is confined to *Eudyptula minor* subspecies.
7. We have not seen specimens of this louse species from this host, but we include the record following Clay (1967:154).
8. Wise (1977:62) recorded *Naubates fuliginosus* (Taschenberg, 1882) following Clay & Moreby (1970:217), who listed it from this host (as *Diomedea exulans*) at Macquarie I. As species of the genus *Naubates* are not regular parasites on members of the family Diomedeidae, we regard the above record as straggler.
9. Wise (1977:63) recorded *Perineus concinnus* (Kellogg & Chapman, 1899) following Harrison (1937:29). Pilgrim & Palma (1979:177) showed that it was a misidentification of *Perineus circumfasciatus* Kéler, 1957.
10. Bedford (1930:170) described *Naubates pterodromi* from three females taken from *Pterodroma macroptera*. As *Naubates* females from several species of *Pterodroma* are indistinguishable, males are needed for a correct identification.

We have found that males from this host fit Harrison's description of *N. heteroproctus* [see Notes 11 and 12].

11. Harrison (1937:30) described *Naubates heteroproctus* from this host, but we have reason to believe that this is an erroneous host record. Subsequent material from *Pterodroma lessonii* is not *N. heteroproctus* [see Notes 10 and 12].
12. *Naubates* males from this host agree with the male described as *N. pterodromi* in Kéler (1952:213) from *Pterodroma incerta*. It is possible that Kéler's specimens were not conspecific with *N. pterodromi* Bedford, 1930 because males from *Pt. macroptera* (the type host of *N. pterodromi*) are different from males taken from *Pt. incerta*. However, males from *Pt. macroptera* agree with the description of *N. heteroproctus* Harrison, 1937 [see Note 10], and Kéler's use of the name *N. pterodromi* has been followed by several authors. Therefore, considering that females of these *Naubates* species are not diagnostic and that a stable nomenclature is desirable, we use *N. pterodromi* Bedford, 1930 in the sense of Kéler, 1952.
13. Watt (1971:238, 242) recorded four nymphs of *Saemundssonia* sp. from this host. Re-examination of the material shows that it is a misidentification of *Trabeculus* sp.
14. Watt (1971:236, 242) recorded many specimens of *Halipeurus leucophryna* Timmermann, 1960 from this host. Re-examination of the material shows that it is not *H. leucophryna*.
15. We have not seen specimens from this host collected in the New Zealand subregion, but we include three lice species following Watson (1967:70, 72). However, we do not accept two further records of Watson (1967:72, 74) which we regard as stragglers, i.e. *Halipeurus turtur* Edwards, 1961 and *Trabeculus* sp. The latter was published in Harrison (1937: 37) as *Giebelia hexakon* Waterston, 1914.
16. Wise (1977:60) recorded *Halipeurus (Halipeurus) angusticeps* (Piaget, 1880) following Harrison (1937:31), who examined one female from this host. Re-examination of Harrison's original material shows that it is a misidentification of *Halipeurus (Halipeurus) procellariae* (J. C. Fabricius, 1775) which we regard as a straggler on this host.
17. Kéler (1952:209) listed this bird as one of the recorded hosts for *Ancistrona vagelli* (J. C. Fabricius, 1787) from the Kermadec Is. He presumably meant the closely related Kermadec Storm Petrel, which was formerly numerous on the Kermadec Is (OSNZ, 1970:29). See also entry 69b and Note 18.
18. We have not seen specimens of this louse genus from this host. We include this record following Johnston & Harrison (1912:364) and Watt (1971:233, 243).
19. Wise (1977:57) recorded *Holomenopon clypeilargum* Eichler, 1943 following Price (1971:643), who examined one female from this host. Examination of more specimens, including males, shows that they are not *H. clypeilargum*.
20. Wise (1977:56) recorded *Actornithophilus timidus* (Kellogg, 1896) following Watt (1971:233, 243). We regard this species as a junior synonym of *A. ochraceus*, following Clay (1962:243).

21. Wise (1977:64) recorded *Quadraceps orarius* (Kellogg, 1896) following Johnston & Harrison (1912:368) and Hopkins & Clay (1952:314). We regard this species as a junior synonym of *Q. charadrii*, following J. M. Martens (pers. comm.).
22. Wise (1977:61) recorded this subspecies as *Lunaceps numenii phaeopi* (Denny, 1842) following Emerson (1972:95).
23. Wise (1977:65) recorded *Saemundssonia limosae* (Denny, 1842) following Harrison (1937:21-22, 46), who examined one male from this host. Watson (1967:74) referred to apparently the same material as *Saemundssonia* sp. We have not been able to examine the specimen reported by Harrison.
24. Wise (1977:65) recorded *S. melanocephalus* (Burmeister, 1838) following Harrison (1937:22, 46), who examined males and females from this host (*Sterna vittata macquariensis* = *S. v. bethunei*). Watson (1967:74) referred to apparently the same material as *Saemundssonia* sp. We have not been able to examine any of the specimens reported by Harrison.
25. Watt (1971:238, 244) recorded four nymphs of *Sturnidoecus* sp. from this host. Re-examination of the material shows that it is a misidentification of *Saemundssonia* sp. and probably a contaminant.

Acknowledgements

We wish to record our indebtedness for numerous identifications and for assistance provided by Dr Theresa Clay, London; Professor K. C. Emerson, Sanibel I., Florida; Professor L. Guimarães, São Paulo; Dr P. Kettle, Wallaceville, Upper Hutt; Dr H. Klockenhoff, Bonn; Mr C. H. C. Lyal, London; Dr J. Martens, Hamburg; Mr C. Moreby, London; Dr B. C. Nelson, Berkeley, California; Dr R. D. Price, St Paul, Minnesota; Professor J. Tendeiro, Lisbon; Dr D. W. Tuff, San Marcos, Texas and Dr R. A. Ward, Washington, D. C.

We are grateful to the many collectors who have sent us much material on which to base this work. One of us (RLCP) received a grant, C/62, from the University of Canterbury to provide some equipment used in the study.

REFERENCES

- BEDFORD, G.A.H. 1930. New genera and species of Mallophaga from South African hosts. 16th Report of the Director of Veterinary Services and Animal Industry, Union of South Africa: 158-173.
- CLAY, T. 1953. Revisions of the genera of Mallophaga. I. The *Rallicola*-complex. Proc. Zool. Soc. London 123(3):563-587.
- CLAY, T. 1962. A key to the species of *Actornithophilus* Ferris with notes and descriptions of new species. Bull. Brit. Mus. (Nat. Hist.), Entomology 11(5):191-244, plates 4-11.
- CLAY, T. 1964. Insects of Campbell Island. Phthiraptera. Pacific Insects Monograph 7:230-234.
- CLAY, T. 1967. Mallophaga (Bitting lice) and Anoplura (Sucking lice). Part I: *Austrogonioides* (Mallophaga) parasitic on penguins (Sphenisciformes). Antarctic Research Series, Entomology of Antarctica 10: 149-155. J. L. Gressit (Ed.) Washington, D.C.
- CLAY, T. 1971a. A new species of *Austrogonioides* (Phthiraptera: Philopteridae) from a duck (Anseriformes). J. Aust. Ent. Soc. 10(4): 293-298.
- CLAY, T. 1971b. A new genus and two new species of Boopidae (Phthiraptera: Amblycera). Pacific Insects 13(3-4):519-529.
- CLAY, T. 1972. The species of *Rallicola* (Insecta: Phthiraptera: Ischnocera) parasitic on kiwis (*Apteryx*). NZ J. Sci. 15(1):70-76.
- CLAY, T.; MOREBY, C. 1970. Mallophaga and Anoplura of Subantarctic islands. Pacific Insects Monograph 23:216-220.
- EMERSON, K. C. 1972. Checklist of the Mallophaga of North America (North of Mexico). Part I. Suborder Ischnocera. Deseret Test Center, Dugway, Utah, 200 pp.

- HARRISON, L. 1987. Mallophaga and Siphunculata. Australasian Antarctic Expedition 1911-14. Sci. Rep., Ser. C. — Zoology and Botany 2(1):1-47, 3 plates.
- HOPKINS, G. H. E.; CLAY, T. 1952. A checklist of the genera & species of *Mallophaga*. Trustees of the British Museum, London, 362 pp.
- HORNING, D. S.; PALMA, R. L. & PILGRIM, R. L. C. 1980. The lice (Insecta: Phthiraptera) from the Snares Islands, New Zealand. Nat. Mus. NZ Misc. Ser. No. 3:1-17.
- JOHNSTON, T. H.; HARRISON, L. 1912. On a collection of Mallophaga from the Kermadecs. Trans. NZ Inst. 44:363-373.
- KELER, S. von. 1952. On some Mallophaga of sea-birds from the Tristan Da Cunha Group and The Dyer Island. J. Ent. Soc. Southern Africa 15(2): 204-238.
- MARTENS, J. M.; PALMA, R. L. 1981. Species distribution of genus *Quadraceps* (Mallophaga: Philopteridae) on New Zealand endemic plovers. NZ J. Zool. 8(1):83-85.
- ORNITHOLOGICAL SOCIETY OF NEW ZEALAND. 1970. Annotated checklist of the birds of New Zealand, including the birds of the Ross Dependency. The Checklist Committee (F. C. Kinsky, Convener). A. H. & A. W. Reed, Wellington, 96 pp., 4 maps.
- ORNITHOLOGICAL SOCIETY OF NEW ZEALAND. 1980. Amendments and additions to the 1970 annotated checklist of the birds of New Zealand. The Checklist Committee (F. C. Kinsky, Convener). Notornis (Supplement) 27:1-23.
- PILGRIM, R. L. C. 1980. The New Zealand flea fauna. Pp. 173-184 in R. Traub & H. Starcke, (Eds), Fleas. Balkema, Rotterdam.
- PILGRIM, R. L. C.; PALMA, R. L. 1979. A redescription of *Perineus concinnus* (Mallophaga: Philopteridae). Pacific Insects 21(2-3):172-178.
- PRICE, R. D. 1971. A review of the genus *Holomenopon* (Mallophaga: Menoponidae) from the Anseriformes. Annals Ent. Soc. America 64(3):633-646.
- TANDAN, B. K. 1972. The species of *Apterygon* (Insecta: Phthiraptera: Amblycera) parasitic on kiwis (*Apteryx*). NZ J. Sci. 15(1):52-69.
- WATSON, K. C. 1967. The terrestrial arthropoda of Macquarie Island. III. Parasites of mammals and birds. ANARE Sci. Rep., Ser. B (I) Zoology, No. 99: 70-76.
- WATT, J. C. 1971. Ectoparasitic insects on birds and mammals of the Kermadec Islands. Notornis 18(4):227-244.
- WISE, K. A. J. 1977. A synonymic checklist of the Hexapoda of the New Zealand Sub-region. The smaller orders. Bull. Auck. Inst. Mus. 11:1-176.

APPENDIX 1

Suggestions for collecting bird lice

With dead birds, it is preferable to send whole carcasses, each enclosed in a separate plastic bag, together with adequate collecting data, to National Museum, Wellington. If this is inconvenient, or if the carcase is unsuitable for postal or freight transport, feathers sent in a plastic bag are a good substitute.

With live birds, spray a pet insecticide (e. g. "Nuvan" aerosol) among the bird's feathers, and hold the bird in a plastic bag for a few minutes. Send the sample so obtained, as it is, in the bag, together with all the dust and insecticide that have come off the bird. Inside each bag, put a label with the name of the bird, date, locality, and name of the collector. Seal the bag with adhesive tape.