


NOTORNIS



QUARTERLY BULLETIN
of the
Ornithological Society of New Zealand



Volume Nine, Number Six, September, 1961

NOTORNIS

In continuation of New Zealand Bird Notes

BULLETIN OF THE ORNITHOLOGICAL SOCIETY OF NEW ZEALAND
(Incorporated)

Registered with the G.P.O., Wellington, as a Magazine

Edited by R. B. SIBSON, King's College, Auckland S.E.7

Annual Subscription: Ordinary Member, £1; Endowment Member, 30/-
Life Membership, £20 (for members over thirty years of age)

OFFICERS 1960-61

President — Mr. A. BLACKBURN, 10 Score Road, Gisborne
North Island Vice-President — Dr. R. A. FALLA, Dominion Museum,
Wellington

South Island Vice-President — Mrs. L. E. WALKER, 15 Cornwall Street,
Vauxhall, Dunedin

Editor — Mr. R. B. SIBSON, King's College, Auckland, S.E.7

Assistant Editor — Mr. A. BLACKBURN, 10 Score Road, Gisborne

Treasurer — Mr. H. R. MCKENZIE, North Road, Clevedon

Secretary — Mr. G. R. WILLIAMS, Zoology Department,
Lincoln College, Christchurch, Canterbury

Members of Council:

Mr. E. G. TURBOTT, Canterbury Museum, Christchurch

Mr. J. C. DAVENPORT, 2 Lanark Road, Tokoroa

Mr. F. C. KINSKY, Dominion Museum, Wellington

Banding Convener — Mr. F. C. KINSKY, Dominion Museum, Wellington

Organiser of Nest Records Scheme — Mr. J. C. R. CLARIDGE,
82 Mungavin Avenue, Porirua

Contents of Vol. 9, No. 6: September, 1961

Preliminary Analysis of Records of "Storm-Killed" Sea Birds from New Zealand, 1939-59	185
Short Notes — South Island Robin Eating Venison; Blue Reef Heron at Lake Taupo; Silvereyes Moving at Night Over Christ- church; Bellbirds Attacking Sparrows; Harrier Attacked by Spur-winged Plovers; Greenfinches and Silvereyes Stripping Cotoneaster; Grey-backed Storm Petrel and Small Passerines at the Brothers Lighthouse; Unusual Death of Morepork; Kiwis in Bay of Plenty	199
Plate XXXVII — Pair of Rock Wrens at Nest-Near Homer Tunnel	201
Plate XXXVIII — Red-fronted Parakeet at Nest on Little Barrier	202
Plate XXXIX — Whitehead on Little Barrier	203
Plate XL — Banded Rail	204
Plate XLI — Young White-faced Heron	205
Plate XLII — Bittern in Full Threat Display	206
Plate XLIII — Close-up of Bittern in Full Threat Display	207
Plate XLIV — Westland Black Petrel Near its Burrow	208
Reprints in Library	211

NOTORNIS

VOLUME NINE, NUMBER SIX - SEPTEMBER, NINETEEN SIXTY-ONE

PRELIMINARY ANALYSIS OF RECORDS OF " STORM-KILLED " SEA BIRDS FROM NEW ZEALAND, 1939-59

Compiled by P. C. BULL* and B. W. BOESON

*Animal Ecology Division, Department of Scientific and Industrial Research, Lower Hutt

I ABSTRACT

Published and unpublished records held by the Ornithological Society of New Zealand give the identity and the date and place of collection of 6,960 specimens of oceanic birds that have been found dead on New Zealand beaches during the period 1939 to 1959. Of the 50 species represented, 38 belong to the order Procellariiformes. Prions (*Pachyptila* spp.) constitute 64 per cent. of the total records and Shearwaters (*Puffinus* spp.) 20 per cent. The records of most of the species, but particularly of the prions, are characterised by occasional periods of very heavy mortality ("wrecks") which sometimes involve a single species, sometimes several species. Records of the Fairy Prion (*Pachyptila turtur*) are presented in some detail to show distribution and seasonal movements; comparable data for other species are given in summary. Many of the species breed in the South Indian Ocean, and banding of birds at breeding stations there and elsewhere in the Subantarctic can be expected to yield many interesting recoveries from New Zealand.

II INTRODUCTION

The recording of sea birds found dead on New Zealand beaches has been one of the New Zealand Ornithological Society's activities almost from its inception. Several such records appear in the Society's first Annual Report (1939-40), and many others are included in the "Summarised Classified Notes" sections of subsequent publications. These records provide a useful source of information for anyone studying the distribution and seasonal movements of birds living off the New Zealand coast. A defect of the records, however, is that, although people usually report the finding of rare species, they sometimes neglect to record the commoner ones, and consequently the records may not give a true picture of the relative abundance of the different species. Further, most of the records give no indication of how much beach was examined, and, without this information, the finding of many birds could mean either that many were present or that the observer covered an unusually long stretch of coastline.

To correct these defects in the records, a special Beach Patrol Scheme was initiated in October 1951 (Notornis 4: 142), and members were asked to record on special cards *all* the sea birds they found, together with details of the date, locality, length of beach patrolled, freshness of specimens and weather conditions during preceding days. The Scheme flourished for a few years and then gradually lapsed. A considerable number of cards were sent in by members, but no summaries were published. The continued inclusion of records of dead sea birds in the Summarised Classified Notes caused some confusion in that

records were sometimes published in the notes, sometimes merely entered on the patrol cards and sometimes preserved both on the cards and in the notes.

The Beach Patrol Scheme was revived in 1960 (Notornis 8: 268) and has met with a good response from members. It was recently decided to discontinue the publication in the "Classified Notes" of records of dead sea birds, and instead to publish an annual summary of the cards sent in; it is hoped that the first of these summaries, that for the year 1960, will appear shortly.

The present compilation deals only with records made prior to 1960. It attempts to describe the nature of the material so far collected, to point out some of the defects in it, and to illustrate some of the ways in which the data can be used to obtain information on species that are rarely seen except at breeding colonies or from ships at sea where specific identification is often impossible.

III THE RECORDS EXAMINED

A. Beach Patrol Cards

The total number of cards held, for patrols made prior to 1960, is 344, but this number may be increased because several observers have promised to make out cards for patrols at present recorded in private field notebooks. Although 32 observers contributed to the above total, only nine provided more than ten cards each and only three sent in more than 40 cards each.

The monthly and yearly distribution of patrols, irrespective of locality, is shown in Table 1 below.

Table 1: Number of Beach Patrols each Month, 1951-59

	—1951	1951	1952	1953	1954	1955	1956	1957	1958	1959	Total
Jan.	4	—	6	3	4	2	1	—	—	2	22
Feb.	4	—	4	6	7	—	1	—	1	2	25
Mar.	—	—	6	11	9	—	—	—	1	2	29
Apr.	1	—	7	21	8	1	1	1	1	6	47
May	—	6	5	14	1	1	—	2	5	6	40
June	—	—	6	5	3	—	—	2	3	8	27
July	—	—	4	11	9	2	—	—	5	8	39
Aug.	1	2	7	7	2	1	—	—	2	6	28
Sept.	—	1	2	—	2	—	—	3	5	5	18
Oct.	—	5	6	6	2	1	—	3	2	1	26
Nov.	1	7	4	8	—	—	1	1	4	4	30
Dec.	1	5	—	2	1	—	—	2	2	—	13
Total	12	26	57	94	48	8	4	14	31	50	344

The number of cards sent in has varied markedly from year to year. Following the initiation of the Beach Patrol Scheme in 1951, the number of patrols increased progressively to reach a peak of 94 in 1953, and then declined to the ignominious figure of four in 1956. In subsequent years the figures rose again to a modest peak of 50 in 1959, but this latter increase was largely due to the contributions of a very few observers working in the Wellington district. Twelve cards record details of patrols made before the scheme was started; these cards, presumably made from entries in field notebooks, cover the years 1943 to 1949. The cards are representative of all seasons of the year;

the combined patrols, over the whole period, exceed 20 for every month except September (18) and December (13); one or more patrols were made in nearly every month of the years 1952, 1953, 1954, 1958 and 1959.




Fig. 1 — Distribution of Beach Patrols. Each dot indicates a locality in which one or more patrols have been made. For purposes of analysis the localities have been grouped in zones; these are indicated as follows:

- A-B Auckland West Coast, Hokianga to Kawhia
- C-D Auckland East Coast, Bay of Islands to East Cape
- E-F Wellington West Coast, Wanganui to Cape Terawhiti
- F-G Wellington South Coast, Cape Terawhiti to Cape Palliser
- H-I North Canterbury, Motunau to Banks Peninsula

The geographic distribution of the patrols (Fig. 1) is very uneven. Over 90 per cent. of the patrols relate to beaches in the vicinity of Auckland or Wellington, and there are no patrols at all from several long stretches of coastline; for example, there are none from the whole length of the west coast of the South Island (Dr. R. A. Falla advises that dead birds are rare on beaches near Hokitika and Okarito). For purposes of analysis, the Auckland patrols have been divided into Western and Eastern zones, and the Wellington ones into Western and Southern zones (Fig. 1). A few patrols from the Christchurch-Motunau Island area have been placed in a fifth zone and all others grouped together as "miscellaneous." In most of these zones several patrols were made each year from 1951 to 1954, but only in the Wellington West Coast Zone, which has the largest number of patrols, is there any approach to an unbroken record for the full period 1951 to 1959 (Table 2).

Table 2: Number of Beach Patrols on Various Coastlines, 1951-59

	—1951	1951	1952	1953	1954	1955	1956	1957	1958	1959	Total
Auck. East	-	7	14	32	10	-	-	-	1	1	65
Auck. West	-	7	23	14	13	-	-	-	-	-	57
Wgtn. West	7	1	6	28	14	5	3	13	29	35	141
Wgtn. Sth.	5	2	3	13	7	3	1	1	1	14	50
Nth. Canty.	-	4	5	2	-	-	-	-	-	-	11
Miscell.	-	5	6	5	4	-	-	-	-	-	20
Total	12	26	57	94	48	8	4	14	31	50	344

The length of beach patrolled is indicated on only 266 of the cards, and the total distance covered by these patrols is 647 miles. The contribution made by each zone to this total is shown in Table 3, together with the average number of petrels found per mile in each zone.

Table 3: Patrols for which Mileage was Recorded

Zone	No. Patrols	Total No. Miles Covered	Total No. Petrels Found	Birds Per Mile
Auck. East	33	94	132	1.4
Auck. West	48	195	1195	6.1
Wgtn. West	121	230	644	2.8
Wgtn. Sth.	44	63	69	1.1
Nth. Canty.	10	44	52	1.2
Miscell.	10	21	12	0.6
Total	266	647	2104	3.3

The number of birds found per mile in the Auckland West Coast Zone is more than twice that of any other zone. This may indicate that more birds are washed up in this zone than in others, or, less likely, that, in this zone, a higher proportion of the patrols took place after strong gales than elsewhere.

The identification of some of the petrels is a matter of considerable difficulty, and the specimen record cards vary with regard

to evidence of correct identification. Some cards have detailed measurements or reference to a preserved skin or skeleton, while others have been signed by an experienced ornithologist who has examined the original specimens. Some cards, however, provide no indication of the grounds on which the identification was made. In view of these considerations, anyone using the Society's records will need to inspect the cards and published records himself. Since the present work is intended as a compilation rather than a critical study, all the records have been accepted at face value; subspecific determinations, however, have been omitted.

B. Records from "Notornis"

For the present purpose, "Notornis" is taken to include also the Society's earlier publication, "N.Z. Bird Notes." The material covers a period of 20 years (1939-59) and also includes a few records from earlier years. The published notes thus provide some indication of sea bird mortality during the 12 years immediately preceding the introduction of the Beach Patrol Scheme, and, even in later years, they include many birds not recorded on beach patrol cards. For the present purpose, all the published records of each species were listed. The list for each species was then compared with the beach patrol cards and any obvious duplications were deleted from the list of published records, together with records of birds found dead at breeding colonies or of healthy birds caught for banding. In this way the two sets of material were made reasonably comparable and duplication was reduced to a minimum.

The published records usually give no indication of how much beach was examined, nor of the basis on which the specimens were identified. In some instances it is even difficult to decide whether a record refers to a living bird or to a dead one.

IV PRELIMINARY ANALYSES

A. Records of *Pachyptila turtur*

Records of Fairy Prion (*P. turtur*), a common and fairly easily recognised species, are presented below in some detail as an example of the kind of data that are available for many other species.

In Table 4 all the Society's records of *P. turtur* have been grouped according to the month and year in which the birds were collected. Nearly all the records of birds found prior to 1951 are from "Notornis" whereas most of the later records are from beach patrol cards. An entry of "0" in the table signifies that no *P. turtur* was found even though some beach was examined that month as indicated by a beach patrol card or a published record concerning the finding of other petrels; in the absence of such indication the space is left blank. The table gives some measure of observer effort during the 20-year period and may be useful in helping to decide whether the lack of records of other species is to be interpreted as a true scarcity of birds or merely an absence of observers.

It is apparent that the observations provide a much better coverage of the second half of the 20-year period than of the first half. Observations are lacking for 48 per cent. of the months in the first ten years but for only seven per cent. of those in the next decade.

Table 4: Records of *Pachyptila turtur*

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Total
-1940							12						12
1940	0	1		1	1			0					3
1941	0			1	2				2				5
1942		0			0	0	0			0		0	-
1943	35	15	4		0		1			3			58
1944					0		0		0				-
1945		0			0		0				0		-
1946	0	0	0	0			1	49	4	0		0	54
1947		2		0	0	0			0			0	2
1948	4	0		0	4	4	4	2	3	1	9	6	37
1949	2	2			4	0	17		2		35	4	66
1950		0	1	0	0	0	2	146	0		0	0	149
1951	8	0	0	1	9	0	0	5	2	52	26	9	112
1952	0	0	3	1	1	2	0	0	1	8	5	1	22
1953	95	11	32	7	3	6	31	8	0	12	26	3	234
1954	1	3	1	5	0	12	128	15	62	3	4	5	239
1955	6	11	1	0	1	0	13	4	18	3		2	59
1956	9	41		0	1		14	3	2	0	0	0	70
1957		0	0	0	0	2			4	3	2	3	14
1958	0	0	0	1	3	6	4	12	13	2	44	0	85
1959	0	0	2	0	2	3	1	2	17	0	6	0	33
Total	160	86	44	17	31	35	228	246	130	87	157	33	1254

It is also likely that patrols were fewer or shorter in the first decade than in the second because the percentage of months in which no *P. turtur* was found (even though some beach was examined) was 50 per cent. during the period 1940-49, but only 32 per cent. during 1950-59. The same conclusion may be drawn from the fact that the average number of *P. turtur* found per month (excluding months with no observations) rose from 3.6 in 1940-49 to 9.1 in 1950-59.

The most striking feature of Table 4 is that the number of birds reported varies greatly from month to month and from year to year. For instance, of the 160 birds found in January (left hand column), 95 (59%) were recorded in January 1953, and a further 35 (22%) in January 1943; birds found in January of the remaining 18 years totalled only 30 (19% of the 20-year total). The occasional periods of heavy mortality may occur at different times of the year. For

instance more than 30 *P. turtur* were found in January (1943 and 1953), February (1956), March (1953), July (1953 and 1954), August (1946 and 1950), September (1954), October (1951) and November (1949 and 1958).

It is assumed above that these high figures indicate unusually severe mortality, but it could be argued that they are perhaps due merely to increased activity of observers. This possibility can be checked by reference to beach patrol cards that show the length of beach examined in the combined Auckland and Wellington West Coast Zones (Tables 5 and 6). Table 5 shows the average number of *P. turtur* found per mile of beach in each year from 1951 to 1959. Excluding the years 1955 and 1956 when only a few miles of beach were examined, the values vary from 2.5 birds per mile in 1951 to 0.2 birds per mile in 1952. At least some beach was examined in every month of these two years (Table 4) so it is unlikely that the low value for 1952 could be explained by inadequate patrolling; indeed, on the basis of miles covered, 1952 was a year of rather high patrolling activity (Table 5). The number of birds found per mile on certain

Table 5: Mean Number of *P. turtur* Found per Mile in Different Years
(Auckland and Wellington West Coasts combined)

	— 1951	1951	1952	1953	1954	1955	1956	1957	1958	1959	Total
No. Miles Patrolled	4	24	90	104	49	6	2	16	72	58	425
No. <i>P. turtur</i> found	9	60	16	195	56	4	41	10	72	30	493
Birds per Mile	2.3	2.5	0.2	1.9	1.1	(0.7)	(20.5)	0.6	1.0	0.5	1.2

occasions was sometimes much higher than the average values given in Tables 5 and 6. For instance, 31 *P. turtur* were found during a patrol of one mile on the Auckland West Coast in October 1951, and 38 were found over a distance of two miles on the Wellington West Coast in February 1956. It is apparent therefore that there are certain occasions when mortality is particularly severe; such occasions are sometimes referred to as "wrecks." Although this term is convenient for describing periods of unusually severe mortality, there is no evidence to suggest that wrecks differ, except in degree, from the more frequent occasions when the birds occur in only moderate numbers.

The variation in the monthly averages (Table 6) is even more marked than in the yearly ones and ranges from 0.2 birds per mile in April to 6.3 in January.

Table 6: Mean Number of *P. turtur* Found per Mile in Different Months
(Auckland and Wellington West Coasts combined)

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Total
No. Miles Patrolled	16	14	50	56	35	51	62	20	35	25	43	19	425
No. <i>P. turtur</i> found	100	50	37	10	13	17	51	13	36	53	103	10	493
Birds per Mile	6.3	3.7	0.7	0.2	0.4	0.3	0.8	0.7	1.0	2.1	2.4	0.5	1.2

The three lowest values (all under 0.5 birds per mile) occur in April, May and June. Further, during the whole period 1940 to 1959, the highest number of *P. turtur* found in any of the above three months was only 12 (June, 1954); except for December (only nine birds) the other months all have a maximum figure of at least 30 birds (Table 4). These results may indicate that the mortality of *P. turtur* tends to be lower in April, May and June than during the rest of the year or that the species is scarce in offshore waters at that season. Any such conclusion, however, must be regarded as provisional because the figures would be completely changed if a wreck should occur in the April-June period, and there are as yet too few records of wrecks to give any assurance that they are really rarer at this season than at others.

Although the species *P. turtur* is present off the New Zealand coast throughout the year, a detailed study of the measurements of specimens found at different seasons might show that there is in fact considerable seasonal movement — the disappearance of birds of one race being balanced by the arrival of those of another.

The geographic distribution of records of *P. turtur* is almost the same as that of beach patrols (Fig. 1). Of the 1,254 birds listed in Table 4, 52 per cent. are from the Auckland West Coast, 36 per cent. from the Wellington West Coast, five per cent. from North Canterbury, 4 per cent. from the Auckland East Coast-Bay of Plenty zone and only 3 per cent. from other areas. The species has been found along almost the whole length of the West Coast of the North Island; on the East Coast specimens have been reported from North Auckland, Firth of Thames, Bay of Plenty and Gisborne. In the South Island *P. turtur* has been found at Cape Farewell, New Brighton (Christchurch), Lake Ellesmere and Dunedin. Probably the species is to be found on all New Zealand coasts where tide, currents and shoreline conditions are such that birds can be washed ashore. Fairy Prions seem to be washed up in greater numbers on West Coast Beaches than on Eastern or Southern ones, but the number of patrols from the latter beaches is still too small to prove this. The highest mortalities recorded for other than West Coast beaches are 0.4 birds per mile in the Bay of Plenty in July, 1953, 1.0 per mile on the Wellington South Coast in April, 1953, and 1.3 per mile in Canterbury in October, 1951. The published records include a few birds from as far inland as Taihape and Queenstown.

B. Records of Other Species

Most of the records refer to species in the Order Procellariiformes (albatrosses, petrels and shearwaters, storm petrels and diving petrels) and these are listed in Table 7. In addition, there are 182 records of penguins; *Eudyptula minor* (150 records) is by far the most numerous species, but there are also a few records of *Megadyptes antipodes*, *Eudyptula albosignata* and *Eudyptes sclateri*. Other groups represented total only 14 specimens and comprise the following species: *Phaethon rubricauda* (1), *Sula leucogaster* (1), *Fregata ariel* (2), *Stercorarius skua* (5), *S. pomarinus* (1), *S. parasiticus* (2), *Sterna bergii* (1), and *S. fuscata* (1). Records of common New Zealand species of shags, gannets, gulls and terns, together with a few records of land birds, have been excluded from this compilation because in these species better information is usually available from the study of living birds. The paragraphs that follow deal only with the petrels and their allies.

Table 7: Summary of Records of Petrels and Their Allies

SPECIES	Total No. Records	% from "Notornis"	No. Records from each of zones				
			CD	AB	EF	FG	HI
<i>Diomedea exulans</i>	42	90	1	19	1	8	5
<i>D. epomophora</i>	19	95	—	10	2	6	1
<i>D. melanophris</i>	16	81	2	12	—	—	2
<i>D. chrysostoma</i>	130	61	—	104	23	3	—
<i>D. bulleri</i>	11	73	1	5	1	4	—
<i>D. cauta</i>	122	78	2	33	15	63	7
<i>Phoebastria palpebrata</i>	15	73	2	9	1	1	1
<i>Macronectes giganteus</i>	78	62	2	54	8	2	3
<i>Daption capensis</i>	66	68	5	31	18	7	3
<i>Fulmarus glacialisoides</i>	7	(86)	—	3	2	—	1
<i>Halobaena caerulea</i>	58	84	1	51	5	—	—
<i>Pachyptila vittata</i>	1026	81	51	596	211	2	107
<i>P. salvini</i>	1720	68	6	1611	101	1	—
<i>P. desolata</i>	163	64	15	116	16	2	14
<i>P. belcheri</i>	146	62	3	116	26	—	—
<i>P. turtur</i>	1254	52	54	650	457	18	56
<i>Puffinus carneipes</i>	60	63	23	32	2	1	—
<i>P. bulleri</i>	140	71	24	57	19	31	6
<i>P. griseus</i>	806	64	40	610	77	28	33
<i>P. tenuirostris</i>	102	84	20	61	17	2	1
<i>P. gavia</i>	234	53	41	115	49	6	17
<i>P. assimilis</i>	20	80	5	9	5	—	—
<i>Procellaria cinerea</i>	4	(75)	—	3	1	—	—
<i>P. parkinsoni</i>	5	(100)	2	—	—	1	—
<i>P. westlandica</i>	2	(100)	—	1	1	—	—
<i>P. aequinoctialis</i>	12	58	1	9	2	—	—
<i>Pterodroma macroptera</i>	52	60	23	13	—	11	2
<i>P. lessoni</i>	154	71	2	126	18	2	2
<i>P. inexpectata</i>	33	67	2	24	5	1	1
<i>P. brevirostris</i>	22	73	1	12	8	—	1
<i>P. pycrofti</i>	6	(100)	6	—	—	—	—
<i>P. leucoptera</i>	1	(100)	—	1	—	—	—
<i>P. cooki</i>	17	65	9	7	1	—	—
<i>P. nigripennis</i>	1	(100)	—	—	1	—	—
<i>Oceanites oceanicus</i>	1	(0)	1	—	—	—	—
<i>Garrodia nereis</i>	6	(100)	—	—	1	—	2
<i>Pelagodroma marina</i>	22	64	14	4	1	1	2
<i>Pelecanoides urinatrix</i>	191	49	95	45	35	6	7
TOTALS	6764	66	454	4549	1130	207	274

Brackets indicate percentages based on less than 10 records

Zones: CD — Auckland East Coast-Bay of Plenty (see Fig. 1)

AB — Auckland West Coast

FG — Wellington South Coast

EF — Wellington West Coast

HI — North Canterbury Coast

The prions (*Genus Pachyptila*) are by far the most numerous group and comprise 64 per cent. of the total records. Of the other groups, the shearwaters make up 20 per cent., other petrels seven per cent., albatrosses five per cent., diving petrels three per cent. and storm petrels less than one per cent. The Lesser Broad-billed Prion (*P. salvini*) is the most abundant species with 1,720 specimens. There are 1,254 records of the Fairy Prion (*P. turtur*), 1,026 of the Broad-billed Prion (*P. vittata*) and 806 of the Sooty Shearwater (*P. griseus*); except for the Fluttering Shearwater (*P. gavia*) (234 records), the other species are each represented by less than 200 records.

Although there are more records of *P. salvini* than of any other species, 83 per cent. of the specimens were found during June and July of 1954. The number of records of *P. turtur*, the next most abundant species, is considerably below that of *P. salvini*, but the records are more evenly distributed (Table 4) and the greatest number of specimens recorded in any period of two consecutive months constitutes only 12 per cent. of the total records. It is apparent therefore that *P. turtur* is more often found than *P. salvini*, even though the total number of specimens is greater for *P. salvini*. A comparable, but less obvious difference, may occur between some of the shearwaters. Taking the maximum number of birds found in any period of two consecutive months as a percentage of the total records, it was found that the highest value for *P. griseus* was 26 per cent., but for *P. gavia* only 16 per cent. The data concerning the diving petrel *P. urinatrix*, though not analysed in detail, appear to resemble those for *P. turtur* and *P. gavia* in consisting of many records of small numbers of specimens rather than of a few "wrecks" involving many birds.

People visiting beaches at times when there has been no recent heavy mortality may find the above three species more often than other species and thus tend to regard them as common and the record not worthy of publication. This possibility is examined in Table 7. The total number of records for each species is shown in the first column of figures; some of these records were from "Notornis" and the rest from beach patrol cards. The percentage of the records that came from "Notornis" (second column of figures) is much lower for *P. turtur* (52 per cent.), *P. gavia* (53 per cent.) and *P. urinatrix* (49 per cent.) than for any of the other species with comparable numbers of records (60 per cent to 84 per cent. for all species with more than 50 records). The fact that the beach patrol scheme rather than "Notornis" yielded so high a proportion of the total records for these three species is best explained on the hypothesis that people have not bothered to publish the finding of small numbers of these relatively common species. It is probable that "wrecks" of these species do occur (see Table 4 for *P. turtur*), but the constant occurrence of small numbers of dead birds between the "wrecks" means that the latter come to constitute a smaller proportion of the total records. *P. turtur*, *P. gavia* and *P. urinatrix* are present off the New Zealand coast throughout the year and they all breed in large numbers on islands in the vicinity of Auckland and Wellington so it is not surprising that they figure frequently in the beach patrols, most of which are from beaches near these centres (Table 3).

The wrecks of petrels sometimes involve mainly a single species and at other times several species. For instance, a wreck, consisting almost entirely of *P. vittata*, occurred in August, 1946. In July, 1954,

on the other hand, the records include 1,135 *P. salvini*, 128 *P. turtur* and over one hundred *P. vittata*. A southerly gale in February, 1947, resulted in the stranding in Palliser Bay over 40 Salvin's Mollymawks (*Diomedea cauta salvini*) and unusually large numbers of Buller's Shearwaters (*P. bulleri*) and Grey-faced Petrels, (*P. macroptera*). These wrecks are recorded in "Notornis" together with information on the weather conditions that preceded them (see Turbott and Sibson, 1946; Wodzicki, 1947; Cunningham, 1948; and Davenport and Sibson, 1955). Earlier wrecks, involving mostly of *P. salvini*, occurred in 1918 and 1932 (Falla, 1940).

Dr. R. A. Falla has pointed out that one might separate winter wrecks (in which food shortage is sometimes suspected as an underlying cause) from wrecks of young birds caught by strong winds within a few days of leaving their burrows. These very young birds can be distinguished by the nature of the bill and plumage; the dates on which they occur are sometimes useful in indicating the time at which young birds leave inaccessible nesting colonies (R. A. Falla, personal communication).

The present data are of limited value for showing the geographic distribution of the different petrels because most of the observations refer to beaches in the vicinity of Auckland and Wellington and there are few if any records from several long stretches of coastline (Fig. 1). A glance at the five right hand columns of Table 7 shows that, in general, the species present are substantially the same in all zones in which regular observations have been made. A notable exception is the Grey-faced Petrel (*P. macroptera*) which has not been found on the Wellington West Coast despite intensive patrolling in this area; the eleven specimens of *P. macroptera* from the Wellington South Coast were all found together after the gale of February, 1947 (Cunningham, 1947). Despite the similarity of the species lists, there are considerable differences between some of the zones with regard to the relative abundance of certain birds. For instance, prions constitute 68 per cent. of the total records from the Auckland West Coast, but only 28 per cent. of those from the East Coast. Conversely, Diving Petrels (*P. urinatrix*) are more abundant on the Auckland East Coast (21 per cent. of all records) than in any other zone (less than four per cent. of total records). Except for *P. macroptera*, the records for the Auckland and Wellington West Coasts are very similar, both with regard to the species present and to their numerical importance.

There are marked differences between the species with regard to the months in which most of the specimens are recorded. This is illustrated in Table 8 for a few species in each of several groups of petrels. Some species are found in good numbers during all months of the year (e.g. *P. gavia* and *P. turtur*), others are rare or absent in mid winter (Flesh-footed and Buller's Shearwaters, *P. carneipes* and *P. bulleri*) and others again occur mainly during the winter months (*P. salvini* and the Blue Petrel, *H. caerulea*). These differences of course are related to the seasonal movements of the species and would be clearer still were it not that many of the records provide no indication of how long the specimen had been lying on the beach before it was found. The occurrence of "wrecks" also increases the difficulty of interpreting Table 8, since the high average for some months is due to the single occurrence of large numbers of birds; as more records accumulate, wrecks may be found in other months and perhaps modify the seasonal patterns shown in the table.

Table 8: Seasonal Distribution of Records

Species	Number Specimens Recorded in Each Month												Total Records
	J	F	M	A	M	J	J	A	S	O	N	D	
<i>Diomedea chrysostoma</i>	3	1	6	3	5	8	66	12	17	3	3	3	130
<i>D. cauta</i>	7	57	8	8	10	6	11	6	2	2	4	1	122
<i>Macronectes giganteus</i>	7	1	1	3	3	8	20	17	10	5	2	2	78
<i>Daption capensis</i>	2	1	-	-	2	4	17	21	6	6	6	1	66
<i>Halobaena caerulea</i>	-	1	1	5	16	1	18	9	9	7	-	-	58
<i>Pachyptila vittata</i>	151	1	2	-	22	76	290	410	31	11	30	3	1026
<i>P. salvini</i>	-	-	1	-	3	18	83	22	11	2	6	-	146
<i>P. belcheri</i>	160	-	1	17	31	35	228	246	130	87	157	33	1254
<i>P. turitur</i>	17	11	10	6	7	2	5	-	-	-	1	6	60
<i>Puffinus carneipes</i>	7	33	4	12	29	7	5	-	-	5	22	16	140
<i>P. bulleri</i>	43	27	21	28	125	23	13	4	1	11	364	146	806
<i>P. griseus</i>	10	14	24	33	14	8	21	48	14	18	22	8	234
<i>Pterodroma macroptera</i>	6	11	3	1	6	6	2	2	3	-	4	6	52
<i>P. lessoni</i>	7	2	7	8	11	15	32	34	9	14	7	8	154
<i>Pelecanoides urinatrix</i>	35	5	4	6	5	2	14	38	11	7	21	43	191

For some species, such as *P. salvini*, the nearest known breeding places are on islands in the south Indian Ocean. In view of the many specimens washed up on New Zealand beaches, a banding programme on these and other subantarctic islands should provide some interesting recoveries from New Zealand. The species found in largest numbers on New Zealand beaches, and therefore most likely to yield recoveries, are indicated in Table 7.

V DISCUSSION

The collection, preservation, analysis and publication of the data listed in Table 7 have cost many people a great deal of time and also a considerable amount of money. The main purpose of this compilation is to summarise what has been collected so far and thus provide a basis for considering any changes that may be required in the Scheme.

It is tempting to advocate measures that would ensure a better coverage of beaches in all parts of the country, but the consequent increases in clerical work would be more than the Society could hope to handle on the present voluntary basis. Before considering alternative methods of improving the Scheme it may be useful to summarise its main objectives. In the opinion of the compilers, these objectives are:

- (1) To add to the list of species recorded from New Zealand waters and to ensure that specimens of particular scientific interest are sent to Museums.
- (2) To collect information on the distribution and seasonal movements of birds living off the New Zealand coast.
- (3) To record the extent and frequency of wrecks, and the conditions that accompany them, so as to provide a better understanding of their cause.
- (4) To increase the chances of recovery of banded birds which would give unequivocal information as to the places of origin of New Zealand off-shore birds.
- (5) To provide data which will allow the linking of sea bird distribution with the results of current studies on the distribution of plankton, the surface-living animals on which the birds feed.

It can be argued, and with justification, that the number of dead birds on the beach at a given time is not a valid indication of the abundance of birds offshore. If the records cover a long enough period, however, it may be possible to detect certain well marked trends that can be interpreted in terms of distribution and movements. For instance if a species such as *P. macroptera* is found regularly in the northern part of the country, but rarely if at all in the south, it seems reasonable to conclude that the normal range of this species is in the more northern waters. With the Cape Pigeon, records from the more northerly part of the country mostly occur in winter, and this can be interpreted as indicating a northerly extension of the range in winter provided records from further south show that mortality does in fact occur during summer. Again, if *P. turtur* is recorded throughout the year, but *P. salvini* only in winter, it seems safe to conclude that the latter species is absent from New Zealand waters during the summer.

In years of heavy *P. griseus* mortality, records of the distribution of dead birds, together with a record of weather conditions, provide some indication of routes followed by the birds while migrating along the coast. In all these instances, however, reliable conclusions can be drawn only if the same pattern emerges year after year. The value of the records is thus proportional to the length of the period they cover and to their continuity throughout this period.

In view of the above and the limits inherent in any purely voluntary scheme, it is considered that the objectives of the Beach Patrol Scheme can best be attained by giving first priority to the maintenance of regular patrols on the Auckland and Wellington West Coast beaches. Birds are washed up in large numbers on these beaches, the existing records already cover a period of 20 years, the beaches are close to large cities where there are a number of observers who can co-operate to ensure continuity of patrols, and finally the results to date are sufficiently similar in the two areas to justify the use of data from one area to cover a temporary break in the continuity of records from the other. In due course it may be possible to link the records from these areas with observations made from ships passing along the coast, and thus obtain a further check on the limitations of the beach patrol data.

The collection of records on distribution in New Zealand as a whole could be achieved by having emergency observers in as many districts as possible. These observers would agree to inspect their local beaches when advised that heavy mortality was occurring in the study areas, or when local weather conditions indicated the likelihood of seabird mortality. Species new to the New Zealand list, and specimens of rare species, can occur anywhere in New Zealand, but are to be expected particularly in the far north and in the far south, and it is important to find observers to cover these areas. The above suggestions for the more efficient use of available resources are in no way intended to discourage those who prefer to do casual patrols outside the Scheme's main programme; the results of such patrols will always be welcome.

In conclusion, it must be emphasised that the success of the whole scheme depends on the correct identification of the birds found. People taking part should remember that, until detected, a single faulty identification may obscure the significance of several accurate records made by other people. It should also be stressed that negative records (no birds found) are valuable and that all cards should include information concerning the number of miles patrolled, the freshness of all specimens found and weather conditions that may have contributed to the presence or absence of dead birds.

VI ACKNOWLEDGEMENTS

The data presented in this compilation have been collected by so many people that all cannot be acknowledged individually. The names of those responsible for the many published notes are already recorded in the appropriate numbers of "*Notornis*"; the beach patrol cards summarised here were sent in by D. H. Brathwaite, J. Britton, P. C. Bull, C. N. Challies, R. Colegrove, A. C. Couttes, D. E. Crockett, J. M. Cunningham, W. C. Clark, J. C. Davenport, R. K. Dell, M. C. Downes, Dr. R. A. Falla, Dr. C. A. Fleming, J. Frame, M. B. Gill, P. C. Harper, H. Hodgkins, Miss N. MacDonald, W. R. Marsden,

H. R. McKenzie, K. H. Miers, A. C. Prentice, N. T. Parham, S. C. Rutherford, H. Secker, R. B. Sibson, M. M. Taylor, E. G. Turbott, Mrs. L. E. Walker, H. Warburton and A. H. Watson.

The compilers are also grateful to Drs. R. A. Falla and K. Wodzicki and to Mr. R. H. Taylor, who very kindly read through the manuscript, and to Mr. F. J. Tindall (Soil Bureau, D.S.I.R.), who drew Fig. 1.

VII REFERENCES

- Cunningham, J. M., 1948: Seabird Mortality in February, 1947; *N.Z. Bird Notes* 2: 188-193.
- Davenport, J. C.; Sibson, R. B., 1955: Notes on two rare Petrels (*H. caerulea* and *Pt. brevirostris*); *Notornis* 6: 115-117.
- Falla, R. A., 1940: The Genus *Pachyptila* Illiger; *Emu* 40: 218-236.
- Turbott, E. G.; Sibson, R. B., 1946: Petrels cast ashore by August gales, 1946, with special reference to *Pachyptila vittata*; *N.Z. Bird Notes* 2: 19-23.
- Wodzicki, K., 1947: Destruction of Prions by storm, August, 1946. (With special reference to Wellington Province). *N.Z. Bird Notes* 2: 64-69.



SHORT NOTES

SOUTH ISLAND ROBIN EATING VENISON

The tameness of the Robins (*P. australis*) in the Eglinton Valley is proverbial, but I have never found them so tame as in May 1961. All eleven birds which I contacted at various parts of the valley (five of them in an area of about ten acres) readily accepted small pieces of bread and though they did not actually feed from my hand, most of them would quite happily take bread fragments dropped on the ground just by my foot. They were not at all interested in bits of biscuit. The pieces of bread were usually carried in the bill to a perch on branch or stump, and most birds returned for more — one bird had twelve pieces. Two birds on the roadside were particularly tame; both were I think cocks, one rather aggressive and the other rather cowed. The aggressive bird took its food to a perch, but the other preferred to hop under the car and eat its portions. After feeding these birds for a little time I left them and went into the bush for about half an hour; when I returned they were still there, the first bird on a perch, the second bird on the ground by the front wheel of the car, body held semi-erect but slightly crouched, wings slightly drooped and tail spread, with the tips of the feathers touching the ground; its whole attitude was expressive of supplication and expectancy.

Robins were watched taking insects and worms from the forest floor, and one bird was seen "marking time" and later collecting an insect from the moss where it had been treading. A deer had recently been killed, skinned and some of the flesh removed. In the process of cutting the meat some small fragments of flesh had been scattered on the ground around the dead beast. As I first came upon the dead deer I saw a Robin take and eat one of the meat fragments, and on my return half an hour later I watched the bird again hop up to the carcase and take another piece. I do not know if this carnivorous behaviour has been previously recorded.

— A. T. EDGAR

BLUE REEF HERON AT LAKE TAUPO

On 5/7/61, my wife and I stopped on the Taupo town waterfront to examine a party of Little Shags at the outer end of a groyne about forty yards from the lake shore. A Blue Heron (*Egretta sacra*) was seen trotting along the groyne towards the shags, catching a fish, probably a "Cockabully," on the way. From the road above the lake we used binoculars to make sure that it was indeed a Blue Heron. While visiting Major G. F. and Mrs. Yerex we learned that there had been a report of a Blue Heron but Major Yerex had not investigated it because White-faced Herons (*Ardea novaehollandiae*) frequent the lake and are usually reported as "Blue" or "Reef" Herons. He accompanied us to the lake shore where we now found two Blue Herons, apparently quite at home in their new-found "sea." Two were seen again by Mrs. McKenzie on 6th and by Dr. J. S. Armstrong on 12th. A single bird, or the two, were seen several times between these dates. Sometimes they were on the stones of the groynes, sometimes padding along the sandy beach.

— H. R. & H. M. MCKENZIE

[There are no recent inland records of the Reef Heron. However, Buller (2nd Edition, p. 131) reports "a pair was seen on the Taupo Lake in October 1875, and I observed one on the wing at Lake Rotoiti in October 1884." — Ed.]



SILVEREYES MOVING AT NIGHT OVER CHRISTCHURCH

On 11/4/61 at 9 p.m., I heard flocks of Silvereyes flying over my home, in Shirley, Christchurch. I did not pay much attention to them, but I did note that they seemed to be travelling roughly from east to west. It was quite dark at the time. The following evening I heard flocks, flying in the same direction, from 9 p.m. to 10.30 p.m., and they may have continued longer. A flock would be heard slowly coming up from the east and then would pass overhead and the flight calls would die away towards the west. Sounds carry well at night and so distances are hard to estimate, but they seemed to be flying fairly high — about the 100 ft. level. Judging from the frequency of calls, there must have been at least twenty birds in each flock, maybe many more, and they came over at a rate of about one flock every five minutes or less. This would mean about 400 birds flew over while I listened on 12/4/61. On both nights there was no wind so, if they were migrating, they were almost certainly not off course.

Stead (*Life Histories of N.Z. Birds*) states that there is a northward migration of Silvereyes in the Christchurch area but, as the Canterbury breeding population is low, there is no return movement in Spring. Oliver (*N.Z. Birds 1955*) accounts for the greater numbers in winter by a movement into settled areas in search of food. Fleming (*Emu*, 42, 209) found no evidence of a migration in Auckland, but suggests aggregation and local wandering. The easterly movement of the birds, over this area, does not agree either with Stead's northerly migration, or a dispersal in search of food, but movement at night does suggest some kind of migration.

— DAVID DAWSON



[M. F. Soper

XXXVII — Pair of Rock Wrens (**Xenicus gilviventris**) at nest near Homer Tunnel, described in Notornis IX, 158-161.



[G. J. H. Moon

XXXVIII — Red-fronted Parakeet (*C. novaezelandiae*) at nest on Little Barrier. Now scarce on the main islands, this parakeet remains plentiful on many offshore islands.



[G. J. H. Moon

XXXIX—The Whitehead (*Mohoua albicilla*) has a somewhat peculiar distribution in the North Island. It is very common on Little Barrier, where this photograph was taken.



[G. J. H. Moon

XL — The elusive Banded Rail (*Rallus philippensis*) is not uncommon along the mangrove creeks of northern New Zealand. The beauty of its markings is seldom noted.



[F. C. Kinsky

XLI — This young White-faced Heron (*Ardea novaehollandiae*) fell from a nest near Foxton in October, 1960, and was cared for at the Dominion Museum.

XLII — Dr. Soper's excellent photographs of a Bittern (*Botaurus poicilophyllus*) at the nest were published in *Notornis* VIII, 45 & 46. In this photograph the full throat display is shown.



IF. C. Kinsky



[F. C. Kinsky

XLIII — Close-up of a Bittern during threat display in a Southland garden, May 1961. Note the eyes. This bird had a slightly injured wing.



[Peter Morrison

XLIV — Westland Black Petrel *Procellaria westlandica*) near its burrow on a bushed ridge in Westland, as described in Notornis VII, 230-232.

BELLBIRD ATTACKING SPARROWS

On 20/4/61 a happy-sounding (male?) Bellbird (*A. melanura*) was heard in the tall gum-trees at the bottom of my garden in New Plymouth. Thence, he moved to an oak tree near the house. The sight of a dozen House Sparrows (*P. domesticus*) milling over a lump of bread at the feeding-table on the lawn seemed to put ideas of mischief into his head, for he made a 50ft. dive, worthy of any falcon or sparrowhawk, at the sparrows. No attempt was made to strike but the Bellbird passed a foot above them and rose into a copper beech nearby. The Sparrows also scattered into the same tree; where the Bellbird sorted out a victim and put it to flight around the lawn and trees at a speed of which I never thought either was capable. While the Sparrow took shelter in some kowhai trees, the Bellbird returned to the oak to await the Sparrows who were soon busy again at the feeding-table. Then down swept the Bellbird to repeat the assault. This went on some half a dozen times till he tired of the game.

— C. T. GIBSON

★

HARRIER ATTACKED BY SPUR-WINGED PLOVERS

Towards dusk on 30/6/61, I witnessed the following attack on a Harrier (*C. approximans*) by a party of five Spur-winged Plovers (*L. novaehollandiae*). This took place over paddocks immediately southwest of the Meteorological Office, Invercargill. My attention was drawn to the scene by the amount of noise the plovers were creating. The mode of attack appeared systematic and was observed as follows:—

One of the party would swoop and attack the Harrier from the rear. Completing the attack, it would rise to the level of the remaining birds; and as this occurred, another of the party would swoop in, again from the rear. This second bird having attacked, yet another would swoop in, and so on. The object of this mode of attack was seemingly to keep the Harrier in a vulnerable position near the ground. This they achieved, and when I lost sight of the birds against a line of conifers, the Harrier was still near the ground and the attack was continuing. By this time the birds were 800 yards away.

— E. J. LE HUQUET

★

GREENFINCHES AND SILVEREYES STRIPPING COTONEASTER

I was interested in A. T. Edgar's notes on Greenfinches (*C. chloris*) eating cotoneaster berries at Titirangi. For some years now a cotoneaster bush at Puketitiri, Hawkes Bay (2000 feet a.s.l.) has been regularly visited by numerous Greenfinches as well as Silvereyes (*Z. lateralis*) which feast off the berries.

It is quite common for up to fifteen Silvereyes and six or eight Greenfinches to be in the bush at once. They are usually present between April and May. They start with the topmost berries, furthest from the ground and work down the bush, completely stripping it of berries as they go. In fact, this last winter the bush was completely devoid of berries within a few weeks of their ripening. It is a large bush, too.

— PAM. M. LEWIS

GREY-BACKED STORM PETREL AND SMALL PASSERINES AT THE BROTHERS LIGHTHOUSE

The night of 17/5/61 was very still and calm. At 2000 hours I was called to the lighthouse tower to examine a bird which had been caught while fluttering at the light. It was obviously a storm-petrel and was identified as belonging to the gray-backed species (*Garrodia nereis*). Before it was released measurements and sketches were made and forwarded to the Dominion Museum. The identification was confirmed. Mid-May would be an unusually late date for the much commoner White-faced Storm Petrel (*P. marina*) still to be in local waters.

At the same time two small birds were seen eating moths on the outside of the tower. They proved to be Silvereyes (*Z. lateralis*). Goldfinches (*C. carduelis*) were also heard and seen flying around.

— A. WRIGHT

★

UNUSUAL DEATH OF MOREPORK

On 3/1/61, while walking under pohutukawas on the seaward side, beyond the small pa at Kauri Point, Bay of Plenty, which was being excavated by the Auckland University Archaeological Society, I found the dessicated skeleton, with some flesh and wing and tail feathers still adhering, of a Morepork (*N. novaeseelandiae*). It was gripping a wire on a barbed-wire fence, and its neck was caught in a half-loop, where the wire went round a post, the end then twisted back around the main wire; the loop was on the free end of the wire. The bird had apparently flown into the fence in the dark, and had caught its neck in the half-loop. The feet had an "agonised" appearance in their grip on the wire, and were difficult to free without breaking.

Moreporks were plentiful in the vicinity.

R. J. SCARLETT

★

KIWIS IN BAY OF PLENTY

Four specimens of the North Island Kiwi (*Apteryx australis mantelli*) were reported in eastern Bay of Plenty between June and September, 1959.

A pair were injured by dogs near Oponae in June, and were placed in the care of the late Mr. J. D. Clark at Opotiki. On his death these were taken over by Mr. P. F. Farnworth at Whakatane, where I twice had the opportunity of examining the birds. Both made good progress, and one was remarkable for its very dark colouring, almost black. The birds were later sent to the Auckland Zoo.

In September a third kiwi was found suffering injuries caused either by a dog or through being struck by a car. It was attended for several weeks by the veterinary surgeons at the Rangitaiki Plains Dairy Co. Ltd., at Edgecumbe, but finally died. The ranger, Mr. D. S. Mackay, collected the dead bird.

At the end of September a kiwi chick, believed to be only about a week old, was killed by a fox terrier on the farm of Mr. F. Mansell at Awakeri. Mr. Mansell previously found two kiwis on his property a few years ago, and says he has noted signs of the birds having scratched under logs for food.

W. T. PARHAM

REPRINTS IN LIBRARY

- Alexander, H. G.: Report on the bird-song inquiry organised by the British Trust for Ornithology. *British Birds*, vol. 36, p. 65-109.
- Alexander, W. B.: The index of heron population, 1942. *British Birds*, v. 36, p. 206-8.
 The index of heron population, 1943. *British Birds*, v. 37, p. 205-6.
 The index of heron population, 1944. *British Birds*, v. 38, p. 232-4.
 The index of heron population, 1946. *British Birds*, v. 40, p. 204-7.
 The index of heron population, 1947. *British Birds*, v. 41, p. 146-8.
 The index of heron population, 1948. *British Birds*, v. 42, p. 81-83.
 The index of heron population, 1949. *British Birds*, v. 43, p. 78-80.
 The index of heron population, 1950. *British Birds*, v. 44, p. 123-126.
 The index of heron population, 1951. *British Birds*, v. 45, p. 64-67.
 The index of heron population, 1952. *British Birds*, v. 46, p. 100-103.
- Alexander, W. B., and others: Observations on the breeding birds of Lundy in 1942. *British Birds*, v. 38, p. 182-191.
- Alexander, W. B.: The ornithology of inn signs. *Bird Notes*, v. 26, no. 1, p. 26-29.
 1945-47: The woodcock in the British Isles. *Ibis*, v. 87-89.
- Allen, F. G. H., 1956: The nesting of the Bay-headed Bee-eater. *Malayan Nature Journal*, v. 10, May 1956.
 1955: The nesting of the Bluewinged Pitta (*Pitta megarhyncha* Schleg). *Malayan Nature Journal*, v. 10, p. 67-68, Dec. 1955.
 1953: The nesting of the Whitecrowned Forktail, *Enicurus leschenaulti frontalis*. *Malayan Nature Journal*, v. 8, p. 107-8, Dec. 1953.
 1957: Some notes on the birds of Cameron Highlands. *Malayan Nature Journal*, v. 11, p. 39-41, March 1957.
- Amadon, Dean, 1950: Australian mud nest builders. *Emu*, v. 50, p. 123-127, 1950.
 1943: Bird weights as an aid in taxonomy. *Wilson Bulletin*, v. 55, no. 3, Sept. 1943, p. 164-177.
- Armstrong, E. A., 1958: Notes on the wren in the Aran Islands, Ireland. *British Birds*, v. 51, Jan. 1958, p. 29-35.
- Arnold, M., and others, 1953: Bagueage d'oiseaux migrateurs au Cap Bon (printemps 1953). *Bulletin de la Societe des Sciences Naturelles de Tunisie*, t. 6, 1952-1953, p. 217-222.
 1955: Bagueage d'oiseaux migrateurs au Cap-Bon. *Bulletin de la Societe des Sciences Naturelles de Tunisie*, t. 8, 1955, fasc. 3-4, p. 337-340.
 1953: Liste commentee des oiseaux bagues repris en Tunisie (1936-1953). *Bulletin de la Societe des Sciences Naturelles de Tunisie*, t. 6, 1952-1953, p. 205-215.
- Bechervaise, J., 1954: Birds of Heard Island. *Victorian Naturalist*, v. 71, p. 71-72.
- Bell, R. S., 1912: Breeding habits of White Tern (*Gygis alba*) on Kermadec Group. *Emu*, v. 12, p. 26-30.
- Bernhoft-Osa, A., 1956: Fuglenotiser fra Bjan i Setesdal og Norefjell i Buskerud. *Stavanger Museums Arbok*, 1956, p. 159-164.
 1957: Noen bidrag til jaerens fuglefauna. *Stavanger Museums Arbok*, 1957, p. 113-118.
- Betts, M. M., 1955: The behaviour of a pair of Great Tits at the nest. *British Birds*, v. 48, Feb. 1955, p. 77-82.

- 1958: The behaviour of adult tits towards other birds and mammals near the nest. *British Birds*, v. 51, Nov. 1958, p. 426-429.
- 1954: Experiments with an artificial nestling. *British Birds*, v. 47, p. 229-231.
- 1955: The food of titmice in oak woodland. *Journal of Animal Ecology*, v. 24, no. 2, November 1955, p. 282-323.
- 1956: Further experiments with an artificial nestling gape. *British Birds*, v. 49, p. 213-215.
- 1956: A list of insects taken by titmice in the forest of Dean (Gloucestershire). *Entomologist's Monthly Magazine*, v. 92, p. 67-71.
- Betts, M. M., 1958: Notes on the life-history of *Ernarmonia conicolana* (Heyl.) (Lep., Eucosmidae). *Entomologist's Monthly Magazine*, v. 94, p. 134-137.
- Braestrup, F. W., 1956: The significance of the strong "Oceanic" affinities of the vertebrate fauna on Rennell Island. *Natural History of Rennell Island, British Solomon Islands*, v. 1, p. 135-148.
- Buddle, G. A., 1941: Birds of the Poor Knights. *Emu*, v. 41, p. 56-58.
- 1938: Notes on the birds of Canton Island. *Records of the Auckland Institute and Museum*, v. 2, no. 3, p. 125-132.
- 1946: A second visit to the Poor Knights. *Emu*, v. 45, p. 315-318.
- Bull, P. C., 1948: Field notes on waders in the South-West Pacific with special reference to the Russell Islands. *Emu*, v. 47, p. 165-176.
- 1946: Notes on the breeding cycle of the Thrush and Blackbird in New Zealand. *Emu*, v. 46, p. 198-208.
- 1953: Observations on a marked population of Blackbirds at Lower Hutt. *Notornis*, v. 5, no. 5, p. 149-156.
- 1943: The occurrence of *Pterodroma leucoptera* in New Zealand. *Emu*, v. 42, p. 145-152.
- Burton, J. F., and Owen, D. F., 1954: Insect migration in S.W. France. *Entomologist's Monthly Magazine*, v. 90, p. 66-69.
- Castan, R., 1955: Baguage d'oiseaux migrateurs a Gabes au cours du printemps de 1955. *Bulletin de la Societe des Sciences Naturelles de Tunisie*, v. 8, fasc. 3-4, p. 309-312.
- Clark, V. I., and Fleming, C. A., 1948: Hutton's Shearwater (*Puffinus gavia huttoni* Math.) in New Zealand. *N.Z. Bird Notes*, v. 2, no. 8, April, 1948, p. 187-8.
- Clark, G. M., 1958: One new and one previously unreported species of nasal mite (Acarina, Speleognathidae) from North American birds. *Proc. of Helminthological Society of Washington*, v. 25, no. 2, p. 78-86.
- Clark, W. C., and Dawson, E. W., 1957: The Trans-Tasman dispersal of the White-fronted Tern (*Sterna striata*, Gm) *Notornis*, v. 7, no. 3, p. 65-69.
- Cowan, A. B., and Herman, C. M.: Winter losses of Canada Geese at Pea Island, North Carolina. *Proc. of the Southeastern Assoc. of Game and Fish Commissioners Meeting, Daytona Beach, Florida, Oct. 2-5, 1955*, p. 172-4.
- Cunningham, J. M., 1946: The flocking of the Silvereye; with special reference to 1944. *Emu*, v. 45, p. 212-223.
- 1946: Methods of Silvereye trapping in New Zealand. *Emu*, v. 46, p. 209-214.
- 1948: Seabird mortality in February, 1947. *N.Z. Bird Notes*, v. 2, no. 8, April, p. 188-193.

- Dawson, E. W., 19—: Advances in the knowledge of the small subfossil birds of New Zealand, and their biographical significance. From XVth International Congress of Zoology, Sec. V, Paper 22.
 1950: Bird notes from Little Barrier. *Notornis*, v. 4, p. 27-31.
 1951: Bird notes from Stewart Island. *Notornis*, v. 4, p. 146-149.
 1955: The birds of the Chatham Islands 1954 Expedition. v. 6, p. 78-82.
 1958: Food of young Black-billed Gulls (*Larus bulleri*) in a breeding colony, North Canterbury. *Notornis*, v. 8, p. 32-38.
 1957: The generic name of the extinct New Zealand gallinule. *Nature*, v. 179, p. 1307-8.
 1958: Obituary notice: Walter Reginald Brook Oliver (1883-1957). *Proc. of the Malacological Society of London*, v. 33, Part 1.
 1958: Re-discoveries of the New Zealand subfossil birds named by H. O. Forbes. *Ibis*, v. 100, p. 232-237.
 1952: Subfossil Nestor (*Psittacidae*) from New Zealand. *Emu*, v. 52, p. 259-272.
- Delacour, J., and Mayr, E., 1945: Notes on the taxonomy of the birds of the Ptilippines. *Zoolgica*, v. 30, p. 105-117.
- Dell, R. K., 1952: The Blue Petrel in Australian waters. *Emu*, v. 51, p. 147-154.
- Ealey, E. M. H., 1954: Analysis of stomach contents of some Heard Island birds. *Emu*, v. 54, p. 204-210.
- Falla, R. A., 1948: Birds of the Solanders. *N.Z. Bird Notes*, v. 3, p. 52-55.
 1946: An undescribed form of the Black Petrel. *Records of the Cant. Museum*, v. 5, p. 111-113.
- Fisher, J., 1953: The fulmar: review. by Ralph S. Palmer. *Auk*, v. 70, p. 377-379.
- Fitter, R. S. R.: Report of the Black Redstart inquiry for 1944. *British Birds*, v. 38, p. 247-249.
- Fleming, C. A., and Wodzicki, K. A., 1952: A census of the Gannet (*Sula serrator*) in New Zealand. *Notornis*, v. 5, p. 39-78.
- Fleming, C. A., 1947: Contributions to the gannet census. III. Horuhoru gannetry (Waiheke), October 2, 1946. *N.Z. Bird Notes*, v. 2, p. 109-114.
 1950: New Zealand flycatchers of the genus *Petroica* Swainson. 2 pts. *Trans. Roy. Soc. N.Z.*, v. 78, p. 14-17 and 127-160.
 1943: Notes on the life history of the Silver-eye based on colour-banding. *Emu*, v. 42, p. 193-217.
- Fleming, C. A., and Serventy, D. L., 1943: The races of *Puffinus assimilis* in Australia and New Zealand. *Emu*, v. 43, p. 113-125.
- Fleming, C. A., 1948: The Snares Islands Expedition, 1947. *N.Z. Bird Notes*, v. 2, p. 181-184.
- Fowler, S., 1947: A landing on Pedra Branch. *Proc. Royal Zool. Soc. N.S.W.*, 1947, p. 22-6.
- Gentili, J., 1949: Foundations of Australian bird geography. *Emu*, v. 49, p. 85-129.
- Gibb, J., 1950: The breeding biology of the great and blue titmice. *Ibis*, v. 92, p. 507-539.
 1951: The birds of the Maltese Islands. *Ibis*, v. 93, p. 109-127.
 1954: Feeding ecology of tits, with notes on treecreeper and goldcrest. *Ibis* v. 96, p. 513-543.
 1954: Population changes of titmice, 1947-1951. *Bird Study*, v. 1, p. 40-48.

- 1955: Feeding rates of Great Tits. *British Birds*, v. 48, p. 49-58.
1956: Territory in the genus *Parus*. *Ibis*, v. 98, p. 420-429.
1957: Food requirements and other observations on captive tits. *Bird Study*, v. 4, p. 207-215.
- Gibb, J., and Hartley, P. H. T., 1957: Bird foods and feeding-habits as subjects for amateur research. *British Birds*, v. 50, p. 278-291.
- Gibb, J. A., 1958: Predation by tits and squirrels on the eucosmid *Ernarmonia conicolana* (Heyl.) *Journal of Animal Ecology*, v. 27, p. 375-396.
- Gifford, E. W., 1925: The Gray-hooded Quail-dove (*Gallicolumba rubescens*) of the Marquesas Islands, in captivity. *Auk*, v. 42, p. 388-396.
1927: Grayson's Pigeon (*Zenaidura graysoni*) in captivity. *Auk*, v. 44, p. 513-9.
- Glauert, L., 1949: Some Western Australian Mallophaga. *Emu*, v. 49, p. 129-131.
- Goodacre, M. J., 1959: The origin of winter visitors to the British Isles. 3 pts. 1959. *Bird Study*, v. 6.
- Goodacre, M. J., and Lack, D., 1959: Early breeding in 1957. *British Birds*, v. 52, p. 74-83.
- Goodbody, I. M., 1952: The post-fledging dispersal of juvenile titmice. *British Birds*, v. 45, p. 279-285.
- Goodhart, C. B., 1958: Thrush predation on the snail *Cepaea hortensis*. *Journal of Animal Ecology*, v. 27, p. 47-57.
- Gregory, J. T., 1952: The jaws of the Cretaceous toothed birds, *Ichthyornis* and *Hesperornis*. *Condor*, v. 54, p. 73-78.
- Gurr, L., 1947: Measurements of birds. *N.Z. Bird Notes*, April 1947, p. 57-61.
1952: Some food of the North Island Kiwi (*Apteryx australis*). *Notornis*, v. 4, p. 209-210.
- Hartley, P. H. T., 1948: The assessment of the food of birds. *Ibis*, v. 90, p. 361-381.
1949: The biology of the Mourning Chat in winter quarters. *Ibis*, v. 91, p. 393-413.
1953: An ecological study of the feeding habits of the English titmice. *Journal of Animal Ecology*, v. 22, p. 261-288.
1954: Wild fruits in the diet of British thrushes: a study in the ecology of closely allied species. *British Birds*, v. 47, p. 97-107.
19—?: Economic ornithology — does it exist? *Bird Notes*, v. 23, No. 2.
- Herman, C. M., 1954: Haemoproteus infections in waterfowl. *Proc. of the Helminthological Society of Washington*, v. 21, p. 37-42.
- Herman, C. M., and Price, D. L.: the occurrence of Hepatazoon in the Gray squirrel (*Sciurus carolinensis*). *Journal of Protozoology*, v. 2, p. 48-51.
- Herman, C. M., and others, 1943: Food habits and intensity of coccidian infection in native valley quail in California. *Journal of Parasitology*, v. 29, p. 206-8.
1954: Studies on avian malaria in vectors and hosts of encephalitis in Kern County, California. I. Infections in avian hosts. *American Journal of Tropical Medicine and Hygiene*, v. 3, p. 676-695.
1955: Causes of winter losses among Canada geese. *Trans. of the 20th North American Wildlife Conference*, p. 161-165.

- Hoffman, K., 1958: Repetition of an experiment on bird orientation. *Nature*, v. 181, p. 1435-7.
- Holgersen, H., 1955: 24 ars ringmerkingsrapporter. Stavanger Museums Arbok 1955, p. 141-152.
- 1955: Utvidelse av Ornitologisk Stasjon, Revtangen. Stavanger Museums Arbok 1955, p. 135-140.
- 1956: Kortnebbgjess (*Anser arvensis brachyrhynchus*) i den kalde ettervinteren 1956. Stavanger Museums Arbok 1956, p. 151-158.
- 1957: Sjeldne gjester i den Norske fuglefauna. Stavanger Museums Arbok, 1957, p. 103-111.
- 1960: Fugl i Norske Biland. - 2. Bouvet-ya. *Sterna*, v. 4, p. 1-9.
- Hope, C. E. and Shortt, T. M.: Southward migration of adult shorebirds on the west coast of James Bay, Ontario. *Auk*, v. 61, p. 572-576.
- Howard, H., 1950: Fossil evidence of avian evolution. *Ibis*, v. 92, p. 1-21.
- 1955: New records and a new species of *Chendytes*, an extinct genus of diving geese. *Condor*, v. 57, p. 135-143.
- Howard, P. F., 1954: Banding of the Black-browed Albatross at Heard Island and Macquarie Island.
- 1956: Banding of Giant Petrels at Heard Island and Macquarie Island — II. *Emu*, v. 56, p. 401-4.
- Hurrell, H. G., 1948: Simultaneous watch for migrant Swifts, May 11th, 1947. *British Birds*, v. 41, p. 138-145.
- Johansen, H., 1959: Resultater af ringmaerkningerne i 1953 og 1954, Vidensk. Medd. fra Dansk Naturh. Foren., Bd. 121, p. 75-139.
- Junge, G. C. A., 1952: Resultaten van het ringonderzoek betreffende de vogeltrek, ingesteld door het Rijksmuseum van Natuurlijke Historie te Leiden, XXXVII (1950). 2 pts. *Limosa*, v. 25, p. 66-85, 113-131.
- Klopfer, P. H., 1958: Influence of social interactions on learning rates in birds. *Science*, v. 128, p. 903.
- Koskimies, J., 1957: Polymorphic variability in clutch size and laying date of the Velvet Scoter, *Melanitta fusca* (L.) *Ornis Fennica*, v. 34, p. 118-128.
- 1957: Terns and gulls as features of habitat recognition for birds nesting in their colonies. *Ornis Fennica*, v. 34, p. 1-6.
- 1957: Variations in size and shape of eggs of the Velvet Scoter, *Melanitta fusca* (L.) *Archivum Societatis Zoologicae Botanicae Fennicae 'Vanamo'*, v. 12, p. 58-69.
- Krieg, H., and Schuhmacher, E., 1936: Beobachtungen an sudamerikanischen Wildhuhnern. *Verh. Orn. Ges. Bay* v. 21, p. 1-18.
- Lack, D., 1943: The age of some more British Birds. *British Birds*, v. 36, p. 193-221.
- 1943: The age of the Blackbird. *British Birds*, v. 36, p. 166-175.
- 1944: The problem of partial migration. *British Birds*, v. 37, p. 122-150.
- 1946: Clutch and brood size in the Robin. *British Birds*, v. 39, p. 98-135.
- 1946: Competition for food by birds of prey. *Journal of Animal Ecology*, v. 15, p. 123-129.
- 1946: Do juvenile birds survive less well than adults? *British Birds*, v. 39, p. 258-264.
- 1946: The names of the Geospizinae (Darwin's finches). *Bulletin of the British Ornithologists' Club*, b. 67, p. 16-22.

- 1947: A further note on the taxonomy of the Robin, *Erithacus rubecula* (Linnaeus). Buletin of the British Ornithologists' Club, v. 67, p. 51-54.
- 1947-48: The significance of clutch-size. 3 pts. *Ibis*, v. 89, p. 302-353, v. 80, p. 25-45.
- 1947: The significance of clutch-size in the Partridge (*Perdix perdix*). *Journal of Animal Ecology*, v. 16, p. 19-25.
- 1948: Further notes on clutch and brood size in the Robin. *British Birds*, v. 41, p. 98-100 and 130-137.
- 1948: Natural selection and family size in the Starling. *Evolution*, v. 2, p. 95-110.
- 1948: Notes on the ecology of the Robin. *Ibis*, v. 90, p. 252-279.
- 1948: The significance of litter-size. *Journal of Animal Ecology*, v. 17, p. 45-50.
- 1949: The apparent survival-rate of ringed Herons. *British Birds*, v. 42, p. 74-9.
- 1949: Comments on Mr. Skutch's paper on clutch-size. *Ibis*, v. 91, p. 455-8.
- 1949: Family size in certain thrushes (*Turdidae*). *Evolution*, v. 3, p. 57-66.
- 1949: The significance of ecological isolation. *Genetics, paleontology, and evolution*, ed. by G. L. Jepsen and others.
- 1949: Vital statistics from ringed Swallows. *British Birds*, v. 42, p. 147-150.
- 1950: Birds and the field centres. C.P.F.S. Annual Report, 1949-50, p. 24-33.
- 1950: Breeding seasons in the Galapagos. *Ibis*, v. 92, p. 268-278.
- 1950: The breeding seasons of European birds. *Ibis*, v. 92, p. 288-316.
- 1950: Family-size in titmice of the genus *Parus*. *Evolution*, v. 4,
- 1950: Population ecology in birds: a review. *Proc. of the Xth International Ornithological Congress, Uppsala, 1950*, p. 409-448.
- 1951: The breeding biology of the Swift *Apus apus*. *Ibis*, v. 93, p. 501-546.
- 1951: Geographical variation in *Erithacus rubecula*. *Ibis*, v. 93, p. 629-630.
- 1951: Migration through the Pyrenees. *Bulletin of the British Ornithologists' Club*, v. 70.
- 1952: Reproductive rate and population density in the Great Tit: Klijver's study. *Ibis*, v. 94, p. 167-173.
- 1954: Call-notes, *Erithacus* and convergence. *Ibis*, v. 96, p. 312-314.
- 1954: The evolution of reproductive rates. *Evolution as a process*, by J. S. Huxley, p. 143-156.
- 1954: The stability of the Heron population. *British Birds*, v. 47, p. 111-121.
- 1954: Two Robin populations. *Bird Study*, v. 1, p. 14-17.
- 1954: Visible migration in S.E. England, 1952. *British Birds*, v. 47, p. 1-15.
- 1955: British tits (*Parus* spp.) in nesting boxes. *Ardea*, v. 43, p. 50-84.
- 1955: Summary report on nesting tits. *Bird Study*, v. 2, p. 199-201.
- 1955: The summer movements of Swifts in England. *Bird Study*, v. 2, p. 32-40.

- 1955: Visible migration through the Pyrenees. *Acta XI Cong. Int. Orn.*, 1954, p. 176-8.
- 1956: Further notes on the breeding biology of the Swift *Apus apus*. *Ibis*, v. 98, p. 606-619.
- 1956: A review of the genera and nesting habits of Swifts. *Auk*, v. 73, p. 1-32.
- 1956: Seaward flights of Swifts at dusk. *Bird Study*, v. 3, p. 37-42.
- 1956: The species of *Apus*. *Ibis*, v. 98, p. 34-62.
- 1956: Variations in the reproductive rate of birds. *Proc. of the Royal Society, B*, v. 145, p. 329-333.
- 1957: The Chaffinch migration in North Devon. *British Birds*, v. 50, p. 10-19.
- 1957: The first primary in Swifts, *Auk*, v. 47, p. 385-6.
- 1957: Notes on nesting Nightjars. *British Birds*, v. 50, p. 273-7.
- 1958: Migrational drift of birds plotted by radar. *Nature*, v. 182, p. 221-3.
- 1958: A new race of the Whiterumped Swift. *Journal of the Bombay Natural History Society*, April 1958.
- 1958: A quantitative breeding study of British tits. *Ardea*, v. 46, p. 91-124.
- 1958: The return and departure of Swifts *Apus apus* at Oxford. *Ibis*, v. 100, p. 477-502.
- 1958: The significance of the colour of turdine eggs. *Ibis*, v. 100, p. 145-166.
- 1958: Swifts over the sea at night: weather movements of Swifts, 1955-1957. *Bird Study*, v. 5, p. 126-142.
- 1959-60: Migration across the North Sea studied by radar. 2 pts. *Ibis*, v. 101 and 102.
- 1959: Migration and orientation: migration across the sea. *Ibis*, v. 101, p. 374-399.
- 1959: Some British pioneers in ornithological research, 1859-1939. *Ibis*, v. 101, p. 71-81.
- 1959: Watching migration by radar. *British Birds*, v. 52, p. 258-267.
- 19—?: The position of the Edward Grey Institute, Oxford. *Bird Notes*, v. 23, p. 231-6.
- Lack, D. and Arn, H., 1947: Die Bedeutung der Gelegegrosse beim Alpensegler. *Der Ornithologische Beobachter*, v. 44, p. 190-210.
- 1953: Die mittlere Lebensdauer des Alpenseglers. *Der Ornithologische Beobachter*, v. 50, p. 133-137.
- Lack, D. and Lack, E., 1949: Passerine migration through England. *British Birds*, v. 42, p. 320-326.
- 1951: Migration of insects and birds through a Pyrenean pass. *Journal of Animal Ecology*, v. 20,
- 1952: The breeding behaviour of the Swift. *British Birds*, v. 45, p. 186-215.
- 1952: Further changes in bird-life caused by afforestation. *Journal of Animal Ecology*, v. 20, p. 173-9.
- 1952: Visible migration at Land's End. *British Birds*, v. 45, p. 81-96.
- 1953: Visible migration through the Pyrenees: an autumn reconnaissance. *Ibis*, v. 95, p. 271-309.
- 1958: The nesting of the Long-tailed Tit. *Bird Study*, v. 5, p. 1-19.
- Lack, D., and Owen, D. F., 1955: The food of the Swift. *Journal of Animal Ecology*, v. 24, p. 120-136.

- 1957: Survival in relation to brood-size in tits. *Proc. Zoological Soc. London*, v. 128, p. 313-326.
- Lack, D., and Ridpath, M. G., 1955: Do English woodpigeons migrate. *British Birds*, v. 48, p. 289-292.
- Lack, D., and Schifferli, A., 1948: Die Lebensdauer des Stares. *Der Ornithologische Beobachter*, v. 45, p. 107-114.
- Lack, D., and Silva, E. T., 1949: The weight of nestling robins. *Ibis*, v. 91, p. 64-78.
- Lack, D., and Southern, H. N., 1949: Birds on Tenerife. *Ibis*, v. 91, p. 607-626.
- Lack, D., and Varley, G. C., 1945: Detection of birds by radar. *Nature*, v. 156, p. 446.
- Lack, D., and Williamson, K., 1959: Bird-migration terms. *Ibis*, v. , p. 255-6.
- Lack, E. 1950: Breeding season and clutch-size of the wood warbler. *Ibis*, v. 92, p. 95-8.
- Lampio, T., 1945: Uhtuan etela-, lansi- ja pohjoispuolen linnustosta. *Ornis Fennica*, v. 22, p. 45-56.
- 1946: Havaintoja ita-karjalan riistaelaimistosta. *Suomalaisen Elain- ja Kasvitieteellisen Seuran Vanamon Elaintieteellisia Julkaisuja*, Osa 11, p. 19-31.
- 1946: Tvarminnen elaintieteellisen aseman ja sen lahiympariston linnuston viimeaikaisesta kehityksesta. *Ornis Fennica*, v. 23, p. 33-49.
- 1947: Taman talven turkisriistatilanteesta [On the stocks of fur game in Finland in winter 1947/48] *Metsalehdesta*, No. 47-48.
- 1948: Luontaiset edellytykset naame oravatalouden perustana: Squirrel economy in Finland based on natural prerequisites. *Suomen Riista II*, p. 97-147.
- 1949: Piirteita ketun ravinnosta totuus repolaisesta paljastumassa [On the food of red fox in Finland]. *Metsastys ja Kalastuslehdesta*, No. 5.
- 1949: Riistamailla lieva tautivuosi. [The game disease year 1948/49]. *Metsalehdesta*, No. 7.
- 1949: Samsonfragan [The Samson fox problem]. *Finlands Jakt-och Fisketidskrift*, No. 2, p. 78-84.
- 1950: Heinakurpan esiintymisestä Suomessa. *Ornis Fennica*, v. 27, p. 31-38.
- Lane, G. B., and Wodzicki, K. A., 1944: A visit to the Ohau River Estuary, New Zealand, on 12th March, 1944. *Avicultural Magazine*, No. 7.
- Lockie, J. D., 1955: The breeding and feeding of Jackdaws and Rooks with notes on Carrion Crows and other Corvidae. *Ibis*, v. 97, p. 341-369.
- 1955: The breeding habits and food of Short-eared Owls after a vole plague. *Bird Study*, v. 2, p. 53-69.
- 1956: The food and feeding behaviour of the Jackdaw, Rook and Carrion Crow. *Animal Ecology*, v. 25, p. 421-428.
- 1956: Winter fighting in feeding flocks of Rooks, Jackdaws and Carrion Crows. *Bird Study*, v. 3, p. 180-190.
- Lockwood, A. P. M., 1959: The extra-haemolymph sodium of *Asellus aquaticus* (L.). *Journal of Experimental Biology*, v. 36, p. 562-565.
- McCartan, L., and Simmons, K. E. L., 1956: Territory in the Great Crested Grebe (*Podiceps cristatus*) re-examined. *Ibis*, v. 98, p. 370-378.

- Madoc, G. C., and Allen, F. G. H., 1952: Ornithological notes made at Fraser's Hill, March 1951. *Buletin of the Raffles Museum, Singapore*, No. 24, p. 164-182.
- Marler, P., 1957: Specific distinctiveness in the communication signals of birds. *Behaviour*, v. 11, p. 13-39.
- Marples, B. J., 1942: A study of the Little Owl, *Athene noctua*, in New Zealand. *Trans. Royal Soc. of N.Z.*, v. 72, p. 237-252.
- Marples, B. J., and Gurr, L., 1943: A mechanism for recording automatically the nesting habits of birds. *Emu*, v. 43, p. 67-71.
- Mason, C. R.: Do you know your birds? *Buletin of the Massachusetts Audubon Society*.
- Mason, E. A.: What feather is that? An old evil revived in the millinery trade. *Ibid*.
- Invite your bird neighbors to dinner. *Ibid*.
- Mason, C. R., and Taylor, W. A.: Help the birds find a home. *Ibid*.
- Matvejev, S., and Sigunov, A., 1952-3: Climographs of East Yugoslavia and samples of their Application in the ecology. *Recuel des Traveux de l'Acad. Serbe des Sciences*, 29. *Institut d'Ecologie et de Biogeographie* No. 3. In Russian with English summary.
- Mayaud, N., 1955: *Bibliographie*. *Alauda*, v. 23, p. 139-144.
- Mayr, E., 1944: Bird watching, a hobby and a science. *Auk*, v. 61, p. 151-2.
- 1944: The family name of the Australian honey-eaters. *Auk*, v. 61, p. 465.
- 1946: The number of species of birds. *Auk*, v. 65, p. 64-9.
- 1946: What needs to be learned about Australian ducks. *Emu*, v. 45, p. 229-232.
- 1949: A second Australian record of the Corn Crake (*Crex crex*). *Emu*, v. 48, p. 243.
- 1950: Speciation in birds. *Proc. Xth International Ornith. Congr., Uppsala, 1950*, p. 91-131.
- 1952: German experiments on orientation of migrating birds. *Biological Reviews*, v. 27, p. 394-400.
- 1952: The problem of land connections across the South Atlantic, with special reference to the Mesozoic. *Introduction and Conclusion 2*. *Bulletin of the American Museum of Nat. Hist.*, v. 99, art. 3 (*Proc. of the Symposium on the South Atlantic Basin, 1949*).
- 1952: *Turdus Musicus Linnaeus*. *Ibis*, v. 94, p. 531-4.
- 19—?: *Artbildung und Variation in der Halcyon-Chloris-Gruppe*. *Ornithologie als biologische Wissenschaft. Festschrift zum 60 Geburtstag von Erwin Stresemann*, p. 55-60.
- Mayr, E., and Gilliard, E. T., 1952: Altitudinal hybridization in New Guinea honeyeaters. *Condor*, v. 54, p. 325-337.
- Moreau, R. E., 1948: Ecological isolation in a rich tropical avifauna. *Journal of Animal Ecology*, v. 17, p. 113-126.
- 1949: Friedmann on African cuckoos. *Special review*. *Ibis*, v. 91, p. 529-37.
- 1950: The migration system in perspective. *Proc. Xth Int. Ornith. Congr., Uppsala, 1950*, p. 245-8.
- 1949: The breeding of a Paradise Flycatcher. *Ibis*, v. 91, p. 256-279.
- 1950: The breeding season of African birds — 1. Land birds. *Ibis*, v. 92, p. 223-267.

- 1951: Africa since the Mesozoic: with particular reference to certain biological problems. *Proc. Zool. Soc. Lond.* v. 121, p. 869-913.
- 1951: The British status of the Quail and some problems of its biology. *British Birds*, v. 44, p. 257-276.
- 1951: Geographical variation and plumage sequence in *Pogonocichla Ibis*, v. 93, 383-401.
- 1952: The place of Africa in the Palaearctic migration system. *Journal of Animal Ecology*, v. 21, p. 250-271.
- 1953: Migration in the Mediterranean area. *Ibis*, v. 95, p. 329-364.
- 1954: The distribution of African evergreen-forest birds. *Proc. Linnean Soc. London*, Session 165, p. 35-46.
- 1954: The main vicissitudes of the European avifauna since the Pliocene. *Ibis*, v. 96, p. 411-431.
- 1955: Ecological changes in the Palaearctic region since the Pliocene. *Proc. Zool. Soc. London*, v. 125, p. 253-295.
- 1956: The Iberian Peninsula and migration. *Bird Study*, v. 3, p. 1-25.
- 1956: Quail in the British Isles, 1950-53. *British Birds*, v. 49, p. 161-6.
- 1958: The *Malimbus* spp. as an evolutionary problem. *Revue de Zoologie et de Botanique africaines*, v. 57, p. 241-255.
- 1958: Some aspects of the Musophagidae. *Ibis*, v. 100, p. 67-112.
- 1959: Les problemes de la migration a travers le Maroc. *Alauda*, v. 27, p. 81-96.
- 1959: Weaver-bird problems. *Ibis*, v. 101, p. 334-335.
- 19—?: The African Mountain Wagtail *Motacilla clara* at the nest. *Ornithologie als biologische Wissenschaft (Festschrift zum 60. Geburtstag von Erwin Stresemann)* p. 183-191.
- Moreau, R. E., and Chapin, J. P., 1951: The African Emerald Cuckoo, *Chrysococcyx cupreus*. *Auk*, v. 68,
- Moreau, R. E., and Monk, J. F., 1957: Autumn migration in southwest Portugal. *Ibis*, v. 99, p. 500-8.
- Moreau, R. E., and Moreau, M. W., 1954: Notas otonales sobre aves de Levante e islas Pityusas. *Ardeola*, v. 1, p. 3-32.
- 1956: Acerca de la migracion otonal en el Estrecho de Gibraltar. *Ardeola*, v. 1, p. 59-69.
- Myers, M. T., 1955: The breeding of Blackbird, Song Thrush and Mistle Thrush in Great Britain. Part I. Breeding Season. *Bird Study*, v. 2, p. 2-24.
- Nordstrom, G., 1957: Die Vogelberingung in Finnland im Jahre 1956. *Memoranda Societatis pro Fauna et Flora Fennica*, v. 33, p. 69-108.
- Norris, C. A., 1945: Summary of a report on the distribution and status of the Corn-Crake. *British Birds*, v. 38, p. 142-8, 162-8.
- 1947: Report on the distribution and status of the Corn-Crake. Part 2. A consideration of the causes of the decrease. *British Birds*, v. 40, p. 226-244.
- Old brecks or new forests? A protest to the Editor of *Bird Notes* from Thomas Titmouse, Esq. and the Hon. Reg. *Regulus* (Thetford Chase, Norfolk). *Bird Notes*, v. 24, No. 6.
- Oliver, W. R. B., 1912: Geographic relationships of the birds of Lord Howe, Norfolk, and the Kermadec Islands. *Trans. N.Z. Inst.*, v. 44, p. 214-221.

- 1913: Further notes on the birds of the Kermadec Islands. *Trans. N.Z. Inst.*, v. 45, p. 92-3.
- 1922: The birds of Little Barrier Island, New Zealand. *Emu*, v. 22, p. 45-51.
- 1930: The New Zealand double-crested shags: with description of a new species. *Trans. Roy. Soc. N.Z.*, v. 61, p. 138-9.
- Owen, D. F., 1953: Migration at the Kentish Knock lightship. *British Birds*, v. 46, p. 353-64.
- 1954: Protocalliphora in birds' nests. *British Birds*, v. 47, p. 236-43.
- 1954: The Swift *Apus apus* L. as a predator of aphids. *Journal of Soc. for British Entomology*, v. 5; p. 82-5.
- 1954: The winter weights of titmice. *Ibis*, v. 96, p. 299-309.
- 1955: Additional records of Protocalliphora (Diptera) in birds' nests. *British Birds*, v. 48, p. 225-229.
- 1955: Coleoptera taken by Swifts (*Apus apus* L.). *Journal of Soc. for British Entomology*, v. 5, p. 105-9.
- 1955: The food of the Heron *Ardea cinerea* in the breeding season. *Ibis*, v. 97, p. 276-295.
- 1956: The food of nestling Jays and Magpies. *Bird Study*, v. 3, p. 257-65.
- 1957: Neottiophilum praeustum in birds' nests. *British Birds*, v. 50, p. 160-4.
- 1959: The breeding season and clutch-size of the Rook *Corvus frugilegus*. *Ibis*, v. 101, p. 235-9.
- Owen, D. F., and Burton, J. F., 1953: Observations sur la migration dans le Sud-Ouest de la France. *Alauda*, v. 21, p. 223-239.
- Owen, D. F., and Le Gros, A. E., 1954: Spiders caught by Swifts. *Entomologist's Gazette*, v. 5, p. 117-120.
- Owen, D. R., and others, 1955: Observaciones ornitologicas otonales en el nore de Espana. *Ardeola*, v. 2, p. 57-78.
- Parkes, K. C., 1959: The northeastern races of the Long-billed March Wren (*Telmatodytes palustris*). *Annals of the Carnegie Museum*, v. 35, p. 275-281.
- Parkhurst, P., and Lack, D., 1946: The clutch-size of the Yellowhammer. *British Birds*, v. 39, p. 358-364.
- Parmenter, L., and Owen, D. F., 1954: The Swift, *Apus apus* L., as a predator of flies. *Journal of the Soc. for British Entomology*, v. 5, p. 27-33.
- Phillips, W. J., 1947: The Maori method of taking Grey Duck and Black Teal near Taupo. *Journ. Polynesian Soc.*, v. 56, p. 333-5.
- 1948: Bird life on West Coast, South Island. I. Notes on the birds of Reefton. *N.Z. Bird Notes*, v. 3, p. 18.
- 1948: Bird Notes from Mahia Peninsula. *N.Z. Bird Notes*, v. 3, p. 17.
- 1948: Note on Maori method of taking weka in North Canterbury. *Journal Polyn. Soc.*, v. 57, p. 177-8.
- 1950: Maori "bird calls" or "whistles." *Ethnos*, 1950, p. 201-5.
- Phillips, W. J., and Lindsay, C. J., 1948: Preliminary list of birds of the Rotorua acclimatisation district. *N.Z. Bird Notes*, Oct. 1948, p. 43-50.
- Probine, M. C., and Wodzicki, K. A., 1955: A note on the thermal resistance of the feathered layer of gannet skins. *N.Z. Journ. Science and Techn. B*, v. 37, p. 158-9.

- Richdale, L. E., 1944: The titi wainui or Fairy Prion, *Pachyptila turtur* (Kuhl.) 2 pts. *Trans. Roy. Soc. N.Z.*, v. 74, p. 32-48, 165-181.
1944: The White-faced Storm Petrel. Part III. *Trans. Roy. Soc. N.Z.*, v. 78, p. 335-350.
1945: Supplementary notes on the diving petrel. *Trans. Roy. Soc. N.Z.*, v. 75, p. 42-53.
- Ripley, S. D., 1954: Comments on the biogeography of Arabia with particular reference to birds. *Journal Bombay Nat. Hist. Soc.*, v. 52, p. 241-8.
1954: Olathe Quail in Utah. *Wilson Bulletin*, v. 66, p. 139.
1955: Anatomical notes on *Zavattariornis*. *Ibis*, v. 97, p. 142-5.
1955: Some comments on Vaurie's revision of the *Muscicapini*. *Auk*, v. 72, p. 86-8.
- Robertson, F. H., and Wodzicki, K., 1948: Contributions to the gannet census. XI. White Island gannetries. *N.Z. Bird Notes*.
- Scarlett, R. J., 1953: A sub-fossil hawk from New Zealand. *Records of Canterbury Museum*, v. 6, p. 245-252.
- Schuz, E., 1955: Über den Altersaufbau von Weisstorch-Populationen. *Acta XI Congr. Int. Orn.*, p. 522-8.
- Secker, H. L., 1955: Notes on the New Zealand Pipit. *Emu*, v. 55, p. 104-8.
- Shortt, T. M., 1951: On the juvenile plumage of North American pipits. *Auk*, v. 68, p. 265.
- Sibson, R. B., 1943: Additions to the avifauna of the North Island, New Zealand. *Emu*, v. 43, p. 136-40.
1943: Observations on the distribution of the Wrybill in the North Island, New Zealand. *Emu*, v. 43, p. 49-62.
1945-6: Notes on birds of Sandbach, S.E. Cheshire. *Northwestern Naturalist*, Sept.-Dec. 1945, p. 247-255, March-June 1946, p. 58-67.
- Silva, E. T., 1949: Nest records of the Song-thrush. *British Birds*, v. 42, p. 97-111.
- Sladen, W. J. L., 1953: The Adelie Penguin. *Nature*, v. 171, p. 952.
- Snow, D. W., 1952: A contribution to the ornithology of North-West Africa. *Ibis*, v. 94, p. 473-498.
1955: The abnormal breeding of birds in the winter 1953/54. *British Birds*, v. 48, p. 120-126.
1952: The winter avifauna of Arctic Lapland. *Ibis*, v. 94, p. 133-143.
1953: The migration of the Greenland Wheatear. *Ibis*, v. 95, p. 376-378.
1953: Visible migration in the British Isles: a review. *Ibis*, v. 95, p. 242-270.
1954: The habits of Eurasian tits (*Parus* spp.). *Ibis*, v. 96, p. 565-585.
1954: Trends in geographical variation in Palaearctic members of the genus *Parus*. *Evolution*, v. 8, p. 19-28.
1955: Geographical variation of the Coal Tit, *Parus ater* L. *Ardea*, v. 43, p. 195-226.
1956: Territory in the Blackbird *Turdus merula*. *Ibis*, v. 98, p. 438-447.

- 1955: The breeding of the Blackbird, Song Thrush, and Mistle Thrush in Great Britain. Parts 2 and 3. *Bird Study*, v. 2, p. 72-84, 169-178.
- 1956: The annual mortality of the Blue Tit in different parts of its range. *British Birds*, v. 49, p. 174-7.
- 1958: The breeding of the Blackbird *Turdus merula* at Oxford. *Ibis*, v. 100, p. 1-30.
- Snow, D. W., and Manning, A. W. G., 1954: Observations sur la migration d'automne en Tunisie et Tripolitaine et dans le Fezzan (1953). *Alauda*, v. 22, p. 1-24.
- Snow, D. W., and Ross, K. F. A., 1952: Insect migration in the Pyrenees. *Entomologist's Monthly Magazine*, v. 88.
- Snow, D. W., and others, 1955: Land- and sea-bird migration in North-west Spain, autumn 1954. *Ibis*, v. 97, p. 557-571.
- Snyder, L. L., and Hope, C. E., 1938: A predator-prey relationship between the Short-eared Owl and the meadow mouse. *Wilson Bulletin*, v. 50, p. 110-112.
- Southern, H. N. and Reeve, E. C. R., 1941: Quantitative studies in the geographical variation of birds — The Common Guillemot (*Uria aalge* Pont.). *Proc. Zool. Soc. Ser. A*, v. 111, p. 255-276.
- Southern, H. N., and Tucker, B. W., 1945: The Manx Shearwater on Lundy. *British Birds*, v. 38, p. 124-9.
- Sowls, L. K., 1950: Techniques for waterfowl-nesting studies. *Trans. 15th North American Wildlife Conference, 1950*, p. 478-489.
- Spencer, R., 1955: Report on bird-ringing for 1954. *British Birds*, v. 48, p. 461-498.
- Stonehouse, B., 1956: The King Penguin of South Georgia. *Nature*, v. 178, p. 1424-6.
- 1958: Notes on the ringing and the breeding distribution of the Giant Petrel *Macronectes giganteus*. *Ibis*, v. 100, p. 204-8.
- Tedd, J. G., and Lack, D., 1958: The detection of bird migration by high-power radar. *Proc. Roy. Soc. London, B.*, v. 149, p. 503-510.
- Thomas, J. F., 1943: Report on the Redshank inquiry 1939-40 organised by the British Trust for Ornithology. *British Birds*, v. 36, p. 5-34.
- Thomson, A. L., 1944: The migration of the Sandwich Tern; results of British ringing. *British Birds*, v. 37, p. 62-9.
- Thomson, A. L., and Moreau, R. E., 1957: Feeding habits of the Palm-nut Vulture *Gypohierax*. *Ibis*, v. 99, p. 608-613.
- Thorpe, W. H., 1958: Further studies on the process of song learning in the Chaffinch (*Fringilla coelebs gengleri*). *Nature*, v. 182, p. 554-7.
- 1958: The learning of song patterns by birds, with special reference to the song of the Chaffinch *Fringilla coelebs*. *Ibis*, v. 100, p. 535-570.
- Turbott, E. G., 1946: A survey of Grey Duck foods. *Annual report of the Auckland Acclimatisation Society, 1946*, p. 1-13.
- Turbott, E. G., and Bull, P. C., 1954: A bird census and some recent observations on birds on Great Island, Three Kings Group. *Records Auckland Inst. & Museum*, v. 4, p. 245-262.
- Vaurie, C., 1950: Variation in *Oenanthe lugubris*. *Ibis*, v. 92, p. 540-4.
- 1955: The geographical variation of *Dicrurus hottentottus* in India. *Ibis*, v. 97, p. 153-5.

- Watson, J. S., and Wodzicki, K. A., 1954: The Myna in the Rotorua-Taupo district. *Notornis*, v. 5, p. 253.
- Wehr, E. E., and Herman, C. M., 1956: *Lophortofilaria californiensis* n.g. n.sp. (Filarioidea, Dipetalonematidae) from California Quail, *Lophortyx californicus*, with notes on its microfilaria. *Journal of Parasitology*, v. 42, p. 42-4.
- Weitnauer, E., and Lack, D., 1955: Daten zur Fortpflanzungsbiologie des Mauerseglers (*Apus apus*) in Oltingen und Oxford. *Ornithologische Beobachter*, v. 52, p. 137-141.
- Werth, I., 1947: The growth of a young Cuckoo. *British Birds*, v. 40, p. 331-4.
- 1947: The tendency of Blackbird and Song-thrush to breed in their birthplaces. *British Birds*, v. 40, p. 328-330.
- Westerskov, K., 1949: An ecological comparison between the avifaunas of Ohio and Denmark. *Ohio Journal of Science*, v. 49, p. 15-31.
- 1950: Bestandsskifte-perioden og legealderen hos lirypen. [The turnover period and life length of the willow grouse, *Lagopus lagopus*]. *Naturen*, nr. 11, p. 321-327.
- 1950: Methods for determining the age of game bird eggs. *Journal of Wildlife Management*, v. 14, p. 56-67.
- 1951: Om Aldersfordeling og Goldhed i Agerhønebestanden. *Dansk Jagttidende*, May 1951, p. 26-8. English summary.
- Willett, G., 1920: Additional notes on the avifauna of Forrester Island, Alaska. *Condor*, v. 22, p. 138-9.
- 1923: Bird records from Craig, Alaska. *Condor*, v. 25, p. 105-6.
- Williams, G. R., 1960: Distribution of specimens of the Kakapo, *Strigops habroptilus* Gray, in some museums throughout the world. *Records of the Dominion Museum*, v. 3, p. 219-227.
- 1960: The Takahe (*Notornis mantelli* Owen, 1848): a general survey. *Trans. Roy. Soc. N.Z.*, v. 88, p. 235-258.
- Williamson, K., 1948: Fair Isle Bird Observatory. First Report. Part I — Detailed notes on species. *Scottish Naturalist*, v. 61, p. 19-31.
- 1949: Work at a bird observatory. *Scottish Educational Journal*, Aug. 5, 1949.
- 1950: Fair Isle Bird Observatory. Note on selected species, autumn, 1949. *British Birds*, v. 43, p. 48-52.
- Wodzicki, K. A., 1947: Destruction of Prions by storm, August, 1946. *N.Z. Bird Notes*, April 1947, p. 64-9.
- 1956: Breeding of the House Sparrow away from man in New Zealand. *Stray Feathers*, v. 56, p. 146-7.
- Wodzicki, K. A., and Eyles, J. R., 1946: White phase of the Reef Heron or matuku-moana, *Demigretta sacra* (Gmelin) at Wairau Bay. *N.Z. Bird Notes*, v. 1, No. 10.
- Wodzicki, K. A., and McMeekan, 1947: The Gannet on Cape Kidnappers. *Trans. Roy. Soc. N.Z.*, v. 76, p. 429-452.
- Wodzicki, K. A., and Robertson, F. H., 1955: Observations on diving of the Australasian Gannet (*Sula bassana serrator* Gray). *Notornis*, v. 6, p. 72-6.
- Wood, C. A.: The fossil eggs of Bermudan birds. *Ibis*, 1923, p. 193-207.
- Zink, G., 1955: Die Verwendung der Randlochkarte für die Wiederfund-kartei. *Acta XI Congr. Int. Orn.* 1954, p. 662-4.

NOTICES

BLACK-BACKED GULL STUDY AT WELLINGTON

A study of the Black-backed Gull is being carried out at Victoria University of Wellington. In this connection Wellington members of the Ornithological Society are asked if they would kindly help by collecting any freshly dead birds that they may find and forwarding them either to R. A. Fordham at the Zoology Department, V.U.W., or to Mr. F. C. Kinsky at the Dominion Museum. A brief note with the specimen, stating the locality and date of finding would be very helpful.

★

BACK NUMBERS OF "NOTORNIS"

Members are reminded that back numbers of *Notornis* and the earlier *N.Z. Bird Notes* are obtainable from the Society. Enquiries about costs and the parts still held in stock should be made to:— Mrs. Hetty McKenzie, Box 45, Clevedon, Auckland.

Other publications available are: *The Takaha* (5/-); *Identification of Albatrosses* (1/-); *Reports and Bulletins, 1939-1942*, with Index, (12/-), Index Alone 1/6. These precede Vol. I of *N.Z. Bird Notes* and record the first three years of the Society's work.

As there is a steady demand for back numbers of *Notornis* and especially for the earlier *N.Z. Bird Notes* (1943-1950), members are asked to offer to the Society, for gift or sale, past numbers which they no longer need.

★

BANDING REPORTS FOR SALE

The full and detailed Ninth, Tenth and Eleventh Annual Reports of the Banding Committee for the years ending 31/3/59 (38 pages), 31/3/60 (42 pages) and 31/3/61 (37 pages) are available at 5/6d. each and may be obtained from Mrs Hetty McKenzie, Box 45, Clevedon, Auckland.

★

NEW MEMBERS up to 26/7/61

Annabell, Mrs. L., Lower Beach Road, Haumoana, H.B.
Chamberlin, B. H., Mataitai, R.D. 2, Clevedon
Clifford, B., 72 Eastbourne Street, Murrays Bay, Auckland
Dene, Miss Eunice M., 23 Austin Street, Wellington
Finer, F. J., 22 Clifford Street, Hawera
Goulding, Miss J. H., 12 Grand View Road, Remuera, Auckland, S.E. 2
Hawley, P. J., Box 62, Whakatane
Howard, Ian, 32 Huia Road, Point Chevalier, Auckland, W. 3
Hutchison, Miss Jan, 25 Aitken Street, Wellington, N. 1
Jackson, Raymond, 319 Whaka Road, Rotorua
Jeffery, Ron., 36 Summit Drive, Mt. Albert, Auckland
Lawson, E. R., 86 Park Avenue, Papatoetoe
Macmillan, D. G., "Omana," Wanstead, Waipukurau
Panckhurst, D. J., 11 Byron Street, Greymouth
Ross, James J., 177 Portage Road, New Lynn, Auckland, S.W. 3
Shaw, Miss Norma, C/o Mr. D. J. Shaw, Box 3, Clevedon
Sumpter, P. C., Box 64, Papakura
Wagener, Allan, Pukenui, R.D. 4, Kaitiaki
Ward, Mrs. E. E., "Te Araroa," Mountain View Road, R.D. 4, Timaru