FALLA, R. A.; SIBSON, R. B.; TURBOTT, E. G. 1979. The New Guide to the Birds of New Zealand. London: Collins.

FLEMING C. A. 1982. George Edward Lodge The Unpublished New Zealand Bird Paintings. Wellington: Nova Pacífica.

KINSKY, F. C. (Convener). 1970. Annotated Checklist of the Birds of New Zealand, including the Birds of the Ross Dependency. Wellington: Reed.

MORSE, P. M. 1981. Wildlife values and wildlife conservation of Buller and North Westland. Fauna Survey Report No. 29. Wildlife Service, Department of Internal Affairs, Wellington.

RIPLEY, S. D. 1977. Rails of the World: A Monograph of the Family Rallidae. Toronto: M. F. Feheley. STEAD, E. F. 1927. The native and introduced birds of Canterbury. Pages 204-225 in Speight, R.; Wall, A.; Laing, R. M. (eds.). The Natural History of Canterbury. Christchurch: Philosophical Institute of Canterbury.

STIDOLPH, R. H. D. 1926. Bird-life around Wellington, N.Z. Emu. 25: 204-207. STIDOLPH, R. H. D. 1931. Birds of the Wairarapa plains. Emu 31: 136-140. WARHAM, J. 1961. The birds of Raine Island, Pandora Cay and Murray Island Sandbank, North

Queensland. Emu 61: 77-93.
WINTERBOURN, M. 1970. The New Zealand species of *Potamopyrgus* (Gastropoda: Hydrobiidae). Malacologia 10: 283-321.

GRAEME ELLIOTT, Zoology Department, Victoria University of Wellington, Box 600, Wellington



SHORT NOTE

Grev-backed Storm Petrel at Doubtful Sound, Fiordland

On the night of 2/11/88 a Grey-backed Storm Petrel (Garrodia nereis) flew aboard RV Munida moored in Deep Cove, Doubtful Sound. The bird was presumably attracted by the lit-up vessel. The weather was overcast with rain. Moderate westerly winds had predominated for some days. The bird was given to me next morning by the crew, A. Heineman and P. Meredith. It was released on 4/11/88.

The lack of plankton-feeding birds plus very low surface salinity at the time suggested low available zooplankton. It is therefore unlikely that the storm petrel had been feeding in Doubtful Sound. Potential food abundance in Doubtful Sound would have been even lower in August, when Cooper (1980) recovered two G. nereis under similar circumstances. Breeding grounds are not known in the Fiordland region.

Similar recoveries were reported by Wright (1973), Esler (1978), Cooper (1980), and Morrison (1981, 1983). All were from the southern South Island remote from food sources. Seventeen birds have been recovered by beach patrols since 1953, mostly on the North Island west coast and all between May and December.

LITERATURE CITED

COOPER, W. 1980. Grey-backed Storm Petrels at Doubtful Sound. Notornis 27:396. ESLER, L. 1978. Grey-backed Storm Petrel in Classified Summarised Notes. Notornis 25: 332-349. MORRISON, K. 1981. Grey-backed Storm Petrel in Classified Summarised Notes. Notornis 28: 57-85. MORRISON K. 1983. Grey-backed Storm Petrel on Routeburn Track. Notornis 30: 108. WRIGHT, A. 1973. Grey-backed Storm Petrel in Classified Summarised Notes. Notornis 20: 346-376.

DAVID HAWKE, Chemistry Department, University of Otago, Dunedin