

- LATHAM, P. C. M. 1981. Another Arctic Tern at the Tarawera River mouth. *Notornis* 28: 213-214.
- MARPLES, G.; MARPLES, A. 1934. Sea Terns or Sea Swallows, their habits, language, arrival and departure. London: Country Life.
- MAYR, E. 1945. Birds of the Southwest Pacific. New York: MacMillan.
- RICHARDSON, R. A. 1953. A distinction in flight between Arctic and Common Terns. *British Birds* 46: 411-412.
- SERVENTY, D. L.; SERVENTY V. N.; WARHAM, J. 1971. The handbook of Australian sea-birds. Sydney: A. H. & A. W. Reed.
- SERVENTY, D. L.; WHITTELLI, H. M. 1976. Birds of Western Australia. 5th ed. Perth: University of Western Australia Press.
- SIMPSON, K. 1972. Birds in Bass Strait. Sydney: BHP & A. H. & A. W. Reed.
- SLATER, P. 1970. A field guide to Australian birds. Non-passerines. Adelaide: Rigby.
- SMYTHIES, B. E. 1981. The birds of Borneo. 3rd ed. Kota Kinabalu and Kluang Lumpur: Sabah Society and Malayan Nature Society.
- SONOBE, K.; ROBINSON, J. W. (eds). 1982. A field guide to the birds of Japan. Tokyo: Wild Bird Society of Japan.
- VANDE WEGHE, J-P. 1966. La Sterne pierregarin (*Sterna hirundo*) et la Sterne arctique (*Sterna paradisaea*). Identification et passage en Belgique. Aves 3: 1-5, plates 1-4 (English translation of identification section, plus plates and editorial notes on separation of Forster's Tern (*Sterna forsteri*), published in 1970 in *California Birds* 1: 33-36).
- VAURIE, C. 1965. The birds of the Palearctic fauna. Non-passeriformes. London: H. F. & H. G. Witherby.
- WADE, P. (ed.) 1975. Every Australian bird illustrated. Adelaide: Rigby.
- WATLING, R. 1982. Birds of Fiji, Tonga and Samoa. Wellington: Millwood.
- WATSON, G. E. 1975. Birds of the Antarctic and Sub-Antarctic. Washington: American Geophysical Union.

P. C. M. LATHAM, *c/o Papamoa Beach Post Office, via Te Puke, Bay of Plenty*



SHORT NOTE

Second New Zealand record of the Bird of Providence

On 22 September 1984 I found a beach-wrecked Bird of Providence (*Pterodroma solandri*) on the west coast of Northland between Maunganui Bluff and Omamari. The bird was identified by D. E. Crockett and subsequently the identification was confirmed by M. J. Imber. This is only the second record of this species in New Zealand, the first being that found by Falla on Muriwai Beach in January 1921. The specimen from Northland is now held at the National Museum.

Description: Back grey-brown. Head dark brown. Uppertail brown. Face and forehead freckled. Belly white in centre, grey on sides — each feather mostly white with a 10 mm grey tip on the end. Small pale tip right on the end of some feathers. Underside of tail grey-brown.

Underwing: primaries grey-brown on outer vane and tips, paler to white on basal half of inner vane.

Feet: black — paler at base of toes *Measurements* (mm)

Legs: paler — brown

Bill: black — large

Culmen: 33.22

Bill width: 14.40

Bill depth: 16.08

Tarsus: right 42.14, left 41.36

Mid-toe & claw: right 59.0, left 59.10

Wing: 310

Length: 420

PATRICK MILLER, 29 Murdoch Crescent, Whangarei