


# NOTORNIS

Ornithology of the Southern Pacific



ISSN 0029-4470

Volume 68 2021  
Journal of the Ornithological Society of New Zealand



## Volume 68, Part 1, March 2021

## CONTENTS

**Papers**

|                                                                                                                          |                                                                                                                |    |
|--------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------|----|
| Nocturnal activity of the western weka ( <i>Gallirallus australis australis</i> ) in an open environment                 | Lamb, S.D.; Taylor, H.R.; Powlesland, R.                                                                       | 1  |
| Clutch sizes and hatching success of Canada geese nesting in Canterbury, New Zealand                                     | Adams, J.S.; Williams, M.                                                                                      | 13 |
| The food and foraging of <i>Collocalia</i> and <i>Aerodramus</i> swiftlets: a review                                     | Tarburton, M.K.; Collins, C.T.                                                                                 | 26 |
| Erect-crested penguins on the Bounty Islands: population size and trends determined from ground counts and drone surveys | Mattern, T.; Rexer-Huber, K.; Parker, G.; Amey, J.; Green, C-P.; Tennyson, A.J.D.; Sagar, P.M.; Thompson, D.R. | 37 |
| Dominance interactions among New Zealand albatrosses and petrels at ecotourist boats                                     | Martin, P.R.; Briskie, J.V.                                                                                    | 51 |

**Short notes**

|                                                                                                                                                                             |                                                  |    |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------|----|
| Individual long-distance migrant <i>Chrysococcyx</i> cuckoos repeat carbon and nitrogen stable isotope ratios after moulting in non-breeding range on successive migrations | Holdaway, R.N.; Sagar, P.M.; Briskie, J.V.       | 65 |
| Second record of Buller's shearwater breeding outside of the Poor Knights Islands                                                                                           | Crowe, P.; Willis, K.; Butler, D.; Whitehead, E. | 72 |
| Vessel survey observations confirm wintering dispersion of northern giant-petrel ( <i>Macronectes halli</i> ) juveniles in southern-central Perú; what is their origin?     | Quiñones, J.; Romero, C.; Zavalaga, C.           | 76 |
| Longevity records for Chatham Island pigeon ( <i>Hemiphaga chathamensis</i> ) and New Zealand pigeon ( <i>H. novaeseelandiae</i> )                                          | Clout, M.N.; Robertson, H.A.                     | 86 |
| First description of the egg of the enigmatic pied cuckoo-dove, with a literature review on the breeding biology of the congeneric crested and great cuckoo-dove            | Marcuk, V.; De Boer, D.                          | 89 |

## Volume 68, Part 2, June 2021

## CONTENTS

**Papers**

- |                                                                                                                          |                                                                                                              |            |
|--------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------|------------|
| Increasing urban abundance of tūī ( <i>Prosthemadera novaeseelandiae</i> ) by pest mammal control in surrounding forests | <b>Fitzgerald, N.; Innes, J.; Watts, C.; Thornburrow, D.; Bartlam, S.; Collins, K.; Byers, D.; Burns, B.</b> | <b>93</b>  |
| Estimating the distribution, population status, and trends of New Zealand scaup ( <i>Aythya novaeseelandiae</i> )        | <b>Greene, B.S.</b>                                                                                          | <b>108</b> |
| Breeding ecology of a translocated population of great spotted kiwi ( <i>Apteryx haastii</i> )                           | <b>Toy, R.; Toy, S.</b>                                                                                      | <b>131</b> |
| Measuring conservation status in New Zealand birds: re-evaluating banded dotterel and black-fronted tern as case studies | <b>Craig, J.L.; Mitchell, N.D.</b>                                                                           | <b>147</b> |

**Short notes**

- |                                                                                                                                                                                     |                                                                    |            |
|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------|------------|
| Common starling ( <i>Sturnus vulgaris</i> ) laying dates, 1970–2019, have not changed in New Zealand, in contrast to those in Denmark                                               | <b>Flux, J.E.C.; Flux, M.M.</b>                                    | <b>161</b> |
| Northward expansion of the non-breeding range of Otago shag ( <i>Leucocarbo chalconotus</i> ) along the Canterbury coast towards Banks Peninsula, eastern South Island, New Zealand | <b>Crossland, A.C.</b>                                             | <b>166</b> |
| Little shearwaters ( <i>Puffinus assimilis haurakiensis</i> ) as prey for morepork ( <i>Ninox novaeseelandiae</i> )                                                                 | <b>Whitehead, E.A.</b>                                             | <b>170</b> |
| Changes in behaviour of great spotted kiwi ( <i>Apteryx haastii</i> ) following handling                                                                                            | <b>Toy, R.; Toy, S.</b>                                            | <b>173</b> |
| Further evidence in support of grey-backed storm petrels ( <i>Garrodia nereis</i> ) breeding in Fiordland                                                                           | <b>Miskelly, C.M.; Stahl, J.C.; Tennyson, A.J.D.; Bishop, C.R.</b> | <b>177</b> |

**Letters to the Editor****182**

## Volume 68, Part 3, September 2021

## CONTENTS

**Papers**

- |                                                                                                                                           |                                                                                                              |            |
|-------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------|------------|
| A survey of Fiordland crested penguins ( <i>Eudyptes pachyrhynchus</i> ): northeast Stewart Island/Rakiura, New Zealand, September 2019   | <b>Long, R.; Litchwark, S.</b>                                                                               | <b>183</b> |
| Field sexing techniques for Fiordland crested penguins (tawaki; <i>Eudyptes pachyrhynchus</i> )                                           | <b>White, J.; Mattern, T.; Ellenberg, U.; Garcia-Borboroglu, P.; Houston, D.M.; Seddon, P.J.; Mays, H.L.</b> | <b>188</b> |
| Breeding petrels of northern and central Fiordland, with a summary of petrel populations for the Fiordland region                         | <b>Miskelly, C.M.; Bishop, C.R.; Tennyson, A.J.D.</b>                                                        | <b>194</b> |
| Garden birds at Rangiora, Christchurch, and Kaikōura, South Island, New Zealand: results from banding 1961–2016                           | <b>Rowe, L.K.</b>                                                                                            | <b>208</b> |
| Widespread ground-nesting in a large population of feral rock pigeons ( <i>Columba livia</i> ) in a predator-free and urban native forest | <b>Briskie, J.V.; Shorey, L.</b>                                                                             | <b>224</b> |

**Short notes**

- |                                                                                                                                                                          |                        |            |
|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------|------------|
| Radiocarbon ages for kakapo ( <i>Strigops habroptilus</i> ) (Strigopidae: Strigopinae) from the Pyramid Valley lake bed deposit, north-eastern South Island, New Zealand | <b>Holdaway, R.N.</b>  | <b>234</b> |
| Decline in common mynas ( <i>Acridotheres tristis</i> ) between 1971 and 2019 in central Auckland, New Zealand                                                           | <b>Beauchamp, A.J.</b> | <b>239</b> |

## Volume 68, Part 4, December 2021

## CONTENTS

**Papers**

- |                                                                                                                                                                                        |                                                                                                       |            |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------|------------|
| Diet of the Floreana mockingbird ( <i>Mimus trifasciatus</i> ) during the dry season on Champion and Gardner Islets, Galápagos Islands, Ecuador                                        | <b>Wittmer-Naranjo, C.; Reyes, E.M.R.; Jácome, H.E.T.; Rueda, D.; Sevilla, C.; Ortiz-Catedral, L.</b> | <b>245</b> |
| Vagrant and extra-limital bird records accepted by the Birds New Zealand Records Appraisal Committee 2019–2020                                                                         | <b>Miskelly, C.M.; Crossland, A.C.; Saville, I.; Southey, I.; Tennyson, A.J.D.; Bell, E.A.</b>        | <b>253</b> |
| Seasonal survey of waterfowl (Anatidae), shags (Phalacrocoracidae) and fernbird ( <i>Bowdleria punctata</i> ) at Te Nohoaka o Tukiauau/Sinclair Wetlands, Otago: July 2015 – July 2018 | <b>Thompson, M.P.; Mckinlay, B.</b>                                                                   | <b>266</b> |

**Short notes**

- |                                                                                                                                                                                                    |                                                        |            |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------|------------|
| New Zealand king shag ( <i>Leucocarbo carunculatus</i> ) with deformed primary feathers                                                                                                            | <b>Melville, D.S.; Schuckard, R.</b>                   | <b>274</b> |
| Two new radiocarbon ages for Haast's eagle ( <i>Hieraaetus moorei</i> ) (Aves: Accipitridae) and comments on the eagle's past distribution and possible survival into the 19 <sup>th</sup> century | <b>Holdaway, R.N.</b>                                  | <b>278</b> |
| A previously unpublished first record of the moa by French naval surgeon/botanist P.A. Lesson                                                                                                      | <b>Lee, M.</b>                                         | <b>283</b> |
| Did Molly Falla observe an instance of active tactical deception in the kea ( <i>Nestor notabilis</i> )?                                                                                           | <b>Woodley of Menie, M.A.; Peñaherrera-Aguirre, M.</b> | <b>290</b> |
| Mind the gap: potential implications of the chronology of the South Island adzebill ( <i>Aptornis defossor</i> ) (Aves: Aptornithidae) at Pyramid Valley, North Canterbury, New Zealand            | <b>Holdaway, R.N.</b>                                  | <b>294</b> |

# INDEX OF AUTHORS AND SPECIES

## Volume 68

- Acridotheres tristis*, see myna, Indian
- ADAMS, J.S.; WILLIAMS, M. Clutch sizes and hatching success of Canada geese nesting in Canterbury, New Zealand, 13-25
- adzebill, North Island, 295, 296  
South Island, 236, 279, 294-298
- Aerodramus bartschi*, see swiftlet, Mariana
- elaphrus*, see swiftlet, Seychelles
- fuciphagus*, see swiftlet, white-nest
- germani*, see swiftlet, Germain's
- hirundinacea*, see swiftlet, mountain
- infuscatus*, see swiftlet, Halmahera
- maximus*, see swiftlet, black-nest
- papuensis*, see swiftlet, three-toed
- pelewensis*, see swiftlet, Palau
- salanganus*, see swiftlet, mossy-nest
- sawtelli*, see swiftlet, Atiu
- spodiopygius*, see swiftlet, white-rumped
- terraereginae*, see swiftlet, Australian
- unicolor*, see swiftlet, Indian
- vanikorensis*, see swiftlet, uniform
- albatross, Antipodean, 52, 57  
Buller's, see mollymawk, Buller's  
Gibson's, 51-53, 55-60, 62  
northern royal, 52, 82  
Salvin's, see mollymawk, Salvin's  
southern royal, 51-53, 55-60, 62  
spp., 51-64, 82  
wandering, 52, 57  
waved, 82
- Alisterus scapularis*, see parrot, Australian king
- AMEY, J., see MATTERN, T.; REXER-HUBER, K.; PARKER, G.; AMEY, J.; GREEN, C.-P.; TENNYSON, A.J.D.; SAGAR, P.M.; THOMPSON, D.R.
- Anas castanea*, see teal, chestnut
- chlorotis*, see teal, brown
- clypeata*, see shoveler, northern
- gracilis*, see teal, grey
- platyrhynchos* x *superciliosa*, see duck, mallard x grey hybrid
- rhynchotis*, see shoveler, Australasian, see also shoveler, New Zealand
- Anatidae, 108-130, 266-273
- Anhinga melanogaster*, see darter
- Anomalopteryx didiformis*, see moa
- Anous minutus*, see noddy, black
- Anser anser*, see goose, greylag
- Anthornis melanura melanura*, see bellbird
- Apodidae, 28
- Apodiform spp., 26
- Apteryx australis australis*, see kiwi, South Island brown, or tokoeka, Haast
- australis* 'Haast', see kiwi, South Island brown, or tokoeka, Haast
- haastii*, see kiwi, great spotted
- mantelli*, see kiwi, North Island brown
- rowi*, see kiwi, Okarito brown or rowi spp., see kiwi spp.
- Aptornis defossor*, see adzebill, South Island
- otidiformis*, see adzebill, North Island
- Aptornithidae, 294-298
- Apus pacificus*, see swift, fork-tailed spp., 31
- Ardea modesta*, see heron, white
- Ardenna creatopus*, see shearwater, pink-footed
- gravis*, see shearwater, great
- grisea*, see shearwater, sooty
- spp., see shearwater, spp
- Artamus personatus*, see woodswallow, masked
- Athene noctua*, see owl, little
- Aythya affinis*, see scaup, lesser
- americana*, see redhead
- australis*, see hardhead or duck, white-eyed
- baeri*, see pochard, Baer's
- collaris*, see duck, ring-necked
- farina*, see pochard, common
- fuligula*, see duck, tufted
- innotata*, see pochard, Madagascar
- marila*, see scaup, greater
- novaeseelandiae*, see scaup, New Zealand
- nyroca*, see pochard, white-eyed
- spp., 109, 110
- vallisineria*, see canvasback
- BARTLAM, S., see FITZGERALD, N.; INNES, J.; WATTS, C.; THORNBURROW, D.; BARTLAM, S.; COLLINS, K.; BYERS, D.; BURNS, B.
- BEAUCHAMP, A.J. Decline in common mynas (*Acridotheres tristis*) between 1971 and 2019 in central Auckland, New Zealand, 239-244
- BELL, E.A., see MISKELLY, C.M.; CROSSLAND, A.C.; SAVILLE, I.; SOUTHEY, I.; TENNYSON, A.J.D.; BELL, E.A.
- bellbird, 209, 210, 218, 220, 225
- bird, diving spp., 63  
forest spp., 143  
introduced spp., 224-233  
migrant spp., 69  
native spp., 224-233  
non-passerine spp., 294  
of prey, 275

- wading spp., 272  
 BISHOP, C.R., see MISKELLY, C.M.; STAHL, J.C.; TENNYSON, A.J.D.; BISHOP, C.R.  
 BISHOP, C.R., see MISKELLY, C.M.; BISHOP, C.R.; TENNYSON, A.J.D.  
 bittern, Australasian, 123, 266, 272  
 blackbird, 209, 211  
   Eurasian, 209, 211, 213, 219-221, 225, 230  
 boobook, Southern, 171, 172  
 booby, blue-footed, 250  
   brown, 256  
   Eastern Pacific red-footed, 256  
   Nazca, 248, 250  
   red-footed, 256  
   South-west Pacific red-footed, 256  
   spp., 63, 248  
*Botaurus poiciloptilus*, see bittern, Australasian  
*Bowdleria punctata*, see fernbird  
*Branta canadensis*, see goose, Canada  
   *canadensis maxima*, see goose, Canada  
 BRISKIE, J.V.; SHOREY, L. Widespread ground-nesting in a large population of feral rock pigeons (*Columba livia*) in a predator-free and urban native forest, 224-233  
 BRISKIE, J.V., see MARTIN, P.R.; BRISKIE, J.V.  
 BRISKIE, J.V., see HOLDAWAY, R.N.; SAGAR, P.M.; BRISKIE, J.V.  
 BURNS, B., see FITZGERALD, N.; INNES, J.; WATTS, C.; THORNBURROW, D.; BARTLAM, S.; COLLINS, K.; BYERS, D.; BURNS, B.  
 BUTLER, D., see CROWE, P.; WILLIS, K.; BUTLER, D.; WHITEHEAD, E.  
 BYERS, D., see FITZGERALD, N.; INNES, J.; WATTS, C.; THORNBURROW, D.; BARTLAM, S.; COLLINS, K.; BYERS, D.; BURNS, B.  
  
*Cacatua goffini*, see cockatoo, Goffin's  
*Cacomantis flabelliformis*, see cuckoo, fan-tailed  
*Calidris alba*, see sanderling  
   *minutus*, see stint, little  
   *tenuirostris*, see knot, great  
*Callaeas cinerea*, see kōkako, South Island  
   *wilsoni*, see kōkako, North Island  
*Camarhynchus* spp., see finch, Darwin's spp.  
 canvasback, 108  
*Carduelis carduelis britannica*, see goldfinch, European  
   *chloris*, see greenfinch, European  
   *flammea*, see redpoll, common  
*Catharacta antarctica*, see skua, subantarctic  
   *maccormicki*, see skua, South Polar  
   spp., see skua, spp.  
*Cereopsis novaehollandiae*, see goose, Cape Barren  
*Certhidea* spp., see finch, Darwin's spp.  
*Chaetura pelagica*, see swift, chimney  
 chaffinch, 211, 212, 218, 220, 221, 272  
*Charadrius bicinctus*, see dotterel, banded  
   *leschenaultii*, see plover, greater sand  
   *melodus*, see plover, piping  
   *mongolus*, see plover, lesser sand  
   *rubricollis*, see dotterel, hooded  
   *wilsonia*, see plover, Wilson's  
*Chenonetta finschi*, see duck, Finsch's  
 chicken, domestic, 250  
*Chlidonias albostratus*, see tern, black-fronted  
   *hybridus*, see tern, whiskered  
   *leucopterus*, see tern, white-winged black  
*Chrysococcyx*, Australian spp., 66  
   *lucidus lucidus*, see cuckoo, shining  
*Circus approximans*, see harrier, swamp  
   *eylesi*, see hawk, Eyles', see also *Circus teauteensis*  
   spp., see harrier spp.  
   *teauteensis*, see harrier, extinct  
 CLOUT, M.N.; ROBERTSON, H.A. Longevity records for Chatham Island pigeon (*Hemiphaga chathamensis*) and New Zealand pigeon (*H. novaeseelandiae*), 86-88  
*Cnemidornis calcitrans*, see goose, South Island  
 cockatoo, Goffin's, 292  
*Colinus virginianus*, see quail, bobwhite  
 COLLINS, C.T., see TARBURTON, M.K.; COLLINS, C.T.  
 COLLINS, K., see FITZGERALD, N.; INNES, J.; WATTS, C.; THORNBURROW, D.; BARTLAM, S.; COLLINS, K.; BYERS, D.; BURNS, B.  
*Collocalia esculenta*, see swiftlet, glossy  
   spp., 26-36  
   *troglodytes*, see swiftlet, pygmy  
*Columba janthina janthina*, see pigeon, Japanese wood  
   *livia*, see pigeon, rock  
   *palumbus*, see woodpigeon  
 Columbidae spp., 224-233  
 coot, Australian, 270  
   extinct New Zealand, 280  
*Coracina novaehollandiae*, see cuckoo-shrike, black-faced  
*Coracopsis vasa*, see parrot, greater vasa  
*Corvus albicollis*, see raven, white-necked  
   *corax*, see raven, common  
 CRAIG, J.L.; MITCHELL, N.D. Measuring conservation status in New Zealand birds: re-evaluating banded dotterel and black-fronted tern as case studies, 147-160  
 crane, marsh, 260, 270, 272  
   spotless, 270, 272  
   spp., 266, 272  
 crane spp., 295  
 CROSSLAND, A.C. Northward expansion of the non-breeding range of Otago shag (*Leucocarbo chalconotus*) along the Canterbury coast towards Banks Peninsula, eastern South Island, New Zealand, 166-169  
 CROSSLAND, A.C., see MISKELLY, C.M.; CROSSLAND, A.C.; SAVILLE, I.; SOUTHEY, I.; TENNYSON, A.J.D.; BELL, E.A.  
 CROWE, P.; WILLIS, K.; BUTLER, D.; WHITEHEAD,



- E. Second record of Buller's shearwater breeding outside of the Poor Knights Islands, 72-75
- cuckoo, fan-tailed, 262
- pallid, 259
- shining, 65-71, 225
- cuckoo-dove, crested, 89-91
- great, 89-91
- pied, 89-91
- cuckoo-shrike, black-faced, 259, 262
- Cuculus pallidus*, see cuckoo, pallid
- Cyanoramphus novaeseelandiae*, see parakeet, red-crowned
- sp., see kākāriki
- Cygnus atratus*, see swan, black
- Cypseloidinae, see swift, Neotropical spp.
- dabchick, New Zealand, 123, 259
- Daption capense*, see petrel, Cape
- darter, 261
- DE BOER, D., see MARCUK, V.; DE BOER, D.
- Dendrocygna eytoni*, see duck, plumed whistling
- Dinornis novaeseelandiae*, see moa, giant
- robustus*, see moa, South Island giant
- Dinornithiformes, 235, 278-282, 288-289, 294
- Diomedea antipodensis gibsoni*, see albatross, Gibson's
- wandering
- epomophora*, see albatross, southern royal
- epomophora sanfordi*, see *Diomedea sanfordi*
- exulans*, see albatross, wandering
- sanfordi*, see albatross, northern royal
- Diomededidae, 51-64
- dodo, 232
- dotterel, banded, 147-160
- black-fronted, 148, 260
- hooded, 182
- dove, barbery, 261
- bar-shouldered, 87
- collared, 87
- eastern turtle, 231
- Galápagos, 248, 250
- mourning, 87
- rose-crowned fruit, 253, 258-259, 268
- spp., 86-88
- white-winged, 87
- duck, ferrugineous, see pochard, white-eyed
- Finsch's, 236
- grey, 268, 271
- mallard, 268-271
- mallard x grey hybrid, 268, 271
- paradise, 266, 269-271
- plumed whistling, 254
- ring-necked, 108
- tufted, 108
- white-eyed, 108
- dunnoek, 208, 211, 212, 219-221, 225, 272
- eagle, African crowned, 280
- forest spp., 280
- Haast's, 236, 278-282, 294
- harpy, 280
- egret, little, 256
- Egretta garzetta*, see egret, little
- novaehollandiae*, see heron, white-faced
- sacra*, see heron, reef
- ELLENBERG, U., see WHITE, J.; MATTERN, T.; ELLENBERG, U.; GARCIA-BORBOROGLU, P.; HOUSTON, D.M.; SEDDON, P.J.; MAYS, H.L.
- Elseyornis melanops*, see dotterel, black-fronted
- Emberiza citrinella*, see yellowhammer
- Emeus crassus*, see moa
- emu, 283
- Eudyptes chrysocome*, see penguin, western rockhopper
- chrysolophus*, see penguin, macaroni
- filholi*, see penguin, eastern rockhopper
- moseleyi*, see penguin, northern rockhopper
- pachyrhynchus*, see penguin, Fiordland crested
- robustus*, see penguin, Snares crested
- schlegeli*, see penguin, royal
- slateri*, see penguin, erect-crested
- spp., see penguin, crested spp.
- Eudyptula minor*, see penguin, little
- Euryapteryx curtus*, see moa
- Falco centroides*, see kestrel, nankeen
- novaeseelandiae*, see falcon, New Zealand
- falcon, New Zealand, 7, 280
- fantail, New Zealand, 209-211, 225
- South Island, 209-211, 218, 220, 225
- fernbird, 261
- South Island, 261, 266-273
- finch, Darwin's spp., 246
- FITZGERALD, N.; INNES, J.; WATTS, C.; THORNBURROW, D.; BARTLAM, S.; COLLINS, K.; BYERS, D.; BURNS, B. Increasing urban abundance of tūi (*Prothemadera novaeseelandiae*) by pest mammal control in surrounding forests, 93-107
- FLUX, J.E.C.; FLUX, M.M. Common starling (*Sturnus vulgaris*) laying dates, 1970-2019, have not changed in New Zealand, in contrast to those in Denmark, 161-165
- FLUX, M.M., see FLUX, J.E.C.; FLUX, M.M.
- flycatcher spp., 28
- fowl, domestic, 286
- Fratercula arctica*, see puffin, Atlantic
- Fregata* sp., see frigatebird sp.
- Fregatta maoriana*, see storm petrel, New Zealand
- frigatebird sp., 256
- Fringilla coelebs*, see chaffinch
- Fulica atra*, see coot, Australian
- prisca*, see coot, extinct New Zealand
- Gallirallus australis*, see weka
- australis australis*, see weka, western
- philippensis*, see rail banded
- Gallus gallus*, see junglefowl, red

- gamebird spp., 13-25  
 gannet, Australasian, 256, 260  
 GARCIA-BORBOROGLU, P., see WHITE, J.;  
 MATTERN, T.; ELLENBERG, U.; GARCIA-  
 BORBOROGLU, P.; HOUSTON, D.M.;  
 SEDDON, P.J.; MAYS, H.L.  
*Garrodia nereis*, see storm petrel, grey-backed  
*Gelochelidon nilotica microtarsa*, see tern, gull-billed  
 (Australian subsp.)  
*Geopelia humeralis*, see dove, bar-shouldered  
*Geospiza*, see finch, Darwin's spp.  
*Gerygone igata*, see warbler, grey  
 goldfinch, 211, 212, 218, 220  
 goose, Canada, 13-25, 259, 269-271  
   Cape Barren, 259  
   greylag, 269  
   South Island, 236, 279, 296  
 grebe, Australasian crested, 123  
   Australasian little, 260  
   hoary-headed, 254  
 GREEN, C-P., see MATTERN, T.; REXER-HUBER,  
 K.; PARKER, G.; AMEY, J.; GREEN,  
 C-P.; TENNYSON, A.J.D.; SAGAR, P.M.;  
 THOMPSON, D.R.  
 GREENE, B.S. Estimating the distribution, population  
 status, and trends of New Zealand scaup (*Aythya*  
*novaeseelandiae*), 108-130  
 greenfinch, 211, 212, 219, 220, 225  
 greenshank, common, 257, 262  
 Gruiformes, 295  
 gull, black-backed, 18, 19, 41  
   laughing, 258, 263  
   red-billed, 208  
   spp., 63, 275  
*Gygis alba*, see tern, white  
  
*Haliaeetus albicilla*, see sea eagle, European white-tailed  
*leucogaster*, see sea eagle, white-bellied  
 hardhead, 108  
*Harpia harpyja*, see eagle, harpy  
 harrier, Australasian, see harrier, swamp  
   extinct, 234, 280, see also harrier, Eyles'  
   Eyles', 7, 236, 280  
   spp., 109  
   swamp, 7, 93, 98, 100, 102, 109, 172, 224, 231,  
   280  
 hawk, Eyles', see harrier, Eyles'  
   sp., 280  
*Hemiphaga chathamensis*, see pigeon, Chatham Island,  
 see also paraea  
*novaeseelandiae*, see pigeon, New Zealand, see  
 also kererū  
 heron, Pacific, 261  
   reef, 261  
   spp., 261, 272  
   white, 263  
   white-faced, 261  
*Hieraaetus moorei*, see eagle, Haast's  
  
 hihi, see stitchbird  
*Hirundapus caudacutus*, see needletail, white-throated  
*Hirundo neoxena*, see swallow, welcome  
 hoiho, see penguin, yellow-eyed  
 HOLDAWAY, R.N. Letter to Editor (re *Thinornis rossi*),  
 182  
 HOLDAWAY, R.N. Radiocarbon ages for kakapo  
 (*Strigops habroptilus*) (Strigopidae: Strigopinae)  
 from the Pyramid Valley lake bed deposit,  
 north-eastern South Island, New Zealand, 234-  
 238  
 HOLDAWAY, R.N. Two new radiocarbon ages for  
 Haast's eagle (*Hieraaetus moorei*) (Aves:  
 Accipitridae) and comments on the eagle's past  
 distribution and possible survival into the 19<sup>th</sup>  
 century, 278-282  
 HOLDAWAY, R.N. Mind the gap: potential implications  
 of the chronology of the South Island adzebill  
 (*Aptornis defossor*) (Aves: Aptornithidae)  
 at Pyramid Valley, North Canterbury, New  
 Zealand, 294-298  
 HOLDAWAY, R.N.; SAGAR, P.M.; BRISKIE, J.V.  
 Individual long-distance migrant *Chrysococcyx*  
 cuckoos repeat carbon and nitrogen stable  
 isotope ratios after moulting in non-breeding  
 range on successive migrations, 65-71  
 honeyeater spp., 93-107  
 HOUSTON, D.M., see WHITE, J.; MATTERN, T.;  
 ELLENBERG, U.; GARCIA-BORBOROGLU, P.;  
 HOUSTON, D.M.; SEDDON, P.J.; MAYS, H.L.  
*Hydrobates pelagicus*, see storm petrel, European  
*Hydrochous gigas*, see swift, waterfall  
*Hydroprogne caspia*, see tern, Caspian  
  
 ibis, glossy, 256  
   white, 261  
 INNES, J., see FITZGERALD, N.; INNES, J.; WATTS,  
 C.; THORNBURROW, D.; BARTLAM, S.;  
 COLLINS, K.; BYERS, D.; BURNS, B.  
  
 JÁCOME, H.E.T., see WITTMER-NARANJO, C.;  
 REYES, E.M.R.; JÁCOME, H.E.T.; RUEDA, D.;  
 SEVILLA, C.; ORTIZ-CATEDRAL, L.  
 jaeger spp., 62  
 junglefowl, red, 286  
  
 kāhu, see harrier, swamp  
 kākā, Norfolk Island, 292  
   South Island, 294  
 kākāpō, 180, 234-238, 279  
 kākāriki, 262  
 kea, 136, 280, 290-293  
 kererū, 86-88, 102, 225, 230, see also pigeon,  
   New Zealand  
 kestrel, nankeen, 253, 256, 262  
 kingfisher, sacred, 225, 231  
 kite, black, 256, 262  
 kiwi, 10

- great spotted, 131-146, 173-176  
 little spotted, 142  
 North Island brown, 142, 143, 287  
 South Island brown, 137, 141-143  
 spp., 131, 173, 285, 286
- knot, great, 256
- kōkako, North Island, 102  
 South Island, 262
- kororā, see penguin, little
- kuaka, see petrel, common diving
- Lalage tricolor*, see triller, white-winged
- LAMB, S.D.; TAYLOR, H.R.; POWLESLAND, R.  
 Nocturnal activity of the western weka  
 (*Gallirallus australis australis*) in an open  
 environment, 1-12
- Laridae, 63
- Larus dominicanus*, see gull, black-backed  
*novaehollandiae scopulinus*, see gull, red-billed
- LEE, M. A previously unpublished first record of the  
 moa by French naval surgeon/botanist P.A.  
 Lesson, 283-289
- LETTERS: HOLDAWAY, R.N. (re *Thinornis rossi*), 182
- Leucocarbo carunculatus*, see shag, New Zealand king  
*chalconotus*, see shag, Otago, or shag, Stewart  
 Island  
*ranfurlyi*, see shag, Bounty Island  
*stewarti*, see shag, Foveaux
- Leucophaeus atricilla*, see gull, laughing
- Limicola falcinellus*, see sandpiper, broad-billed
- LITCHWARK, S., see LONG, R.; LITCHWARK, S.
- LONG, R.; LITCHWARK, S. A survey of Fiordland  
 crested penguins (*Eudyptes pachyrhynchus*):  
 northeast Stewart Island/Rakiura, New Zealand,  
 September 2019, 183-187
- Macronectes giganteus*, see petrel, southern giant  
*halli*, see petrel, northern giant  
 spp., 77
- mallard, 268-271
- MARCUK, V.; DE BOER, D. First description of the  
 egg of the enigmatic pied cuckoo-dove, with a  
 literature review on the breeding biology of the  
 congeneric crested and great cuckoo-dove, 89-91
- martin, Australian tree, 259, 262
- MARTIN, P.R.; BRISKIE, J.V. Dominance interactions  
 among New Zealand albatrosses and petrels at  
 ecotourist boats, 51-64
- MATTERN, T.; REXER-HUBER, K.; PARKER, G.;  
 AMEY, J.; GREEN, C-P.; TENNYSON, A.J.D.;  
 SAGAR, P.M.; THOMPSON, D.R. Erect-crested  
 penguins on the Bounty Islands: population size  
 and trends determined from ground counts and  
 drone surveys, 37-50
- MATTERN, T., see WHITE, J.; MATTERN, T.;  
 ELLENBERG, U.; GARCIA-BORBOROGLU, P.;  
 HOUSTON, D.M.; SEDDON, P.J.; MAYS, H.L.
- MAYS, H.L., see WHITE, J.; MATTERN, T.;
- ELLENBERG, U.; GARCIA-BORBOROGLU, P.;  
 HOUSTON, D.M.; SEDDON, P.J.; MAYS, H.L.
- MCKINLAY, B, see THOMPSON, M.P.; MCKINLAY, B.
- Megadyptes antipodes*, see penguin, yellow-eyed
- Megalapteryx didinus*, see moa, upland
- Meliphagidae, 93-107  
 Australian spp., 102
- MELVILLE, D.S.; SCHUCKARD, R. New Zealand king  
 shag (*Leucocarbo carunculatus*) with deformed  
 primary feathers, 274-277
- Milvus migrans*, see kite, black
- Mimoides graysoni*, see see mockingbird, Socorro
- Mimus macdonaldi*, see mockingbird, Española  
*melanotis*, see mockingbird, San Cristobel  
*parvulus*, see mockingbird, Galápagos  
*parvulus bauri*, see mockingbird, Galápagos  
 subsp.  
*trifasciatus*, see mockingbird, Floreana
- MISKELLY, C.M.; STAHL, J.C.; TENNYSON, A.J.D.;  
 BISHOP, C.R. Further evidence in support of  
 grey-backed storm petrels (*Garrodia nereis*)  
 breeding in Fiordland, 177-181
- MISKELLY, C.M.; BISHOP, C.R.; TENNYSON, A.J.D.  
 Breeding petrels of northern and central  
 Fiordland, with a summary of petrel populations  
 for the Fiordland region, 194-207
- MISKELLY, C.M.; CROSSLAND, A.C.; SAVILLE, I.;  
 SOUTHEY, I.; TENNYSON, A.J.D.; BELL, E.A.  
 Vagrant and extra-limital bird records accepted  
 by the Birds New Zealand Records Appraisal  
 Committee 2019-2020, 253-265
- MITCHELL, N.D., see CRAIG, J.L.; MITCHELL, N.D.
- moa, emeid spp., 280  
 giant, 284  
 South Island giant, 280  
 spp., 235, 236, 278-280, 283-289, 294, 295, 297  
 upland, 280
- mockingbird, Española, 246, 250  
 Floreana, 245-252  
 Galápagos, 246, 248  
 Galápagos subsp., 246  
 San Cristobel, 246  
 Socorro, 248
- Mohoua ochrocephala*, see yellowhead
- mohua, 262
- mollymawk, Atlantic yellow-nosed, 253, 255  
 black-browed, 52, 55  
 Buller's, 51, 52, 54, 55, 59-61, 82  
 Indian yellow-nosed, 61  
 northern Buller's, 52, 82  
 Salvin's, 38, 39, 41, 46, 47, 51-53, 55-61, 82  
 shy, 261  
 spp., 51-64  
 Tasmanian (shy), 261  
 white-capped, 51-53, 55-61, 261
- morepork, 170-172
- Morus serator*, see gannet, Australasian
- myna, common, 239-244

- needletail, white-throated, 259, 263  
*Nestor meridionalis*, see kākā  
     *notabilis*, see kea  
     *productus*, see kākā, Norfolk Island  
*Ninox novaeseelandiae*, see morepork  
     *novaeseelandiae boobook*, see boobook,  
     Southern  
 noddy, black, 260  
*Notiomystis cincta*, see stitchbird  
*Numenius phaeopus*, see whimbrel  
  
*Onychoprion fuscatus*, see tern, sooty  
 ORTIZ-CATEDRAL, L., see WITTMER-NARANJO, C.;  
     REYES, E.M.R.; JÁCOME, H.E.T.; RUEDA, D.;  
     SEVILLA, C.; ORTIZ-CATEDRAL, L.  
 ostrich, 284  
 owl, barn, 259, 262  
     laughing, 236  
     little, 2  
     spp., 171  
  
*Pachyornis australis*, see moa  
     *elephantopus*, see moa  
*Pachyptila crassirostris*, see prion, fulmar  
     *desolata*, see prion, Antarctic  
     *turtur*, see prion, fairy  
     *vittata*, see prion, broad-billed  
 parakeet, red-crowned, 261  
 pareia, 86-88, see also pigeon, Chatham Island  
 PARKER, G., see MATTERN, T.; REXER-HUBER,  
     K.; PARKER, G.; AMEY, J.; GREEN,  
     C.-P.; TENNYSON, A.J.D.; SAGAR, P.M.;  
     THOMPSON, D.R.  
 parrot, Australian king, 263  
     greater vasa, 292  
*Passer domesticus domesticus*, see sparrow, house  
 passerine spp., 208-223  
*Pelagodroma marina*, see storm petrel, white-faced  
     *marina dulciae*, see storm petrel, Australian  
     white-faced  
*Pelecanoides garnotii*, see diving petrel, Peruvian  
     *urinatrix*, see petrel, common diving  
 PEÑAHERRERA-AGUIRRE, M., see WOODLEY OF  
     MENIE, M.A.; PEÑAHERRERA-AGUIRRE, M.  
 penguin, African, 189  
     blue, see penguin, little  
     crested spp., 38, 192  
     eastern rockhopper, 38, 46  
     erect-crested, 37-50, 260  
     *Eudyptes* spp., 189  
     Fiordland crested, 38, 183-187, 188-193, 260  
     Humboldt, 83  
     little, 189, 190  
     macaroni, 190, 255, 263  
     northern rockhopper, 189, 190  
     rockhopper, 46  
     royal, 255, 263  
     Snares crested, 38  
     southern rockhopper, see penguin, western  
     rockhopper  
     spp., 63, 161, 189, 275  
     western rockhopper, 189, 190, 254, 255  
     yellow-eyed, 46, 184, 189, 260  
 petrel, Cape, 51-56, 60-62  
     collared, 253, 255, 263  
     common diving, 172, 178, 194, 195, 199-201  
     Cook's, 172  
     Fiji, 261  
     giant, 51-53, 61, 62, 76-85  
     great-winged, 261  
     grey-faced, 72, 74  
     Hawaiian, 74  
     Juan Fernandez, 62, 63, 255, 263  
     Kermadec, 62  
     mottled, 178, 194, 195, 198-203  
     northern giant, 51-53, 55-57, 60, 61, 76-85  
     Peruvian diving, 83  
     Pycroft's, 72, 172  
     southern giant, 61, 77  
     spp., 51-64, 82, 178, 194-207, 231  
     white-chinned, 52, 56, 79  
*Petrochelidon nigricans*, see martin, Australian tree  
*Petroica longipes*, see robin, North Island  
*Pezophaps solitaria*, see solitaire, Réunion  
 Phalacrocoracidae, 266-273  
*Phalacrocorax carbo*, see shag, black  
     *melanoleucos*, see shag, little  
 phalarope, grey, 257  
     red-necked, 257  
*Phalaropus fulicaria*, see phalarope, grey  
     *lobatus*, see phalarope, red-necked  
*Philesturnus carunculatus*, see saddleback, South Island  
*Phoebastria irrorata*, see albatross, waved  
 pigeon, Chatham Island, 86-88, 231  
     feral, see pigeon, rock  
     fruit spp., 86-88  
     Japanese wood, 231  
     New Zealand, 86-88, 102, 225, 230, 236, 294  
     rock, 224-233  
     spp., 86-88  
*Platalea flavipes*, see spoonbill, yellow-billed  
     *regia*, see spoonbill, royal  
*Platyspiza*, see finch, Darwin's spp.  
*Plegadis falcinellus*, see ibis, glossy  
 plover, greater sand, 257  
     grey, 257  
     lesser sand, 257, 262  
     piping, 291  
     shore, 172, 182  
     spur-winged, 262  
     Wilson's, 291  
*Pluvialis squatarola*, see plover, grey  
 pochard, Baer's, 108  
     common, 108  
     Madagascar, 108

- white-eyed, 108  
*Podiceps cristatus*, see grebe, Australasian crested  
*Poliiocephalus poliocephalus*, see grebe, hoary-headed  
*rufopectus*, see dabchick, New Zealand  
*Porphyrio hochstetteri*, see takahē  
*melanotus*, see pūkeko  
*Porzana pusilla*, see crane, marsh  
*tabuensis*, see crane, spotless  
 POWLESLAND, R., see LAMB, S.D.; TAYLOR, H.R.;  
 POWLESLAND, R.  
 prion, Antarctic, 178  
   broad-billed, 178, 194, 195, 198-203  
   fairy, 171, 195  
   fulmar, 38, 39  
*Procellaria aequinoctialis*, see petrel, white-chinned  
*pelagica*, see *Hydrobates pelagicus*  
*vittata*, see *Pachyptila vittata*  
*westlandica*, see petrel, Westland  
 Procellariid spp., 51-64  
 Procellariidae, 51-64, 194-207  
 Procellariiformes, 51-64, 72-75, 170-172, 177-181, 194-  
 207, 253, 255, 256, 261, 263  
*Prosthemadera novaeseelandiae*, see tūi  
*novaeseelandiae chathamensis*, see tūi, Chatham  
 Island  
*Prunella modularis*, see dunnoek  
*Pseudobulweria macgillivrayi*, see petrel, Fiji  
 Psittacine spp., 275  
*Pterodroma brevipes*, see petrel, collared  
*cookii*, see petrel, Cook's  
*externa*, see petrel, Juan Fernandez  
*gouldi*, see petrel, grey-faced  
*inexpectata*, see petrel, mottled  
*macroptera*, see petrel, great-winged  
*macroptera macroptera*, see petrel, great-winged  
*magentae*, see taiko, Chatham Island  
*neglecta*, see petrel, Kermadec  
*pycrofti*, see petrel, Pycroft's  
*sandwichensis*, see petrel, Hawaiian  
*Ptilinopus regina*, see dove, rose-crowned fruit  
 puffin, Atlantic, 291  
*Puffinus assimilis*, see shearwater, little  
*assimilis haurakiensis*, see shearwater, North  
 Island little  
*carneipes*, see shearwater, flesh-footed  
*gavia*, see shearwater, fluttering  
*griseus*, see shearwater, sooty  
*huttoni*, see shearwater, Hutton's  
*pacificus*, see shearwater, wedge-tailed  
*tenuirostris*, see shearwater, short-tailed  
 pūkeko, 270  
 quail, bobwhite, 262  
 QUIÑONES, J.; ROMERO, C.; ZAVALAGA, C.  
   Vessel survey observations confirm  
   wintering dispersion of northern giant-petrel  
   (*Macronectes halli*) juveniles in southern-  
   central Perú; what is their origin? 76-85  
 rail, banded, 260  
   spp., 295  
*Raphus cucullatus*, see dodo  
 raptor, North American spp., 275  
   spp., 281  
 ratite, 284  
 raven, common, 291  
   white-necked, 292  
 redhead, 108  
 redpoll, common, 211, 218, 272  
*Reinwardtoena browni*, see cuckoo-dove, pied  
*crassirostris*, see cuckoo-dove, crested  
*reinwardtii*, see cuckoo-dove, great  
*r. brevis*, 89  
*r. griseotincta*, 89  
 REXER-HUBER, K., see MATTERN, T.; REXER-  
 HUBER, K.; PARKER, G.; AMEY, J.; GREEN,  
 C-P; TENNYSON, A.J.D.; SAGAR, P.M.;  
 THOMPSON, D.R.  
 REYES, E.M.R., see WITTMER-NARANJO, C.;  
 REYES, E.M.R.; JÁCOME, H.E.T.; RUEDA, D.;  
 SEVILLA, C.; ORTIZ-CATEDRAL, L.  
*Rhipidura fuliginosa*, see fantail, New Zealand  
*fuliginosa fuliginosa*, see fantail, South Island  
 ROBERTSON, H.A., see CLOUT, M.N.; ROBERTSON,  
 H.A.  
 robin, North Island, 102  
 ROMERO, C., see QUIÑONES, J.; ROMERO, C.;  
 ZAVALAGA, C.  
 rorua, 131-146, 173-176, see also kiwi, great spotted  
 ROWE, L.K. Garden birds at Rangiora, Christchurch,  
 and Kaikōura, South Island, New Zealand:  
   results from banding 1961-2016, 208-223  
 rowi, 141, 142, see also kiwi, Ōkarito brown  
 RUEDA, D., see WITTMER-NARANJO, C.; REYES,  
 E.M.R.; JÁCOME, H.E.T.; RUEDA, D.;  
 SEVILLA, C.; ORTIZ-CATEDRAL, L.  
 ruru, see morepork  
 saddleback, South Island, 262  
 SAGAR, P.M., see MATTERN, T.; REXER-HUBER,  
 K.; PARKER, G.; AMEY, J.; GREEN,  
 C-P; TENNYSON, A.J.D.; SAGAR, P.M.;  
 THOMPSON, D.R.  
 SAGAR, P.M., see HOLDAWAY, R.N.; SAGAR, P.M.;  
 BRISKIE, J.V.  
 sanderling, 256  
 sandpiper, broad-billed, 257  
   buff-breasted, 253, 257  
   common, 257  
 SAVILLE, I., see MISKELLY, C.M.; CROSSLAND, A.C.;  
 SAVILLE, I.; SOUTHEY, I.; TENNYSON, A.J.D.;  
 BELL, E.A.  
 scaup, greater, 108  
   lesser, 108  
   New Zealand, 108-130, 266, 269-271  
*Sceloglaux albifacies*, see owl, laughing



- SCHUCKARD, R., see MELVILLE, D.S.;  
SCHUCKARD, R.
- seabird spp., 37-50, 51-64, 72-75, 76-85, 166-169, 170-172, 177-181, 183-187, 188-193, 194-207, 253-265, 274-277, 296
- sea eagle, European white-tailed, 276  
white-bellied, 261-262
- SEDDON, P.J., see WHITE, J.; MATTERN, T.;  
ELLENBERG, U.; GARCIA-BORBOROGLU, P.;  
HOUSTON, D.M.; SEDDON, P.J.; MAYS, H.L.
- SEVILLA, C., see WITTMER-NARANJO, C.; REYES,  
E.M.R.; JÁCOME, H.E.T.; RUEDA, D.;  
SEVILLA, C.; ORTIZ-CATEDRAL, L.
- shag, black, 261, 269, 271  
Bounty Island, 38  
Foveaux, 166  
little, 269, 271, see also shag, little pied  
little pied, 269, 271  
New Zealand king, 274-277  
Otago, 166-169  
spp., 167, 168, 266-273  
spotted, 167  
Stewart Island, 260, 276
- shearwater, Buller's, 72-75  
flesh-footed, 72, 74  
fluttering, 72, 74  
great, 255  
Hutton's, 261  
little, 72, 170-172  
North Island little, 170-172  
pink-footed, 255, 261, 263  
short-tailed, 53, 61  
sooty, 53, 61, 72, 178, 194-203, 207  
spp., 254  
wedge-tailed, 62, 63
- shelduck, chestnut-breasted, 254  
paradise, 266, 269-271, 280
- SHOREY, L., see BRISKIE, J.V.; SHOREY, L.
- shoveler, Australasian, 266, 269-271, see also shoveler,  
New Zealand  
New Zealand, 117, 123, 266, 269-271  
Northern, 254, 263
- silveryeye, 208-211, 214-221, 225
- skua, long-tailed, 253, 257  
South Polar, 253, 257, 263  
spp., 62  
subantarctic, 260, 262
- solitaire, Réunion, 232
- SOUTHEY, I., see MISKELLY, C.M.; CROSSLAND,  
A.C.; SAVILLE, I.; SOUTHEY, I.; TENNYSON,  
A.J.D.; BELL, E.A.
- sparrow, 212  
house, 208-211, 214, 215, 218-221
- Spheniscidae, 63  
*Spheniscus demersus*, see penguin, African  
*humboldti*, see penguin, Humboldt
- spoonbill, royal, 261  
yellow-billed, 261
- STAHL, J.C., see MISKELLY, C.M.; STAHL, J.C.;  
TENNYSON, A.J.D.; BISHOP, C.R.
- starling, common, 161-165, 208-211, 213-214, 219-221,  
272
- Stephanoaetus coronatus*, see eagle, African crowned
- Stercorarius longicaudus*, see skua, long-tailed  
spp., see jaeger spp.
- Sterna bergii*, see tern, crested  
*hirundo*, see tern, common  
*paradisaea*, see tern, Arctic  
*striata*, see tern, white-fronted
- Sternula albifrons*, see tern, little  
*nereis davisae*, see tern, fairy
- Stictocarbo punctatus*, see shag, spotted
- stilt, spp., 272
- stint, little, 257
- stitchbird, 172
- storm petrel, Australian white-faced, 253, 256  
European, 285  
grey-backed, 177-181, 195  
New Zealand, 285  
white-faced, 172
- Streptopelia capicola*, see turtle-dove, Cape  
*decaocto*, see dove, collared  
*orientalis*, see dove, eastern turtle  
*risoria*, see dove, Barbary
- Strigopidae, 234
- Strigopinae, 234
- Strigops habroptilus*, see kākāpō
- Struthio*, 283
- Sturnus vulgaris*, see starling, common
- Sula granti*, see booby, Nazca  
*leucogaster*, see booby, brown  
*neboxii*, see booby, blue-footed  
spp., 63  
*sula*, see booby, red-footed  
*sula rubripes*, see booby, red-footed (South-west  
Pacific subsp.)  
*sula websteri*, see booby, red-footed (Eastern  
Pacific subsp.)
- swallow, spp., 26  
welcome, 262
- swan, black, 269-271
- swift, *Apus* spp., 31  
chimney, 28  
fork-tailed, 259, 263  
Neotropical spp., 30-32  
spp., 26  
*Tachymarpis* spp., 31  
waterfall, 27
- swiftlet, Atiu, 29-31  
Australian, 27-30  
black-nest, 26, 28-31, 34, 35  
Germain's, 28, 29  
glossy, 28, 30, 31, 34, 35  
Halmahera, 28  
Indian, 27  
Mariana, 29

- mossy-nest, 28-30  
 mountain, 28  
 Palau, 28  
 pygmy, 28  
 Seychelles, 29, 30  
 spp., 26-36  
 three-toed, 27  
 uniform, 828-30  
 white-nest, 28-31, 34, 35  
 white-rumped, 28-30
- Tachybaptus novaehollandiae*, see grebe, Australasian little  
*Tachymarpis* swift spp., 31  
*Tadorna tadornoides*, see shelduck, chestnut-breasted  
     *variegata*, see shelduck, paradise  
 taiko, Chatham Island, 261  
 takahē, 279, 281  
 takahikare moana, see storm petrel, white-faced  
 TARBURTON, M.K.; COLLINS, C.T. The food and foraging of *Collocalia* and *Aerodramus* swiftlets: a review, 26-36  
 tattler, wandering, 257  
 tawaki, see penguin, Fiordland crested  
 TAYLOR, H.R., see LAMB, S.D.; TAYLOR, H.R.; POWLESLAND, R.  
 teal, brown, 69, 266, 294  
     chestnut, 254  
     grey, 266, 269, 271  
 TENNYSON, A.J.D., see MATTERN, T.; REXER-HUBER, K.; PARKER, G.; AMEY, J.; GREEN, C.-P.; TENNYSON, A.J.D.; SAGAR, P.M.; THOMPSON, D.R.  
 TENNYSON, A.J.D., see MISKELLY, C.M.; STAHL, J.C.; TENNYSON, A.J.D.; BISHOP, C.R.  
 TENNYSON, A.J.D., see MISKELLY, C.M.; BISHOP, C.R.; TENNYSON, A.J.D.  
 TENNYSON, A.J.D., see MISKELLY, C.M.; CROSSLAND, A.C.; SAVILLE, I.; SOUTHEY, I.; TENNYSON, A.J.D.; BELL, E.A.  
 tern, Arctic, 253, 258  
     black-fronted, 147-160, 262  
     Caspian, 262  
     common, 258, 262  
     crested, 258  
     fairy, 261, 262  
     gull-billed, 253, 258, 262, 263  
     little, 258  
     sooty, 63, 260-261  
     whiskered, 258  
     white, 260  
     white-fronted, 262  
     white-winged black, 258, 262  
*Thalassarche bulleri*, see mollymawk, Buller's  
     *bulleri bulleri*, see mollymawk, southern Buller's  
     *bulleri platei*, see mollymawk, northern Buller's  
     *carteri*, see mollymawk, Indian yellow-nosed  
     *cauta cauta*, see mollymawk, shy  
     *cauta steadi*, see mollymawk, white-capped  
     *chlororhynchos*, see mollymawk, Atlantic yellow-nosed  
     *salvini*, see mollymawk, Salvin's  
*Thinornis cucullatus*, see dotterel, hooded  
     *novaeseelandiae*, see plover, shore  
     *rossii*, see plover, shore  
     *rubricollis*, see dotterel, hooded  
 THOMPSON, D.R., see MATTERN, T.; REXER-HUBER, K.; PARKER, G.; AMEY, J.; GREEN, C.-P.; TENNYSON, A.J.D.; SAGAR, P.M.; THOMPSON, D.R.  
 THOMPSON, M.P.; MCKINLAY, B. Seasonal survey of waterfowl (Anatidae), shags (Phalacrocoracidae) and fernbird (*Bowdleria punctata*) at Te Nohoaka o Tukiauu/Sinclair Wetlands, Otago: July 2015 - July 2018, 266-273  
 THORNBURROW, D., see FITZGERALD, N.; INNES, J.; WATTS, C.; THORNBURROW, D.; BARTLAM, S.; COLLINS, K.; BYERS, D.; BURNS, B.  
*Threskiornis molucca*, see ibis, white  
 thrush, song, 209, 211, 219, 220, 225, 230  
 titi, see petrel, Cook's, or petrel, Pycroft's  
 titi wainui, see prion, fairy  
*Todiramphus sanctus*, see kingfisher, sacred  
 tokoeka, Fiordland, 141-143, see also kiwi, South Island brown  
     Haast, 137, 141, 142, see also kiwi, South Island brown  
 TOY, R.; TOY, S. Breeding ecology of a translocated population of great spotted kiwi (*Apteryx haastii*), 131-146  
 TOY, R.; TOY, S. Changes in behaviour of great spotted kiwi (*Apteryx haastii*) following handling, 173-176  
 TOY, S., see TOY, R.; TOY, S. (2 papers)  
 treeswift spp., 26  
 triller, white-winged, 262  
*Tringa hypoleucos*, see sandpiper, common  
     *incana*, see tattler, wandering  
     *nebularia*, see greenshank, common  
*Tryngites subruficollis*, see sandpiper, buff-breasted  
 tūi, 93-107, 294  
     Chatham Island, 102  
*Turdus merula merula*, see blackbird  
     *philomelos*, see thrush, song  
 turtle-dove, Cape, 87  
*Tyto alba*, see owl, barn  
*Vanellus miles*, see plover, spur-winged  
 warbler, grey, 65, 67, 209, 211, 218-220, 225  
 waterbird spp., 108-130  
 waterfowl spp., 13-25, 108-130, 266-273  
 WATTS, C., see FITZGERALD, N.; INNES, J.; WATTS, C.; THORNBURROW, D.; BARTLAM, S.; COLLINS, K.; BYERS, D.; BURNS, B.

- weka, 1-12, 140-143, 280, 281  
     western, 1-12, 136, 140, 141  
 whimbrel, 257  
 WHITE, J.; MATTERN, T.; ELLENBERG, U.;  
     GARCIA-BORBOROGLU, P.; HOUSTON,  
     D.M.; SEDDON, P.J.; MAYS, H.L. Field sexing  
     techniques for Fiordland crested penguins  
     (tawaki; *Eudyptes pachyrhynchus*), 188-193  
 WHITEHEAD, E. Little shearwaters (*Puffinus assimilis*  
     *haurakiensis*) as prey for morepork (*Ninox*  
     *novaeseelandiae*), 170-172  
 WHITEHEAD, E., see CROWE, P.; WILLIS, K.;  
     BUTLER, D.; WHITEHEAD, E.  
 WILLIAMS, M., see ADAMS, J.S.; WILLIAMS, M.  
 WILLIS, K., see CROWE, P.; WILLIS, K.; BUTLER, D.;  
     WHITEHEAD, E.  
 WITTMER-NARANJO, C.; REYES, E.M.R.;  
     JÁCOME, H.E.T.; RUEDA, D.; SEVILLA, C.;  
     ORTIZ-CATEDRAL, L. Diet of the Floreana  
     mockingbird (*Mimus trifasciatus*) during the  
     dry season on Champion and Gardner Islets,  
     Galápagos Islands, Ecuador, 245-252  
 WOODLEY OF MENIE, M.A.; PEÑAHERRERA-  
     AGUIRRE, M. Did Molly Falla observe an  
     instance of active tactical deception in the kea  
     (*Nestor notabilis*)? 290-293  
 woodpigeon, 87  
 woodswallow, masked, 259, 262  
 wren, rock, 180  
 wrybill, 149  
*Xenicus gilviventris*, see wren, rock  
 yellowhammer, 210, 211, 218, 272  
 yellowhead, 262, see also mohua  
 ZAVALAGA, C., see QUIÑONES, J.; ROMERO, C.;  
     ZAVALAGA, C.  
*Zenaida asiatica*, see dove, white-winged  
     *galapagoensis*, see dove, Galápagos  
     *macroura*, see dove, mourning  
*Zosterops lateralis lateralis*, see silvereye  
 Compiled by TIM LOVEGROVE, 6 Deuxberry Ave,  
     Northcote, Auckland 0627